

Holographic Space-Time, Fractional Quantum Hall Effect, Artificial Superlattices, Topological Strings And N=2 Gauge Systems, Black Holes In Higher Spin Gravity, Matrix Quantum Mechanics, Chiral Symmetry Breaking, Anomalous Transport Phenomena, String Theory And Unification, Attention And Intention, Condition Of Phase-Conjugate-Adaptive-Resonance Is Necessary To Completely Specify The Act Of Perception, Hē Phýsis Oudèn Poieî Hálmata: Natura Non Facit Saltus:"Nature Does Not Make [Sudden] Jumps.":A Principle Of Natural Philosophies Since Aristotle's Time, The Exact Phrase Coming From Carl Von Linné : Γόρδιος Δεσμός Górdios Desmós:"Gordian Knot" Models:

[illegible]

¹**Dr. K.N.P.Kumar**

Post Doctoral Scholar, Department of Mathematics, Kuvempu University, Janna Sahyadri, Shankaraghatta, Karnataka, India

²Prof. C.J.Prabhakara

Chairman and Professor Department of Computer Science, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka,
India

²Prof. S.K. Narasimha Murthy

Chairman and Professor Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

²Dr. B.J.Gireesha

Assistant Professor Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

[illegible]

Abstract: Aficionado buff, devotee dilettante, adept aesthete, appreciator arbiter, cognoscente critic have tried to establish relationship betwixt holographic spacetimes and Fraction Quantum Hall Effect. We have given models for some of the studies done in the dovetailing concomitant mathematical modular enucleation. Some notable examples are as follows: **Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin** develop a systematic method for renormalizing the AdS/CFT prescription for computing correlation functions. This involves regularizing the bulk on-shell supergravity action in a covariant way, computing all divergences, adding counterterms to cancel them and then removing the regulator. We explicitly work out the case of pure gravity up to six dimensions and of gravity

coupled to scalars. The method can also be viewed and visualized as providing a holographic reconstruction of the bulk spacetime metric and of bulk fields on this spacetime, out of conformal field theory data. Knowing which sources are turned on is sufficient in order to obtain an asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order so that all infrared divergences of the on-shell action are obtained. To continue the holographic reconstruction of the bulk fields one needs new CFT data: the expectation value of the dual operator. In particular, in order to obtain the bulk metric one needs to know the expectation value of stress-energy tensor of the boundary theory. We provide completely explicit formulae for the holographic stress-energy tensors up to six dimensions. We show that both the gravitational and matter conformal anomalies of the boundary theory are correctly reproduced. We also obtain the conformal transformation properties of the boundary stress-energy tensors.

Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin. The theory of holographic space-time (HST) generalizes both string theory and quantum field theory. It provides a geometric rationale for supersymmetry (SUSY) and a formalism in which **super-Poincare invariance follows from Poincare invariance**. HST unifies particles and black holes, realizing both as excitations of non-commutative geometrical variables on a holographic screen. Compact extra dimensions are interpreted as finite dimensional unitary representations of super-algebras, and have no moduli. Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of super-Poincare invariant limits in which the number of holographic degrees of freedom goes to infinity. Finite radius de Sitter (dS) spaces have no moduli, and break SUSY with a gravitino mass scaling like $\Lambda^{1/4}$.

T. Banks presents a holographic theory of inflation and fluctuations. The **inflaton field** is an emergent concept, describing the geometry of an underlying HST model, rather than "a field associated with a microscopic string theory". **T. Banks** argues that the phrase in quotes is meaningless in the HST formalism. Cite as: arXiv: 1109.2435 [hep-th] **Holographic Space-Time: The Takeaway T. Banks. T. Banks, W. Fischler** use the formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain a firewall, i.e. an in-falling observer encounters highly excited states at a time much shorter than the light crossing time of the Schwarzschild radius. This conclusion is much less dramatic in HST than in the hypothetical models of quantum gravity used in [1]. In HST there is no dramatic change in particle physics inside the horizon until a time of order the Schwarzschild radius. Report number: UTTG-15-12; TCC-015-12; RUNHETC-2012-17; SCIPP 12/11 arXiv: 1208.4757 [hep-th] **Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler. Tom Banks** reviews the holographic theory of space-time and its applications to cosmology. Much of this has appeared before, but this discussion is more unified and concise. He also includes some material on work in progress, whose aim is to understand compactification in terms of finite-dimensional super-algebras. **Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era .**The spinorial geometry method of solving Killing spinor equations is reviewed as it applies to six-dimensional (1,0) supergravity. In particular, it is explained how the method is used to identify both the fractions of supersymmetry preserved by and the geometry of all supersymmetric backgrounds. Then two applications are described to systems that exhibit superconformal symmetry. The first is the proof that some six-dimensional black hole horizons are locally isometric to $\text{AdS}_3 \times \Sigma_3$, where Σ_3 is diffeomorphic to S^3 . The second one is a description of all supersymmetric solutions of six-dimensional (1,0) superconformal theories and in particular of their brane solitons. **M Akyol and G Papadopoulos 2014 Class. Quantum Grav 31 123001 doi:10.1088/0264-9381/31/12/123001 Spinorial geometry, horizons and superconformal symmetry in six dimensions.** Field of squeezed states for gravitational-wave (GW) detector enhancement is rapidly maturing. In this review paper, **S S Y Chua¹, B J J Slagmolen, D A Shaddock and D E McClelland** provide an analysis of the field circa 2013. They begin by outlining the concept and description of quantum squeezed states. This is followed by an overview of how quantum squeezed states can improve GW detection, and the requirements on squeezed states to achieve such enhancement. Next, an overview of current technology for producing squeezed states, using atoms, optomechanical methods and nonlinear crystals, is provided. We finally highlight the milestone squeezing implementation experiments at the GEO600 and LIGO GW detectors. **S S Y Chua et al 2014 Class. Quantum Grav 31 183001 doi:10.1088/0264-9381/31/18/183001 Quantum squeezed light in gravitational-wave detectors. Tom Banks and John Kehayias** present a new framework for

defining fuzzy approximations to geometry in terms of a cutoff on the spectrum of the Dirac operator, and a generalization of it that we call the Dirac-flux operator. This framework does not require a symplectic form on the manifold, and is completely rotation invariant on an arbitrary n -sphere. The framework is motivated by the formalism of holographic space-time, whose fundamental variables are sections of the spinor bundle over a compact Euclidean manifold. The strong holographic principle requires the space of these sections to be finite dimensional. They discuss applications of fuzzy spinor geometry to holographic space-time and to matrix theory. DOI: <http://dx.doi.org/10.1103/PhysRevD.84.086008> © 2011 American Physical Society **Fuzzy geometry via the spinor bundle, with applications to holographic space-time and matrix theory Phys. Rev. D 84, 086008 – Published 25 October 2011 Tom Banks and John Kehayias. Chaolun Wu, Shao-Feng Wu** show that Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry. The key to this formalism is a set of correspondence relations that maps all the field degrees of freedom in the Horava-Lifshitz gravity theory to external background (source) fields among others in the effective action of the quantum Hall effect, according to their symmetry transformation properties. **Chaolun Wu, Shao-Feng Wu** originally derive the **map as a holographic dictionary**, but its form is independent of the existence of holographic duality. This paves the way for the application of Horava-Lifshitz holography on fractional quantum Hall effect. Using the simplest holographic Chern-Simons model, we compute the low energy effective action at leading orders and show that it captures universal electromagnetic and geometric properties of Quantum Hall States, including the Wen-Zee shift, Hall viscosity, angular momentum density and their relations. They identify the shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum. This enables us to distinguish guiding center angular momentum density from the internal one, which is the sum of Landau orbit spin and intrinsic (topological) spin of the composite particles. Effective action shows that Hall viscosity is minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity is half of the guiding center angular momentum density. ArXiv: 1409.1178 [hep-th] **Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect Chaolun Wu, Shao-Feng Wu. Brian Swingle** shows how recent progress in real space renormalization group methods can be used to define a generalized notion of holography inspired by **holographic dualities in quantum gravity**. The generalization is based upon organizing information in a quantum state in terms of scale and defining a higher dimensional geometry from this structure. While states with a finite correlation length typically give simple geometries, the state at a quantum critical point gives a discrete version of anti de Sitter space. Some finite temperature quantum states include black hole-like objects. The gross features of equal time correlation functions are also reproduced in this geometric framework. The relationship between this framework and better understood versions of holography is discussed: Phys. Rev. D 86, 065007 (2012) DOI: 10.1103/PhysRevD.86.065007 arXiv: 0905.1317 [cond-mat.str-el] **Entanglement Renormalization and Holography Brian Swingle. Paul Fendley, 1 Matthew P. A. Fisher² and Chetan Nayak^{3,4}** study the entropy of chiral 2+1-dimensional topological phases, where there are both gapped bulk excitations and gapless edge modes. They show how the entanglement entropy of both types of excitations can be encoded in a single partition function. This partition function is holographic because it can be expressed entirely in terms of the conformal field theory describing the edge modes. They give a general expression for the holographic partition function, and discuss several examples in depth, including abelian and non-abelian fractional Quantum Hall States, and $p + ip$ superconductors. They extend these results to include a point contact allowing tunneling between two points on the edge, which causes thermodynamic entropy associated with the point contact to be lost with decreasing temperature. Such a perturbation effectively breaks the system in two, and we can identify the thermodynamic entropy loss with the loss of the edge entanglement entropy. From these results, we obtain a simple interpretation of the non-integer ‘ground state degeneracy’ which is obtained in 1+1-dimensional quantum impurity problems: its logarithm is a 2+1-dimensional topological entanglement entropy **Topological Entanglement Entropy from the Holographic Partition Function Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}. John Preskill** considers some promising future directions for quantum information theory that could influence the development of 21st century physics. Advances in the theory of the distinguishability of superoperators may lead to new strategies for improving

the precision of quantum-limited measurements. A better grasp of the properties of multi-partite quantum entanglement may lead to deeper understanding of strongly-coupled dynamics in quantum many-body systems, quantum field theory, and quantum gravity. With the discovery of an apparent separation between the classical and quantum classifications of computational complexity [1], and of fault-tolerant schemes for quantum computation [2], quantum information theory has earned a lasting and prominent place at the foundations of computer science. But at present this discipline seems rather isolated from most of the rest of physics. Will this change in the future? How might it change?

One view is that thinking about information theory will lead us to a deeper understanding of the foundations of quantum mechanics. This vision has been vividly expressed by John Wheeler [3]; Bill Wootters [9] and Chris Fuchs [5] have been among its particularly eloquent spokespersons. But I am not convinced in my heart that we are supposed to understand the foundations of quantum mechanics much better than we currently do. **John Preskill** prefers to look in a different direction to anticipate where quantum information may have an impact on physics. Our deepening understanding of quantum information may lead to new strategies for pushing back the boundaries of quantum-limited measurements. Quantum entanglement, quantum error correction, and quantum information processing might all be exploited to improve the information-gathering capability of physics experiments.

The most challenging and interesting problems in quantum dynamics involve understanding the behavior of strongly-coupled many-body systems — systems with many degrees of freedom that undergo large quantum fluctuations. Better ways of characterizing and classifying the features of many particle entanglements may lead to new and more effective methods for understanding the dynamical behavior of complex quantum system. A watchword of quantum information theory is: “Entanglement is a Useful Resource.” It should not be a surprise if entanglement can extend the capabilities of the laboratory physicist. For example, the phenomenon of superdense coding illustrates that shared entanglement can enhance classical communication between two parties [13]. The same strategy can sometimes be used to exploit entanglement to improve the distinguishability among Hamiltonians (an idea suggested by Chris Fuchs [14]). Suppose I wish to observe the precession of spin-1/2 objects to determine the value of an unknown magnetic field. If two spins are available, one way to estimate the value of the unknown field is to allow both spins to precess in the field independently, and then measure them separately. An alternative method is to prepare an entangled Bell pair, expose one of the two spins to the magnetic field while the other is carefully shielded from the field, and finally carry out a collective Bell measurement on the pair. It turns out that in many cases (for example when we have no a priori knowledge about the field direction), the entangled strategy extracts more information about the unknown field than the strategy in which uncorrelated spins are measured one at a time [6]. This separation still holds even if we allow the unentangled strategy to be adaptive; that is, even if the outcome of the measurement of the first spin is permitted to influence the choice of the measurement that is performed on the second spin. **Quantum information and physics: some future directions John Preskill.**

Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q can be deformed into compressible phases by modifying their Hamiltonian, H , by a chemical potential $H \rightarrow H - \mu Q$. We study 2+1 dimensional CFTs upon which an explicit S duality mapping can be performed. **Subir Sachdev** finds that this construction leads naturally to compressible phases which are superfluids, solids, or non-Fermi liquids which are more appropriately called ‘Bose metals’ in the present context. The Bose metal preserves all symmetries and has Fermi surfaces of gauge-charged fermions, even in cases where the parent CFT can be expressed solely by bosonic degrees of freedom. Monopole operators are identified as order parameters of the solids, and the product of their magnetic charge and Q determines the area of the unit cell. He presents implications for holographic theories on asymptotically AdS_4 spacetimes: S duality and monopole/dyon fields play important roles in this connection **Compressible quantum phases from conformal field theories in 2+1 dimensions Subir Sachdev. Andrei T. Patrascu** extends the notion of quantization from the standard interpretation focused on non-commuting observables defined starting from classical analogues, to the topological equivalents defined in terms of coefficient groups in (co)homology. It is shown that the commutation relations between quantum observables become (non)compatibility relations between coefficient groups. Main result is the construction of a new, higher-level form of quantization, as seen from the perspective of the universal coefficient theorem. This idea brings us closer to a consistent quantization of gravity, allows for a systematic description of topology changing string interactions but also gives new, quantum-topological degrees of

freedom in discussions involving quantum information. On the practical side, a possible connection to the fractional quantum Hall effect is explored. ArXiv:1411.4475 [physics.gen-ph] **The quantization of topology, from quantum Hall effect to quantum gravity Andrei T. Patrascu. Robbert Dijkgraafa Cumrun Vafa** show that B-model topological strings on local Calabi–Yau threefolds are large-N duals of matrix models, which in the planar limit naturally give rise to special geometry. These matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by deforming $\mathcal{N} = 2$ theories by a superpotential term that can be directly identified with the potential of the matrix model. Moreover by tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models coupled to 2d gravity, thereby relating non-critical string theories to type II superstrings on Calabi–Yau backgrounds. Copyright © 2002 Published by Elsevier B.V. **Nuclear Physics B Volume 644, Issues 1–2, 11 November 2002, Pages 3–20 Matrix models, topological strings, and supersymmetric gauge theories Robbert Dijkgraafa Cumrun Vafa doi:10.1016/S0550-3213(02)00766-6.** The possible domain structures which can arise in the universe in a spontaneously broken gauge theory are studied. It is shown that the formation of domain wall, strings or monopoles depends on the homotopy groups of the manifold of degenerate vacua. The subsequent evolution of these structures is investigated. It is argued that while theories generating domain walls can probably be eliminated (because of their unacceptable gravitational effects), a cosmic network of strings may well have been formed and may have had important cosmological effects. **T W B Kibble 1976 J. Phys. A: Math. Gen. 9 1387 doi:10.1088/0305-4470/9/8/029 Topology of cosmic domains and strings.** It is an old speculation and prognostication in physics that, once the gravitational field is successfully quantized, it should serve as the natural regulator of infrared and ultraviolet singularities that plague quantum field theories in a background metric. **T Thiemann** demonstrates that at least part of this idea is implemented in a precise sense within the framework of four-dimensional canonical Lorentzian quantum gravity in the continuum. Specifically, he shows that the Hamiltonian of the standard model supports a representation in which finite linear combinations of Wilson loop functionals around closed loops, as well as along open lines with fermionic and Higgs field insertions at the end points, are densely defined operators. This Hamiltonian, surprisingly, does not suffer from any singularities; it is completely finite without renormalization. This property is shared by string theory. In contrast to string theory, however, we are dealing with a particular phase of the standard model coupled to gravity which is entirely non-perturbatively defined and second quantized. Of course, to show that the entire theory is finite requires more: one would need to know what the physical observables are, apart from the Hamiltonian constraint, and whether they are also finite. However, with the results given in this paper this question can now be answered, at least in principle. **T Thiemann 1998 Class Quantum Grav.15 1281 doi:10.1088/0264-9381/15/5/012 Quantum spin dynamics (QSD): V. Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories. T Thiemann and O Winkler** apply the methods outlined in the previous paper of this series to the particular set of states obtained by choosing the complexifier to be a Laplace operator for each edge of a graph. The corresponding coherent state transform was introduced by Hall for one edge and generalized by Ashtekar, Lewandowski, Marolf, Mourão and Thiemann to arbitrary, finite, piecewise-analytic graphs. However, both of these works were incomplete with respect to the following two issues. The focus was on the unitarity of the transform and left the properties of the corresponding coherent states themselves untouched. While these states depend in some sense on **complexified connections**, it remained unclear what the complexification was in terms of the coordinates of the underlying real phase space. In this paper we complement these results: first, we explicitly derive the complexification of the configuration space underlying these heat kernel coherent states and, secondly, prove that this family of states satisfies all the usual properties. (i) **Peakedness in the configuration, momentum and phase space (or Bargmann-Segal) representation.** (ii) **Saturation of the unquenched Heisenberg uncertainty bound.** (iii) (Over) completeness. These states therefore comprise a candidate family for the semiclassical analysis of canonical quantum gravity and quantum gauge theory coupled to quantum gravity. They also enable error-controlled approximations to difficult analytical calculations and therefore set a new starting point for numerical, semiclassical canonical quantum general relativity and gauge theory. The text is supplemented and accentuated with an appendix which contains extensive graphics in order to give a feeling for the so far unknown peakedness properties of the states constructed. **T Thiemann and O Winkler 2001 Class Quantum Grav.18 2561 doi:10.1088/0264-9381/18/14/301 Gauge field theory coherent states**

(GCS): II. Peakedness properties. Kevin H. Knuth^{1,a} and Newshaw Bahreyni¹ present a novel derivation of both the Minkowski metric and Lorentz transformations from the consistent quantification of a causally ordered set of events with respect to an **embedded observer**. Unlike past derivations, which have relied on assumptions such as the existence of a 4-dimensional manifold, symmetries of space-time, or the constant speed of light, **Kevin H. Knuth^{1,a} and Newshaw Bahreyni¹** demonstrate that this now familiar mathematics can be derived as the unique means to consistently quantify a network of events. This suggests that **space-time need not be physical, but instead the mathematics of space and time emerges as the unique way in which an observer can consistently quantify events and their relationships to one another. The result is a potential foundation for emergent space-time. A potential foundation for emergent space-time Kevin H. Knuth^{1,a} and Newshaw Bahreyni¹.** In this topical review, **Alejandro Perez** reviews the present status of the **spin foam formulation** of non-perturbative (background-independent) quantum gravity. The topical review is divided into two parts. In the first part, **Alejandro Perez** presents a general introduction to the main ideas emphasizing their motivation from various perspectives. Riemannian three-dimensional gravity is used as a simple example to illustrate conceptual issues and the main goals of the approach. The main features of the various existing models for four-dimensional gravity are also presented here. We conclude with a discussion of important questions to be addressed in four dimensions (gauge invariance, discretization independence, etc). In the second part, Alejandro concentrates on the definition of the **Barrett–Crane model**. He presents the main results obtained in this framework from a critical perspective. Finally, we review the combinatorial formulation of spin foam models based on the dual group field theory technology. We present the Barrett–Crane model in this framework and review the finiteness results obtained for both its Riemannian and its Lorentzian variants. **Alejandro Perez 2003 Class. Quantum Grav. 20 R43 doi:10.1088/0264-9381/20/6/202 Spin foam models for quantum gravity.** Intrinsic relationship between topological strings and quantum hall effect is studied by many authors. Conformal symmetry, splitting strategy, symmetry in QHE,D branes, is some of the areas in which there has been some progress both in thematic and discursive form. Specific areas of metals and compounds have been explored which we shall not deliberate upon, inconsideration to spatial restraints. Some of them are as follows .Free planar electrons in a uniform magnetic field are shown to possess the dynamical symmetry of area-preserving diffeomorphisms (W-infinity algebra). Intuitively, this is a consequence of gauge invariance, which forces dynamics to depend only on the flux. The infinity of generators of this symmetry act within each Landau level, which is infinite dimensional in the thermodynamic limit. The incompressible ground states corresponding to completely filled Landau levels (integer quantum Hall effect) possess a dynamical symmetry, since they are left invariant by an infinite subset of generators. This geometrical characterization of incompressibility also holds for fractional fillings of the lowest level (simplest fractional Hall Effect) in the presence of Haldane's effective two-body interactions. Although these modify the symmetry algebra, the corresponding incompressible ground states proposed by Laughlin are again symmetric with respect to the modified infinite algebra. **Nuclear Physics B Volume 396, Issues 2–3, 17 May 1993, Pages 465–490 Infinite symmetry in the quantum Hall effect Andrea Cappelli¹, Carlo A. Trugenberger, Guillermo R. Zemba doi:10.1016/0550-3213(93)90660-H.** In a solid, electrons behave differently than in a vacuum. In particular, their charge can break up into fractions of the elementary charge. Theoretical work shows how the electron's spin could help to observe fractional charges directly. An electron in a vacuum seems to be an indivisible particle of charge $-e$ and spin $1/2$, but an electron moving in the active background of a solid can break up into excitations ('quasiparticles') that carry modified values of charge or spin. The most celebrated example is the fractional quantum Hall effect, recognized by the 1998 Nobel Prize: the combination of an applied magnetic field and the Coulomb repulsion between electrons confined to a plane gives a liquid ground state whose quasiparticles carry fractional charge. **Nature Physics 4, 270 - 271 (2008) doi: 10.1038/nphys925 Subject Categories: Quantum physics | Condensed-matter physics Joel Moore Topological order: How spin splits the electron.** The low-lying excitations of a Quantum Hall State on disk geometry are edge excitations. Their dynamics is governed by a conformal field theory on the cylinder defined by the disk boundary and the time variable. We give a simple and detailed derivation of this conformal field theory for integer filling, starting from the microscopic dynamics of $(2 + 1)$ -dimensional non-relativistic electrons in Landau levels. This construction can be generalized to describe Laughlin's fractional Hall states via chiral bosonization, thereby making contact with the effective Chern-Simons theory approach. The conformal field theory dictates the finite-size effects

in the energy spectrum. An experimental or numerical verification of these universal effects would provide a further confirmation of Laughlin's theory of incompressible quantum fluids. **Conformal symmetry and universal properties of Quantum Hall States** Andrea Cappellia, 1, Gerald V. Dunne, Carlo A. Trugenberger, Guillermo R. Zambaa Received 20 November 1992, Accepted 10 February 1993, Available online 25 October 2002 doi:10.1016/0550-3213(93)90603-M Nuclear Physics B Volume 398, Issue 3, 21 June 1993, Pages 531–567. Steven S. Gubser and Mukund Rangamani study the recently proposed D-brane configuration [1] modeling the quantum Hall effect, focusing on the nature of the interactions between the charged particles. Our analysis indicates that the interaction is repulsive, which it should be for the ground state of the system to behave as a quantum Hall liquid. The strength of interactions varies inversely with the filling fraction, leading us to conclude that a Wigner crystal is the ground state at small ν . For larger rational ν (still less than unity), it is reasonable to expect a fractional quantum Hall ground state. **Steven S. Gubser and Mukund Rangamani JHEP05 (2001)041 doi:10.1088/1126-6708/2001/05/041 D-brane dynamics and the quantum Hall Effect** A novel hierarchy of the one-dimensional SU(N) electron models with $1/r^2$ interaction is proposed and solved by the asymptotic Bethe ansatz both for the continuum and lattice cases. The construction of the hierarchy is closely related to that for the fractional quantum Hall effect (FQHE) of the filling factor $\nu = 1/[p_1 - 1/(p_2 - \dots - 1/p_N)]$. Under the chiral constraint the model describes the essential properties of the edge states for the FQHE with the above filling fraction. Furthermore the matrix deduced from the excitation spectrum characterizes the topological order of the FQHE state. DOI: <http://dx.doi.org/10.1103/PhysRevLett.71.275> © 1993 The American Physical Society **Novel hierarchy of the SU(N) electron models and edge states of fractional quantum Hall effect Phys. Rev. Lett. 71, 275 – Published 12 July 1993 Norio Kawakami**. Generality abstract principle, principle sweeping statement, and universality induction, conclusion conjecture, and deduction generalization, judgment logical reasoning, ratiocination rationalization, cause criterion, exigency formula, precept principium, proposal proposition, rule source identification goes in to grand limbo of oblivion and hibernation when it comes to the final statement about the nature's general ledger, insofar as the transactional accountability or the heads thereof. In the following we give a monolith model for the following modules are incorporated and concomitant properties of stability, Solutional behaviour, and asymptotic analysis made. Asymptotic Symmetry And Three-Dimensional Higher Spin Ads Gravity, String Theory, Black Holes And Asymptotic Symmetries, $SL(3) \times SL(3)$ Chern-Simons Theory And Its Asymptotic Symmetry Algebra, **Band Offsets In ZnSe-ZnSs1-X Strained-Layer Superlattices**, Quantum Spin Dynamics (QSD), Gauge Field Theory Coherent States (GCS), Canonical Commutation And Adjointness Relations Of The Quantum Field Algebra Of Diffeomorphism Invariant Gauge Field Theories By Ashtekar, Lewandowski, Marolf, Mourão And Thiemann., Renormalization In The Ads/CFT Correspondence, Ads/CFT Correspondence, Holographic Reconstruction Of Spacetime, Holographic Space-Time: Big Bang To The De Sitter Era, Holographic Space-Time Does Not Predict Firewalls, Factor-Ordering Singularities Free Operator, Spinorial Geometry, Horizons And Superconformal Symmetry, Holographic Space-Time And Matrix Theory, Fuzzy Spinor Geometry And Holographic Space-Time, Horava-Lifshitz Gravity, Non-Perturbative Lorentzian Quantum Gravity, Non-Perturbative Lorentzian Quantum Gravity, Entanglement Renormalization Vis A Vis Holographic Principle, Topological Entanglement Entropy, Thermodynamic Entropy Loss With The Loss Of The Edge Entanglement Entropy, **Spin** Quantum Information, **Outcome Of The Measurement Of The First Spin Is Permitted To Influence The Choice Of The Measurement That Is Performed On The Second Spin**, Compressible Quantum Phases, Quantum Gravity As The Natural Regulator Of The Hamiltonian Constraint, Hamiltonian Constraint Of Matter Quantum Field Theories, Gauge Field Theory Coherent States, Emergent Space-Time, **Spin Foam Formulation** Of Non-Perturbative (Background-Independent) Quantum Gravity, **Signature To Be Euclidean**, Topological Strings, And Supersymmetric Gauge Theories, Topology Of Cosmic Domains And Strings, **Condition Of Phase-Conjugate-Adaptive-Resonance Is Necessary To Completely Specify The Act Of Perception**, **Non-Locality, Near And Far, Attention And Intention**, **Suppression By Cultural Conditioning In Childhood And Subsequent Lack Of Practice Cause The Natural Ability For Conscious, Intuitive Perceptions To Atrophy**, Easy Quantum Physics, Entanglement Witnesses, Chern-Simons-Landau-Ginzburg Theory, Chiral Viscoelastic Response, Gravity Dual Of A Quantum Hall Plateau Transition.

@@@@
 @@@@
 @@@@

Eighth Series Of Structure Levi Strauss Has Also Noticed That Signs Always Offer An Excess. The System Of Language, ‘The Order Of The Known’ Exceeds Actual Speech, Even Attempts At Totalization (48). Laws Pre-Exist Actual Cases. [So We Were Getting Close To A Role For Social Life, But Then It Gets **Metaphysical Again**]. As LS Put It, The Universe Signified Long Before Human Beings Knew What It Was Signifying. By Contrast, The Domination Of Nature Proceeds Partially And Progressively, Step By Step Unlike Social Life Where All Its Goals And Possibilities Given At Once. We’re Back With Two Series, This Time Conceived As Rhythms, Social And Natural. Both Technocrats And Dictators Attempt A False Synthesis Of These Two Rhythms. Levi Strauss Referred To ‘The Floating Signifier’ As A Creative Force And Deleuze Wants To Say It’s ‘The Promise Of All Revolutions’ (49). There Are Also ‘Floated Signifieds’, Which Seem To Be Possibilities Which Have Not Yet Been Realized. These Can Fill The Gap Between Signifier And Signified [And Are Found In Common Sense Expressions Like ‘Gadgets’ Or ‘Whatnot’ – Maybe Connected To The Idea Of A Bricoleur?]. It Implies A Symbolic Content, But Does Not Attempt To Fill It With Specifics. Together, These Possibilities Constitute A Structure, Two Heterogeneous Series, One Signifying, One Signified, Interdependent, And Including Particular Events, Singularities, Emitted By A Differentiator. The Singularities Belong To Neither Series Exclusively And Thus Have **No Coherent Identity** –Each Is An Excess In One Series And A Lack In The Other. The Singularities Can React Back On The Series, So Structures And Events Are Interdependent [So Structures Need A Dynamic Element, And, D Argues, An Excess, An Empty Square Instead Of Total Systematic Closure. Addresses The Old Issue Of The Static Nature Of Structuralism]. The Signifying Series Contains A Series Of Ideal Events, An Internal History. Differentiators Articulate Series

And This Produces A ‘Tangled Tale’ Overall (51). Sense Can Be Found In Either Series. It Is Not Just Signification But The Relation That Produces Signifier And Signified, [The Operation Of The Whole Structure In This Expanded Sense]. Attributable And Ascribable To Constraints On Space Accomodation Model Is Provided In Some Paper, Notwithstanding The Generality And Commonalty Of The Observation And Its Concatenatability With The Other Modules

Ninth Series Of The Problematic: [We Start To Develop The Terminology Of The More General Complexity Theory Approach]. Ideal Events Are Singularities: ‘Turning Points And Points Of Inflection; Bottlenecks, Knots, Foyers And Centres; Points Of Fusion, Condensation And Boiling; Point Of Tears And Joy, Sickness And Health, Hope And Anxiety, “Sensitive” Points. Such Singularities, However, Should Not Be Confused Either With The Personality Of The One Expressing Herself In Discourse, Or With The Individuality Of The State Of Affairs’ Designated By The Proposition, Or Even With The Generality All Universality Of The Concept... The Singularity Belongs To Another Dimension Than That Of Denotation, Manifestation All Significations. It Is Essentially Pre-Individual, Non Personal, And Aconceptual. It Is Quite Indifferent To The Individual And Collective, The Personal And The Impersonal, The Particular And The General... Singularity Is Neutral’ (52). Singularities Produce A Series Stretching Between Them, And Are Themselves Organised In A Structure. ‘The Moment That The Two Series Resonate And Communicate, We Pass From One Distribution To Another’ (53). Singularities Are Displayed In Paradoxes, Themselves Produced By A ‘Paradoxical Agent’ (53). The Events Are Ideal, But They Are Realized Or Actualised In Imperfect States Of Affairs. ‘The Distinction Is Between Event And Accident’ (53) [So Empirical Events Are Entirely Accidental]. It Is A Mistake To See Events As Exhibiting Essences, When They Are ‘Jets Of Singularities’, And To Confuse Events And Accidents, Which Is [Naive] Empiricism. Events Exist In Unlimited Time, ‘Aion, The Infinitive’ (53). Events Set Problematics Which Define Problems And Conditions [So We Have This Isomorphism Between Reality And The Activities Of Mathematicians And Philosophers Mentioned In Delanda]. Events Set The Problematic, While Specific Problems Appear As Singular Points [Which Might Have To Be Further Specified In Detail, By Adding

Specific Values And So On]. Therefore Solutions Arise By Finding The Conditions Which Determine Problems, The Singularities. Problematics Are Neither Subjective Nor Empirical. Problems Can Be Concealed In Solutions, However [See Delanda Again On The Limits Of Attempting Always To Find Solutions To Rather Than Defined Problems]: Solutions Have No Sense If They Do Not Recover This Deeper Structure Of The Origins Of Problems, The ‘Indispensable Horizon Of [What] Occurs Or Appears’ (54). Solutions Certainly Do Not Exhaust Problems, And Answers Do Not Exhaust Questions: This Alludes To Some ‘Ideational Objectivities’ (56 Sic). These Are Sometimes Alluded To By Esoteric Words, And Are Represented In General By The Empty Space, The Blank Word. Mathematics Is Not An Activity Found Exclusively In Human Consciousness, And Its Solutions Should Be Seen As Human Events Which Exposed The Conditions Of A [Real] Problem. Carroll Shows This With His Recreational Mathematics (55) [Which Often Seems To Anthropomorphise Mathematical Constructions—Deleuze Says That Conventional Conceptions Of Human Beings Also Anthropomorphise Their ‘Prepersonal Singularities’ (55)]. Human Feelings ‘Are Constituted In The Vicinity Of These Singularities: Sensitive Crisis Points, Turning Points, Boiling Points, Knots And Foyers’ (55). Events Are Only Known In The Context Of The Problem They Are Determining, And We Need A Language To Describe Events In General In Their Field, And How They Are Realized. Paradox Can Be Seen As A Particular Problem Related To Singular Points, But Again Some Empty Square Or ‘Aleatory Point’ Must Be Involved, Enabling Events To Communicate In An Unusual Way. Paradox Therefore Illustrates The Relation Of Events: It Is ‘The Unique Event, In Which All Events Communicate And Are Distributed’ (56). Paradox Alludes To This ‘Singular Being’, Corresponding To ‘The Question As Such’ (57). Alexander I. Stingl's Blog A Nomadic Scholarship Entity NOTES ON: Deleuze, G (1990) The Logic Of Sense, Trans Mark Lester, Edited By Constantin Boundas, New York: Columbia University Press For Models Kindly See One Of The Papers In The Series. Attributable And Ascribable To Constraints On Space Accommodation Model Is Provided In Some Paper, Notwithstanding The Generality And Commonalty Of The Observation And Its Concatenability With The Other Modules

In Sum, Whitehead Offers A Relational Process Ontology That Promises To Deepen The Constructivist Insights Associated With The Turn To Textuality, But Without Reducing The Universe To “Discourse” And “Materiality”. In This Ontology, Things (Whether Occasions Or Assemblages) Are Definable As Their Relevance To Other Things And In Terms Of The Way Other Things Are Relevant To Them. Things Have Relational Essences. Likewise, Things Do Not Exist Independently Of Temporality But Are Constituted By The History Of Their Specific And Situated Encounters. Every Actual Thing Is Thus “Something By Reason Of Its Activity” (Whitehead, 1927/1985, P. 26). Importantly, This Talk Of “Things” Need Not Incline One Towards Denying The Relevance Of Subjectivity. I Have Thus Taken Issue With A Tendency Illustrated In The Work Of Thrift (2008), Who Appears To Define Cutting-Edge Social Theory As Concerned With “Flow” And “Play” Rather Than With Stability (Since “Non-Foundational Theory Takes The Leitmotif Of Movement”, P. 5 And “Privileges Play”, P. 7); As “A Means Of Going Beyond Constructivism” (P. 5); And As “Trading” In “Modes Of Perception Which Are Not Subject-Based” (P. 7). I Have Tried To Show That, In Fact, The Concept Of Process Is As Much About Stability As About Change. Stability Is To Be Thought Of As An Achievement Resulting From Particular Ways Of Actualizing Potential And Of Patterning Occasions Into Spatial And Temporal Co-Assemblies. Nevertheless, I Have Also Stressed That Becoming Is An Inherently Self-Creative Process, Albeit A Self-Creation Grounded In The Facticity Of A Concrete Inheritance. Whitehead's Category Of Subjective Unity Is Thus An Exemplary Statement Of Constructivism, Which States That: “Self-Realization Is The Ultimate Fact Of Facts. An Actuality Is Self-Realizing, And Whatever Is Self-Realizing Is An Actuality” (Whitehead, 1927–1928/1985, P. 222). The “Capture Of Intensity” And The “Clutch At Vivid Immediacy” Are Thus The Defining Characteristics Of Life (Whitehead, 1927–1928/1985, P. 105). Finally, I Have Suggested That It Is Only On The Basis Of A Deep Extension Of The Concept Of Experience Throughout Nature That Whitehead Is Able To Resoundingly Affirm His Reformed Notion Of The Subjectivist Principle: “That Apart From The Experiences Of Subjects There Is Nothing, Nothing, Nothing, Bare Nothingness” (1927–1928/1985, P. 167). “Scientific Reasoning Is Completely Dominated By The Presupposition That Mental Functionings Are Not Properly Part Of Nature” (Whitehead, 1938/1966, P. 156). (Ibid)

Lewis Carroll Is Wonderful In Exploring Some Of The Paradoxes Of Logic. Following Some Of These Paradoxes Will Lead To The Important Role Of ‘Sense’ In Understanding. [NB Weird Titles Are Deleuze's Own]. **First Series Of Paradoxes Of Pure Becoming:** We Need To Explore The Nature Of An Event. Events Assume Becoming, Since They Refer To States In The Past And The Future In A Way Which ‘Eludes The Present’. This Is Paradoxical But Still Makes Sense. Plato Tried To Distinguish Between Limited Fixed Things And Pure Becoming, But The Two Cannot Be Separated. Instead, A Dualism Is Hidden In Material Bodies. We Need To Introduce The Notion Of A Simulacrum Which Is Neither Copy Nor Model [Massumi Has A Useful Article On This—Roughly, The Issue Is That What We Take To Be Material Reality Is Actually A Simulacrum Of The Virtual, A Limited Condensation ‘Beneath Things’ (2)]. It Also Avoids The Problems Of Subsuming Reality Under The Idea. Sometimes, This Is Indicated By A Summer The Peculiarities Of Language, Which Can Seem To Flow Over Specific Referents. This Provides A Clue That There Is Some Dimension To Language Which Serves To Come To The Aid Of More Specific Attempts To Name And Describe. There Are Implications For Identity. Fully Grasping Becoming Means That Identities Are Infinite, Incorporating Future And Past, Active And Passive, And Even Cause And Effect. Language Attempts To Limit This Infinity, But Still Often Alludes To It [With Open-Ended Statements Or Generalisations]. As With The Alice Stories, This Also Disrupts The Conventional Notion Of The Personal Identity. Normally, This Is Maintained By Some Underlying Commonsense Or Knowledge, As When ‘The Personal Self Requires God And The World In General’ (3). Becoming Threatens This Stability With The Paradox Of Events, Which Can Penetrate Even Commonsense—It Is Not Just A Doubt About Reality, But A Clear Indication Of The ‘Objective Structure Of The Event Itself, Insofar As It Moves In Two Directions At Once, And Insofar As It Fragments The Subject Following This Double Direction’ (3). **Second Series Of Paradoxes Of Surface Effects:** Stoics Divided Things Into Bodies And States Of Affairs, ‘Actions And Passions’ (4). There Is Also Some Cosmic Unifying Quality, Always In The Present. Bodies Can Interact And Cause Effects In Each

Other, But These Effects Are Incorporeal, ‘Logical Or Dialectical Attributes... Not Things Or Facts But Events’ (5). They Have The Kind Of Subsidiary Existence, Acting As Verbs, And They Are Infinitives—The Example Is A Cut Inflicted On The Body, Which Is Seen As An Incorporeal Surface Effect, Compared To The Actuality Of Bodies And Their Mixtures. This Argument Had Important Implications For Understanding The Causal Relation. Specific Bodily Causes Produce Other Bodies, Linked By Some Cosmic Unity Or Destiny. Similarly, Effects Can Be Seen As Having Bonds Between Them, But Effects Can Never Be Causes In Themselves. They Can Only Be “Quasi-Causes’ Following Laws Which Perhaps Express In Each Case The Relative Unity Or Mixture Of Bodies On Which They Depend For Their Real Causes’ (6). These Combinations And Bonds To Provide For Some Emergent Qualities, Which Means That Destiny Can Be Avoided. An Alternative Is Offered By The Epicurean Classification Of Different Kinds Of Causes Which Are Relatively Independent [And So Can Interact], And This Is A Kantian Idea Too. There Is A Reference Back To The Capacities Of Language To Offer ‘A Declension Of Causes...[And]... A Conjugation Of Effects’ (6). Stoic Philosophy Introduces The Notion Of A Something Behind Both Specific Material Beings And Incorporeal Events. The Idea Must Belong To ‘This Impassive Extra-Being Which Is Sterile, Inefficacious, And On The Surface Of Things: The Ideational Or The Incorporeal Can No Longer Be Anything Other Than An “Effect”’ (7). This In Turn Leads To A Change Of Metaphor From Surface/Depth To Just Surface, To A Series Of Effects Which Are Manifestations And Are Of Different Types. We Have A Notion Of Possibilities, Of Ideality Itself, Rather Than The Platonic Idea, But With No ‘Causal And Spiritual Efficacy’ (7). The Simulacrum Now Appears On The Surface, Rather Than Being Hidden In The Depths. Events As Effects Combine Past And Present, Active And Passive, All Of Which Are Located Elsewhere As Causes. The Relation Between Events Can Only Be Quasi-Causes. Stoics Saw Dialectical Analysis Has Explorations Of These Combinations, Once They Had Been Expressed In Propositions—Dialectics As Conjugation. Language Also Enables Us To Go Beyond Events Into The Possible Or Becoming. The Relation Between Propositions And Specifics Is Itself Still Paradoxical—‘Chryssipus Taught “If You Say Something It Passes Through Your Lips, So If You Say “Chariot”, A Chariot Passes Through Your Lips’ (8). It Is Deliberate Nonsense In The Anglo American Sense, Or Humourous Play On The Surface, As Opposed To An Ironic Exploration

Of Depths And Heights. Lewis Carroll Did Something Similar In Alice. [A Commentary On Alice Ensues, Stressing The Surface Rather Than The Underground World, And Picking Up The Disdain Lewis Carroll Felt For Boys Who Did Not Like To Operate At The Surface. Left Handers And Stutterers Can Sometimes Remind Us Of The Paradoxes Of The Surface, However, Which Can Defeat Commonsense Understandings]. **Third Series Of The Proposition:** Describing Events As Propositions Raises The Question Of How Best To Analyse Surface Events. There Are Three Possibilities: Denotation [Roughly, A Direct Connection Between Words And Images Which Represent States Of Affairs, As In Indexical Signs. Here, Propositions Are Either True Or False, And May Be True In All Cases]; Manifestation -- A Relation Between The Proposition And The Person Expressing It, Statements Of Desire And Belief. These Are Causal Relations: 'Desire Is The Internal Causality Of An Image With Respect To The Existence Of The Object Or The Corresponding State Of Affairs' (13). Belief Anticipates Production Of An Effect By A Cause. Manifestation Includes Denotation, Makes It Possible. "'I' Is The Basic Manifester' (13). Manifestation Is 'The Domain Of The Personal, Which Functions As The Principle Of All Possible Denotation' (13). The Issues Here Turn On Veracity Other Than Truth And Falsehood, The Avoidance Of Illusion. Signification, The Relation Of A Word To Universal Or General Concepts, And Connections To Implications Which Have To Follow The Rules Of Syntax. Again, Signifying Involves Conceptual Implications Referring To Other Propositions, As In Premises Or Conclusions. This Involves A Certain Indirect Process, Implication Or Assertion, Instead Of Truth Or Veracity, Which Remain As Possible In Certain Conditions. However, It Is Not Just Formal Logical Operations That Are Involved, But Notions Of Probability Or Even Moral Terms Such As Promise Or Commitment. Error Produces Not Falsehood But Absurdity [Looks Really Close To Habermas And The Three Validity Claims Here]. Signification May Not Be Primary, Since All Language Begins From The Standpoint Of The 'I', But There Is An Assumption That Propositions Must Be Understood By Others And Have A General Force. This Implies That Manifestation Has Primacy. But Signification Is Implied, And [In Social Relations] Would Be The Basis Of Manifestation. It Is The Difference Between Langue And Parole. Particular Utterances Only Make Sense Against The Background Of Constant Concepts. This Is Extended To Particular Desires And Beliefs, As Opposed To 'Simple Opinions' Which Would Not Signify (16). Actual

Utterances Often Involve Truth Claims And General Signification [And Other Presuppositions, Which Are Technically Infinite]. This Is The Paradox Facing Pure Logic, Solved By A Form Of [Smuggling]: ‘Implication Never Succeeds In Grounding Denotation Except By Giving Itself A Ready- Made Denotation, Once In The Premises And Again In The Conclusion’ (16). So Actual Propositions Feature Circular Relations Between Signification, Manifestation, And Denotation. There Might Even Be A Fourth Dimension—Sense, But Introducing This Will Depend On Making Relations Theoretically Consistent—It Is ‘Not Simply A Question Of Fact’ (17). To Approach The Issue, We Ask Whether Sense Might Be Located In One Of The Existing Three. Denotation Concerns Itself With Truth And Falsity, Which Is Too Narrow. The Mere Relation Between Words And Denoted Things Is Too Paradoxical To Always Make Sense, As In The Example Of Speaking The Term ‘Chariot’. Instead, Denotation Presupposes Sense. Manifestation Does Involve Some Manifesting Subject Which Initiates, So May Be Sense Is Itself A Subjective Matter Of Beliefs And Desires Of Persons—But Subjects Only Possess This Ability To Speak Because Of A General System Of Signification In Language. It Looks Like Sense Must Be Identified With Signification—But Signification Is Linked In A Circular Relation With Denotation And Manifestation. Perhaps It’s Necessary To Think Of Different Forms Of Possibility Of Propositions—Logical, Physical, Syntactic And So On. This Might Serve As Foundations For Sense, But This Would Be An External Foundation, Independent Of Speech [I Think The Problem Is The Connections Between Any Foundations And Actual Act Of Speech, Whether Anything Would Escape The Foundations]. The Concept Of Truth In Particular Implies Independence From Form. This Independence, Separate From Conceptual Possibilities In Signification, Is What Constitutes Sense. This Is The Fourth Dimension. It Is ‘An Incorporeal, Complex, And Irreducible Entity, At The Surface Of Things’ (19). There Is Philosophers Have Discovered And Rediscovered This Quality. It Is The Idea Of A Something Again, Beyond The Propositions And The Terms And The Objects Which Are Denoted, Beyond The Subjective I And Things Which Are Expressed. Sense Is Irreducible To Propositions, And It Is And Must Be “Neutral,” Altogether Indifferent To Both Particular And General, Singular And Universal, Personal And Impersonal’ (19). There Is Been Little Agreement About This Possible Fourth Dimension, Whether It Exists Simply In The Form Of Some Enquiry. It Is Not Even Immediately Useful Because It Is Neutral. It Can Only Be

Inferred Indirectly, By Questioning Characteristics Of Propositions As Above— This Is ‘Inspired In Its Entirety By Empiricism... [Avoiding Notions Of Essence Or Idea, And Knowing]... Have To Track Down, Invoked And Perhaps Produce A Phantom At The Limit Of A Length Or Unfolded Experience’ (20). It Might Be What Husserl Called ‘Expression’, Lingering In Terms Such As The Noema, As Pure Appearance, Outside Denotation Or Manifestation, Linked In Complex Ways To Appearances. In The Same Way, Sense Does Not Exist Outside Propositions Exactly, But ‘Inheres Or Subsists’ (21). It Is Not Just An Expression, But An Attribute, Not Just Of The Proposition, But ‘Of The Thing Or State Of Affairs’ (21), [The Potential, ‘To Be Able To Be Green’ Rather Than Just The Denotation ‘Green’ Is The Example Here]. It Is Said Of A Thing, So It Depends On Propositions Which Express It And Is Therefore Not Separate From The Proposition. It Is Something Else, Both The Expressible, And The State Of Affairs: ‘It Turns One Side Towards Things And One Side Towards Propositions’ (22). It Is What Joins Propositions And Things. [It Is A Becoming]. It Operates On The Surface, Rather As Mathematics Does, Or The Nonsense Of Carroll. It Is The Operation Of Sense That Produces [Meaningful] Paradox. **Fourth Series Of Dualities:** Important Dualities Exist Between Causes And Effects, And ‘Corporeal Things And Incorporeal Events’ (23). This Is Extended To A Duality Between Things And Propositions, Bodies And Language. This Is Expressed In Lewis Carroll As A Duality Between Eating Or Speaking—The Former Is A Matter Of Bodies Actions And Passions, And The Latter Movements Of The Surface And ‘Ideational Attributes Or Incorporeal Events’ (23) [Lots Of Examples From Alice About Being Presented To Food And Having Food Presented To You]. The Normal Relationship Can Be Distorted By ‘Verbal Hallucinations... Unrestricted Oral Behaviour... And Various Disorders Of The Surface’ As Bodily Matters Intrude—Stuttering, Left Handedness (24). Sense Is Always Expressed In Propositions, But It Lies In States Of Affairs, It Happens To Things. In This Sense, Bodies And Language Are United In The Production Of Sense, Existing ‘On The Two Sides Of The Frontier Represented By Sense’, Which Constantly Articulates The Differences (24)—Things Include ‘Ideational Logical Attributes Which Indicate Incorporeal Events’, And Propositions Include Both Denotations And Expressions, Names And Adjectives, And Verbs [The Latter Indicating Becoming And Chains Of Events] (24) [Illustrated With Words By Humpty Dumpty]. This Duality In Propositions Represents Two Dimensions, The

@@@@
 @@@@
 @@@@

Page 332808

commission, it is my sincere entreat, earnest beseech, fervent appeal to kindly pardon me and the error is absolutely inadvertent and in deliberate. Let not any sensibilities, susceptibilities, and sentimentalities be hurt. Parson's pattern variables provide a way of categorizing the types of choices and forms of orientation for individual social actors, both in contemporary society and historically .I want to put on **record with humble gratefulness the help by American Physical society, nature and other Noetic institutes of US who sent lot of alerts for my reference which was very valuable**, and could not have been found by me despite assiduous and fervent search. Most important type is that concerning the stability of solutions near to a point of equilibrium. This may be discussed by the theory of Lyapunov. In simple terms, if all solutions of the dynamical system that start out near an equilibrium point x_e stay near x_e forever, then x_e is Lyapunov stable. More strongly, if x_e is Lyapunov stable and all solutions that start out near x_e converge to x_e , then x_e is asymptotically stable. The notion of exponential stability guarantees a minimal rate of decay, i.e., an estimate of how quickly the solutions converge. The idea of Lyapunov stability can be extended to infinite-dimensional manifolds, where it is known as structural stability, which concerns the behavior of different but "nearby" solutions to differential equations. Input-to-state stability (ISS) applies Lyapunov notions to systems with inputs. von Neumann stability is necessary and sufficient for stability in the sense of Lax–Richtmyer (as used in the Lax equivalence theorem): The PDE and the finite difference scheme models are linear; the PDE is constant-coefficient with periodic boundary conditions and have only two independent variables; and the scheme uses no more than two time levels (See Wikipedia) Von Neumann stability is necessary in a much wider variety of cases. It is often used in place of a more detailed stability analysis to provide a good guess at the restrictions (if any) on the step sizes used in the scheme because of its relative simplicity. Albeit forwarded in nine module systematizations, the entire gamut is to be seen in a single shot, and the presentation of nine schedule twenty seven storey models is to circumvent typing of hundreds of superscripts and subscripts. In fact the statement is made inclusive of all previous models, and the variables are definitely different for each schedule, which again is reinstated due to typing of corresponding variables millions of systems, a fastidious and fussy work again. I beg pardon for any inconvenience caused to the readers due to such utilization of convention. I am grateful to Professor Chadralekha MD PhD. Tagore medical College, Chennai for deliberations and discussions on Medicine. To discussions on Physics topics credit goes to Dr. A.S. Krishna Prasad, Former Director DRDO, and Bangalore Chapter. Prof. Sunita MSc. PhD., of MS Ramaiah University helped me with valuable suggestions on Aerodynamics and propellant chemistry. Sir KVB Pantulu, former Chairman of NALCO, ESSAR Steels helped in formatting process and project management advices.

Note: Here we talk of the characteristics of systems which satisfy the condition of cosmological constant or any other for that matter or variable or axiomatic predication. . There are lots of zeroes corresponding and concomitant to the second law of black holes. Infact as many as that of extant and existential blackholes exist. At the outset, it is to be stated that there are hadrons in every system. Supersymmetry between forces and matter, with both open and closed strings; no tachyon; **group symmetry is SO (32) and its axiomatic predications, predication anteriorities, character constitution shall be extant and existential in very many systems, and the characteristics are taken in to consideration in the classification scheme. Many systems have such fundamental instabilities like that of quantum gravity and characteristics of those systems form the citadel and fulcrum, bulwark and manor, mainstay and reinforcement, alcazar and chateau theory has a fundamental instability on which the classification schémas are valid. There are lots of systems which follow the axioms of string theory It is the characteristics of this system which are taken in to consideration in the classification scheme. "All institutionalization involves common moral as well as other values. Collectivity obligations are, therefore, an aspect of every institutionalized role. But in certain contexts of orientation-choice, these obligations may be latent" (Parsons, 1951, p. 99). There are various systems that have the same bastion, support system, stylobate and sentinel as that of the Deleuzean terms and predications and phenomenological methodologies systemized. Look at this beautiful passage whose stability analysis would be eye opener. To "deconstruct" is not the same as to destroy. Deconstruction attempts to undo logical contradictions, to overturn rigid conceptual oppositions while releasing new concepts and meanings that could not be included in the old system. At the heart of Western metaphysics, for example, Derrida finds the opposition between "speech" and "writing." This binary logic functions in an illicit way to establish speech as the means of giving "presence" to the**

world, while writing is deemed derivative and inferior In Derrida's sense of "grammatology," however, all production of meaning is writing and subject to the infinite play of signification. By taking away the transcendental signified and advancing the concept of "différance" (language organized around difference and deferred, or mediated, understandings), Derrida, like Nietzsche, wants to leave us without transcendental illusions, metaphysical unities, and foundations that constrain thought and creativity. **THE POSTMODERN TURN IN PHILOSOPHY: THEORETICAL PROVOCATIONS AND NORMATIVE DEFICITS** by Steven Best and Douglas Kellner <http://www.gseis.ucla.edu/faculty/kellner/kellner.html> Each and every system has electrons, neutrons and protons and for that matter quarks. There shall be strong nuclear force and weak nuclear force. There are many systems that satisfy the criterion specified by the equation, principle or statement in question. Characteristics of the investigating systems form the bastion for the classification scheme and doxa thereof. Systemic differentiation is conducted. Despite gravity being constant, there exists gravity between two objects, and this could be taken as a system. Depending upon some parametric representationalities, functionalities, advantageousness, appropriateness, benefit, facilitation, fittingness, helpfulness, instrumentality, merit, practicality, serviceability, suitability, and usefulness, utility, these systems could be classified in to various categories. In respect of an equation, there shall be many systems that satisfy the given equation. Equations themselves could be by the utilization of the model solved term by term as has been exemplified and illustrated many time in the previous papers. There is lot of systems that could be brought in to the orbit of and gamut of the theory in question which the investigatable systems satisfy the axiomatic predications and postulation alcovishness of the **systems in question. Towards the end of classificational consummation, consolidation, corporation and concatenation we take the characteristics of the systems, the predicational interiorities, ontological consonance and primordial exactitude, accolytish representations, functional topology, apocryphal aneurism and atrophied asseveration, event at contracted points, and other parameters as the bastion and stylobate of the stratification purposes. Such totalistic entities would have easy paradigm of relational content, differentiated system of expressly oriented actions with primary focus and locus** of homologues receptiveness and differentially instrumental activity, variable universalism and particularism, imperative compatibilities and structural variabilities, interactional dynamical orientation, institutionalization and internalisation of pattern variables common attitudinal orientation of constitutionalisation of internalized dispositions, and a qualitative gradient of structural differentiation and ascribed particularistic solidarity abstraction or interactional dynamics, internal differentiation, structural morphology, formal characterization, concept formulation, phenomenological methodologies, constituent structure, transformational minimal condition, paradigmatic feasibilities, programmatic plausibilities, comparative variability, normative aspect of expectational prediction, projection and prognostication as consideration of the investigatory systems. Any scale can be used that is convenient to the classification scheme. It is important to note that the scheme of classification and the stratification doxa must not be adversely antagonistic, inexorably irreconcilable, antithetically antipodal, diametrically opposed, repugnantly retrogressive, inimically inverse, violatively unsimilar, diversely dissimilar, antipathetically antithetical, conflictingly combative, obstructively pugnacious, inimically obstructive, repellently restrictive, disputatiously gainsaying or conformingly pugnacious to the axiomatic predications and postulation alcovishness of the theory in question or the equation representative or constitutive thereof. Sole intention, main objective and primary aim is twofold. One is towards the end of circumvention of the extra equation and the concomitant and corresponding variable therein. Second is avoidance of clustered congest, swarmed huddle, mustered pack, sardine squash, and swamp throng in the scheme of classification. Only thing that is sought out is the consonance in the entire diaspora and body fabric of the systems under study. This statement is true and holds unmistakably true for all the papers and I sincerely entreat readers to remember the statement and read the paper against this background. When we write $A+B$ we mean by that B is being added to A or vice versa. It is like adding milk to water and water to milk. There may or may not be a time gap. As said earlier there may be many systems that satisfy the conditions of the equations and those systems that are investigatory or investigatable are taken in to consideration based on their characteristics in the classification doxa. When there are more than two entities, we can take logarithm and find the value of that factor to be **taken with anti log to obtain prediction and projected values of the model. In case of $A-B$, we are removing B from A and that means B is eating up A. These factors are taken in to consideration in the application of the model to equations. $\log(ab)$ and $\log(a+b)$** is well defined. Model stands out as universal

testament and template for application to each sentence and equation what with the quantification process done and the correlations well defined. In the eventuality of non existence of any connection at some phase, the model would have the accentuation and attrition coefficients and detritions coefficients as zero rendering the equations of concatenation simpler. Projection formula which incorporates in its diaspora the initial values provide authenticative determination, unimpeachable validation, incontrovertible establishment, apodictic evidence, reliable roll out, unfailing cinquecento quattrocento trecento, incontrovertible indication of the final finale, notwithstanding appellation, appellative, brand, cognomen, compellation, designation, flag, handle, identification, label, moniker, nomen, slot, style, surname, tab, tag, term, title designation, appellative, class, classification, denomination, description, epithet, of the classification scepter, scenario, scimitar, schottische. There is pure and impure consciousness in every one. Gratification producing and deprivation producing one's can be easily classified from individual general ledgers. Similar analogy holds for collective general ledger and cosmic general ledger or nature's general ledger. It is also to be noted that while dealing with equations towards the end of consummation of the measure, it is necessary that the two variables are to be classified in to three sections and each one would have the adventitious and decidedly stated relationship whereby the fundamental equations are drawn up and the analysis made. All the parametric representationalities, conditionalities, orientationalities remain unequivocal as stated in the variables stated in to consideration section and are different from module to module. In essence the paper is to be read as holistic one with the sole intention, primary objective and *raison d'être* to build a TOE. Towards the end of circumvention of typing hundreds of superscripts and subscripts which would be a sardine squash and the concomitant operational difficulties, model is presented in piece meal of nine modules. Logarithms are to be taken in respect of those which incorporate more than one variable in bra-ket. Values of $\log(ab)$ and $\log(a+b)$ are readily available. Anti logarithm shall be taken at the end to predict, project, prognosticate the value of the variable. This is true for tensors, vectors and other variables too. Affirmational assertion, and explanation justification, statement of vindication, annotational commentary, and explicational glossary for each and every system changes and physical interpretation of results is one thing that is to be with earnest endeavour and feverish and febrile expediency. Editions never ever mean the same and identity of parameters and this has been explicitly and unmistakably stated in the model *a priori* itself. Akin and analogous, cognate and concurrent, correspondent and congruent, comparable and complementary, synonymous and duple, tantamount and agnate, commensurate and correlative representation is only to highlight the importance and subterfuge the replication of work. It is my fervent solicitation to kindly bear with me for any lapses, notwithstanding the orchestrated efforts for a paper without any minor errors. Postulation predication, conclusive presumption, differential presuppositions, underscored decidedly axiomatic statement of the statement, equation form the bye word or the watch word in the aggrandizement-amplification, caricature and crock, understatement- unembellishment, elocution-emphasis, enunciation-inflection, announcement-argument, articulation, assertion, asseveration, choice of words, commentary and communication, declaration and definition, delivery-diction, elucidation-emphasis, enunciation-execution, explanation, exposition-formulation, idiom, interpretation and intonation tone and tenor of stratification. Now, how do find the reaction of systems to these singularities. You do the same thing a boss does for you. "Problematize" the events and see how you behave. I will resort to "pressure tactics". "intimidation of deriding report", or "cut in the increment" to make you undergo trials, travails and tribulations. I am happy to see if you improve your work; but may or may not be sad if you succumb to it and hang yourself! We do the same thing with systems. systems show conducive response, felicitous reciprocation or behave erratically with inner roil, eponymous radicality without and with blitzzy conviction say like a solipsist nature of bellicose and blustering particles, or for that matter coruscation, trepidational motion in fluid flows, or seemingly perfidious incendiaries in gormandizing fellow elementary particles, abnormal ebullitions, surcharges calumniation and unwarranted(you think so but the system idoes not!) unrighteous fulminations. Amount of emotion or affect that is appropriate or expected in an given form of interaction. Particular individuals and diffuse obligations (see c and d) are associated with affectivity, whereas contacts with many individuals (universalistic) in a bureaucracy may be devoid of emotion and characterized by affective neutrality. Affective neutrality may refer to self discipline and the deferment of gratification (eg Weber's spirit of capitalism). In contrast, affectivity may be associated with expressing emotions. Adams and Sydie also refer to affective neutrality being associated with ego control (p. 15). (Parsons: Wikipedia). So the point that is made here is "like we

problematize the "events" to understand the human behaviour we have to "problematize" the events of systems to understand their behaviour. This statement is made in connection to the fact that there shall be creation or destruction of particles or complete obliteration of the system (blackhole evaporation) or obfuscation of results. Some systems are like "inside traders" they will not put signature atoll! How do you find they did it! Anyway, there are possibilities of a CIA finding out as they recently did! So we can do the same thing with systems to. This is accentuation, corroboration, fortification, .fomentatory note to explain the various coefficients we have used in the model as also the dissipations called door. In the bank example we have clarified that various systems are individually conservative, and their conservativeness extends holistically too. that one law is universal does not mean there is complete adjudication of nonexistence of totality or global or holistic figure. Total always exists and "individual" systems always exists, if we don't bring Kant in to picture! For the time being let us not! Equations would become more ensorcelled and frenzied..... philosophy merges with ontology; ontology merges with univocity of being; analogy has always a theological vision; not a philosophical vision; one becomes adapted to the forms of god; self and world; the univocity of being does not mean that there is one and the same being; on the contrary, beings are multiple and different they are always produced by disjunctive synthesis; and they themselves are disintegrated and disjoint and divergent; membra disjuncta. the univocity of being signifies that that being is a voice that is said and it is said in one and the same "consciousness". In consciousness research, two rival sets of theories can be recognized: (A) Scientific material interpretations of consciousness are based on axioms that view consciousness in the context of highly advanced intentional processing of information in which subject-object relations evolve, and (B) humanistic interpretations of consciousness are based on axioms that view consciousness in the context of, say, "centered pulsations" that enable a conscious agent to act from his or her center of awareness. In this paper I will argue for the selection of axioms that favor humanistic interpretations of consciousness. The Pursuit of Autonomy Interdisciplinary Observations to Human Consciousness: Wautischer, Helmut (1993) the Pursuit of Autonomy. Interdisciplinary Observations to Human Consciousness: Social Neuroscience Bulletin, 6 (4). pp. 52-56. Everything about which consciousness is spoken about being is the same for everything for which it is said like gravity; it occurs therefore as an unique event for everything; for everything for which it happens; eventum tantum; it is the ultimate form for all of the forms; and all these forms are disjointed; it brings about resonance and ramification of its disjunction; the univocity of being merges with the positive use of the disjunctive synthesis, and this is the highest affirmation of its univocity like gravity; it is the eternal resurrection or a return itself, the affirmation of all chance in a single moment, the unique cast for all throws; a simple rejoinder for Einstein's god does not play dice; one being, one consciousness, for all forms and all times. a single instance for all that exists, a single phantom for all the living; a single voice for very hum of voices; or a single silence for all the silences; a single vacuum for all the vacuums; consciousness should not be said without occurring; if consciousness is one unique event in which all the events communicate with each other; univocity refers both to what occurs to what it is said. This is attributable to all states of bodies and states of affairs and the expressible of every proposition. So univocity of consciousness means the identity of the noematic attribute and that which is expressed linguistically and sense fully. Univocity means that it does not allow consciousness to be subsisting in a quasi state and but expresses in all pervading reality; **CONSCIOUSNESS AND ITS UNIQUENESS June 12, 2012 at 8:40pm (See Deleuze Logic of sense, Wikipedia and Stanford encyclopedia for more details)** Atrocious contrivance and device, gratificational primogeniture, calamitous dodge and expedient, depraved, destructive, disastrous, execrable gambit and gimmick, great solace and succor, iniquitous, injurious, loathsome, low, maleficent, malevolent machination and maneuver, promethaleon of candor, frankness, honesty, honor, ingenuousness, innocence, openness, reality, sincerity, truthfulness, spiteful, stinking, ugly, unpleasant, unpropitious play and ploy, progenitor of jurisprudence and circumspection, racket and ruse, savvy, scam, stratagem, subterfuge, tactic and wile, forthrightness, honesty, truthfulness embodied and personified is the structural predisposition and dispensation of the complex. Nobody need have to take the Brahman-Anti Brahman agency to be a recalcitrant, repugnant, and refractory proposition. God gives and also takes away. That is the point made. And of course lesser gods, and God doth follow them and they follow God. That is why they are lesser gods. And wield the power on lesser mortals like us. When we mean gratification and deprivation we mean celestial Brahman Anti Brahman and Terrestrial Brahman and Anti Brahman agencies producing such happiness or sadness by any means such as (blending- amalgamation,

adulterant-adulteration, amalgam-amalgamation, blend- combination, composite- compound, debasement-denaturant, fusion-hybrid, intermixture-reduction, amalgam- mixture, composite- compound, fusion-mishmash, aggregate- alloy, amalgamation-blend, commixture-composite, composition-compost, conglomerate-fusion, goulash-medley, mishmash-stew, blend-coalition, commixture-compound, federation-heating, integration-junction, melting-merger, synthesis-unification, assortment- combination, adulteration-alloy, assimilation-association, batter-blend, brewed combine, composite compound, concoction confection, conglomeration cross of money, orientation(dis), penury creation, murder, mayhem plunder, pillage, apocalypse, Armageddon in various forms and permutation and combination of Manichaeian artifice, agathokakological malevolence, agley chicane fourberie, fraud, furtiveness, gambit, awry and bad ,flagitious mechanizations, cacodemonic gambit, hanky-panky, intrigue, machination, deprecatory and diabolic underhandedness and wiles, energumenical skulduggery, goetic chicanery, lenocinant stratagem, malefic and maleficent strategy, malominous sell, sellout, sham subterfuge, peccable, humbug, imposture, misrepresentation peccant fraud, fraudulence, hoax, quod , sclerate pretense, ruse, sell, sellout, sham stratagems' of maladroitness manoeuvres , sinister ,tortious and venal sophistry, contingency enterprise, endangerment emprise, jeopardizing adventure, jejune jeremiad, avoidable inertia, latent passiveness: with each action accounted and showing balance in the nature's general ledger as also actionlessness:e). All these are recorded in Nature's general ledger. Neuron DNA encapsulates all the actions passions, actions, interactions, transactions or the lack of it at various levels of individual, collective and cosmic levels. In celestial Brahman Anti Brahman ledger there is cessation of dualities. One more point to be made is if $5=3+2$, then we can say 5 gormandizes 3 or 5 gobbles up 2. Quintessentially such a statement arises out of the difference of LHS and RHS term wise. It must be remembered despite all theoretical abstractions and generalizations be it proton mass or Higgs Boson, or topology of a space, it is notwithstanding the fundamentality of nature is just a number. Just everything in Physics and mathematics is just a number. Let us assume I buy rice in Bangalore (District) worth 5k.g. I can always divide 5k.g.s of rice in categories of Bangalore (South), Bangalore (west), Bangalore (East) and Bangalore (South). On similar basis, there are billions of neutrons, protons and concomitant interactions which would satisfy the given numerical value. Example of gravity above makes things clear. Everything atleast is a function of time and someplace or some other coordinates, subordinates or superordinates. Mass of proton is 2.47 does not mean total mass of all protons is 2.47. It is this argument that is used in the classification scheme. By the very nature, self the unmonitoring but the witness consciousness agency identifies itself with something, be it a profession, an actor on the screen, when it does not have a identity. Such examples are legion, a Charlie Chaplin in "The Kid" or Raj Kapoor in "Jagte Raho". These are people who just wanted to live and living without being put under surveillance every moment is the greatest part of their life. This is "Absolute Subjectivity" of Shiva. Such a state is always craving for identification for in pure consciousness all identities are lost. It is this "identitylessness" or having very minimal identity is what pervades most in India and this is due to expansion of individual consciousness. When that happens, be it education, or wealth, the "identity" becomes so well entrenched, they become the exact opposite of "pure consciousness", "impure consciousness" (Shakti). This is a state of "Absolute or relativistic objectivity". There was this case of a lady who went to such an extent of planning, she was sure that whatever she does must work out. When it did not, she lapsed in to "identity crisis". Such thing as this might happen in well developed countries where the competition is extremal and to be achieved at any cost. Realisation that one is not the doer but the witness consciousness is just like watching a movie in which one has acted. It is in this sense "advaita", "non duality" seems to be of cardinal and preeminent importance for the analytical treatment of "identity" and "identity lessness", both happen in space time. "Multiverses" in this sense is also like a different film of an actor released at different theaters and it is in this sense we state that that every object in space time is a multiverse. You must have totally immersed in doing a "string theory" problem or watching a "film", oblivious of surroundings. Essentially creation (space time) also is what is happening in mind and honed in to you billion times. Identification of anything with a designation or signification is one that leads to association within spacetime, which would only have pathological consequences and detrimental ramification as stated above or positive implications of "happiness" because of the identity itself. Search or 'show" has to go on! It gives a sense of consummation. a watching pot never boils; donot go to check nature; you are disturbing it; it will never show its true colours bhakti and Virakthi, gratification and deprivation, raga and dwesha is essential to live in space. it is the quantum interest thereof that produces energy to live

.gratification and deprivation which form the bastion, pillar, post and stylobate and sentinel of all human actions depends on tamas, rajas, and sattva of individual or insentient object. Verily tamas, rajas and satva are the characteristics of space itself, nay the behavioural pattern and attitudinal orientation. There are peaceful places like Vienna or Sringeri and there are violent places like others. Space is warped and woofed by the Akshara, the quantum information of primordial syllable Om. cosmic general ledger is the one vibration of which is brahman; quantum information possessions thereof form the sentinel of all transactions (Gargi to Yajnavalkya: my interpretation) adhocism, nonsequentiality of events, anti sequentiality of events, simultaneity of phenomenon, concurrent happenings, improbability of phenomenon happening itself are characteristics of brahman-anti brahman (Terrestrial and celestial) are the cause of non predictability of happenings like in black hole, or in the areas of operation of brahman-anti brahman (Terrestrial and celestial). brahman AntiBrahman actions, interactions, decisions, circumspection, adhocism, satisfies sensitive to initial conditions; topologically mixing; dense periodic orbits, the axiomatic predications and postulation alcovishness of chaos theory. Like gravity is a characteristic of space, so is tamas, rajas, sattva, with spaces knotted together by quantum information (for models see below). accidents are preplanned actions of Terrestrial and celestial Brahman AntiBrahman. All actions are perpetrated by them, be it train on fire or death. Belief systems play an important role in the determination of tamas, rajas, and satva state of human beings. There is neither cause nor effect. All actions are performed by Terrestrial and celestial Brahman anti Brahman and it is off the record. psychic energy is negative equivalent to $-mc^2$, so that second law of thermodynamics is satisfied on a holistic basis vis-à-vis material energy ParaBrahman is infinite individual, collective and cosmic consciousnesses are infinite (concomitant general ledgers) for they are in the evolution mode with respect to each other in that order. Later emanates from the former as transactions emanate from the conservative state of assets and liabilities. Is not zero true for infinite conservative systems inside space and time? zero is the only consummation, consolidation and conservation. if fifty out of seventy members come out of the room at whatever time you come out, talk about the grand limbo of oblivion and hibernation of lifeless world you are going to and if you are not a celebrity, then it is not coincidence, it is coordination. Nature is playing the same trick on you. that a generalised law is there and you can measure does not make things less good albeit recalcitrant, refractory and rogue particles occur time and again, by and large. Entering in to all living beings through his constant part on the new moon night, he is born therefrom next morning. Absolute ParaBrahman (zero) projects himself again and again on the screen of individual consciousness which appears of depairing despondent world. Some where the balancing act is done on karma. cosmic general ledger reflects upon itself to keep all transactional ties conservative (Brihadarnkayaka Upanishad: 1.5 14) for models see next paper through its divine power (i interpret man creates god; it is attribution and ascription of various Shaktis would the formation of god takes place like space and time) the self assumes (eb) the form of a deity man can contemplate and venerate, even though siva, the pure subject, can never in fact be an object of meditation (adhyeya). Until we realise our true identity with Sankara, he is worshipped and conceived to be a reality alien to ourselves. While we are in the realm of creation, he too is a creation or mode or appearing of the absolute, manifest to us in meditation, through his freedom as an eternal, omniscient being. There is no gulf between the created and the uncreated creator: nothing in reality, although an object of knowledge, ceases to be Siva: this is the reason why meditation [on this or that aspect] of reality bestows its fruit. The world of the senses and mind appears to the well awakened (suprabuddha) as a theophany, an eternal revealing of god in his creation. The doctrine of vibration declares that "there is no state in word, meaning or thought, either at the beginning, middle or end, which is not Siva." 59 to utter any word is, in reality, to intone a sacred formula. every act is a part of Siva's eternal cosmic liturgy, every movement of the body a ritual gesture (mudras), and every thought, god's thought. By what path are you not attainable? What words do not speak of you? in which meditation are you not an object of contemplation? What indeed are you not, o lord? 61 spandashakti, which accounts for the appearing of all things, is also the means by which deity in its many varied forms appears to man. Ksemaraja concludes: the ultimate object of worship of any theistic school differs not from the Spanda principle. The diversity of meditation is due solely to the absolute freedom of Spanda. 62 Sankara is not only the supreme object of devotion; as the static polarity of the absolute, he is the inner reality which holds together its Siva and Sakti outer manifestations. Phenomena are patterns of cognitions projected onto the surface of self-luminous siva-consciousness. There they become apparent, directly revealed to consciousness according to their

manifest form. siva is accordingly symbolized as the ground or surface of awareness, smooth and even like a screen (samabhittitalopama). 64 inscribed on this screen (kutfya) are the countless manifest forms which appear within it rendering it as diverse and beautiful as a fossil ammonite (falagrama). 65 **Siva is the sacred ground upon which the cosmic mandalas are drawn, the absolute surface of inscription which bears the mark (cihna) of the universe. Abhinava writes: the variety of this world can only be manifest if the highest lord, who is essentially the pure light of consciousness, exists; just as a surface is necessary for a picture. if external objects were perceived in isolation then, because 'blue' and 'yellow', etc., are self-confined and the perceptions [we have of them] refer to their objects alone and so are insentient, mute and dumb in relation to one another . . . how would it be possible to be aware that** an object is variegated? but just as depths and elevations can be represented by lines on a smooth wall, and we perceive [a female figure and think], similarly it is possible to be aware of differences in the variegated (contents of experience) only if all the diverse perceptions are connected together on the one wall of the universal light of consciousness. 66 Siva is the perfect artist who, without need of canvas or brush, paints the world pictures. the instant he imagines it, it appears spontaneously, perfect in every respect. The colours he uses are the varying shades and gradations of his own Spanda energy and the medium his own consciousness. Intractable disaffect, Mutinous obdurate, Rebellious recalcitrant, Seditious refractory, Insubordinate insurgent, antagonistically antipodean, discrepant incompatible, Contumacious coadjutant, Conflicting converse, Unyielding obdurate, the antagonistic and anachronistic dispositional ties of Anti Brahman are in fact the sine qua non of Brahman only (See Vishwaroopa of Mahavishnu, which contains the polarised positives on RHS and negatives on LHS). The universe is coloured with the dye of its own nature (svabhava) by the power of Siva's consciousness (a/). Rajanaka rama says: homage to him who paints the picture of the three worlds, thereby displaying in full evidence his amazing genius {pratibha}\ to Sambhu who is beautiful with the hundreds of appearances laid out by the brush of his own unique, subtle and pure energy. 68 analogously, at the microcosmic level, all the cognitions and emotions, etc., which make up the individual personality form the outward flow of essentially introverted consciousness. They are specific pulsations (vise\$aspanda) or aspects of the universal pulsation (samanyaspanda) of pure t consciousness. at the lower level, within the domain of maya, they represent the play in the fettered soul of the three primary qualities (gunas) or feeling-tones' which permeate to varying degrees his daily experience. These are: 1) sattva — the quality of goodness and luminosity which accompanies blissful experience both aesthetic and spiritual. 2) Rajas — the passion or agitation which oscillates between the extremes of 'light' and 'darkness' and characterises inherently painful experiences. 3) Tamas — the torpor and delusion which accompany states of inertia and ignorance. 69 the liberated soul recognises that these three are the natural and uncreated powers of pure consciousness. for him they are manifest respectively as: 1) Sankara 's power of knowledge (jnana) — the light of consciousness (prakasa); 2) the power of action (kriya) — the reflective awareness of consciousness (vimarsa)\ 3) the power of maya — which does not mean here the world of diversity, but the initial subtle distinction which appears between subject and object in pure consciousness. 70(doctrine of vibration: mark S.G. Dyczkowski) .to “deconstruct” is not the same as to destroy. Deconstruction attempts to undo logical contradictions, to overturn rigid conceptual oppositions while releasing new concepts and meanings that could not be included in the old system. at the heart of western metaphysics, for example, derrida finds the opposition between “speech” and “writing.” this binary logic functions in an illicit way to establish speech as the means of giving “presence” to the world, while writing is deemed derivative and inferior in derrida's sense of “grammatology,” however, all production of meaning is writing and subject to the infinite play of signification. By taking away the transcendental signified and advancing the concept of “différance” (language organized around difference and deferred, or mediated, understandings), derrida, like Nietzsche, wants to leave us without transcendental illusions, metaphysical unities, and foundations that constrain thought and creativity. The postmodern turn in philosophy: theoretical provocations and normative deficits by Steven best and Douglas Kelner <http://www.gseis.ucla.edu/faculty/kellner/kellner.html>. you cannot see beyond consciousness for that is the limit of your storage of information or the basis for understanding further information. Turiya is empty bucket; it is from which prajna the measure of arises; turiya measures the number of quantum information stores in other three states of waking, dreams, and dreamless deep sleep (Manduka Upanishad; my interpretation for models next paper).let there be path where angels tread: it is a force never born never dies the brahman an anti brahman verily there are parallel universes in (eb) in the ambit of space and time; the society of

sinners and the sinned with the grammar of evil; parallel universes contain(e) multiverses in them; all physics, philosophy, break down(e) in these universes; astronomical gratification ,deprivation, glorification, mortification, projections, axes, dimensions, rotations, foldings, concurrences, motions, shocks, are the axiomatic predications of multiverses and parallel universes; death is when self (the witness consciousness; register of individual general ledger) leaves the body; ostensibly parallel seem to have (e) an objective; parallel universes donot (e) have objectives; their paradigm of relational content, differentiated system of expressly oriented actions with primary focus and locus of homologues receptiveness and differentially instrumental activity, variable universalism and particularism, imperative compatibilities and structural variabilities, interactional dynamical orientation, institutionalization and internalisation of pattern variables common attitudinal orientation of constituionalisation of internalized dispositions, and a qualitative gradient of structural differentiation and ascribed particularistic solidarity abstraction or interactional dynamics, internal differentiation, structural morphology, formal characterization, concept formulation, phenomenological methodologies, constituent structure, transformational minimal condition, paradigmatic feasibilities, programmatic plausibilities, comparative variability, normative aspect of expectational prediction, affection- appreciation, awareness- discernment, emotion- feeling, gut reaction, heart insight, intuition, judgment, keenness, perceptiveness, rationale, sensation, sense, sensitiveness, sensitivity, sentiment, susceptibility, taste, vibes, admiration, aesthetic sense, affection, appraisal, assessment, attraction, awareness, cognizance, commendation, comprehension, enjoyment, esteem, estimation, grasp, high regard, knowledge, liking, love, perceptual realization, recognitional regard, relish, respect, responsiveness, sensibility, sensitiveness, sensitivity, sympathy, understanding, valuation, acknowledgment, airing, baring, , confessional defenselessness, denudation-denunciation, disclosure, display, divulgence-exhibition, hazard, introduction, jeopardy, laying open, liability, manifestation, nakedness, openness, peril, presentation, publicity, revelation, risk, showing, susceptibility , susceptiveness,susceptivity, unfolding, unmasking, unveiling, vulnerability, vulnerableness, action, affection, appreciation and ardor, behavior, capacity, compassion, concern, cultivation and culture, delicacy, discernment and discrimination, emotion, empathy, faculty, fervor, fondness, heat, imagination, impression and intelligence, intensity and intuition, judgment, keenness- palpability, passion, pathos, pity, reaction, refinement, sensibility, sensitivity, sentiment, sentimentality , sharpness, spirit, sympathy, tangibility, taste, tenderness, understanding, warmth, fond memories, hearts and flowers, homesickness, longing, pining, reminiscence, ,schmaltz, sentimentality wistfulness, yearning satiation is only ostensible; their job is to get the job done; their business is to mind other's business ;all laws of physics and finance breakdown; it is antilaw,piscatorial Piratish and bubonic bucaneerishness; there is free transactions between in multiverses; when reflected on the screen of consciousness they appear as extant existential universe. Transactions are astronomical and gratification deprivation, glorification and mortification incalculable; they have no barriers of region, religion, language which they create for others. theirs objectives are: corruption criminality, debasement debauchery, degradation evil immoral lewdness licentiousness, perversion sinfulness, vice viciousness, vileness wickedness, improbity corruption, atrocity decadence, degeneration degradation, depravity evil, immorality impurity, infamy iniquity, looseness lubricity, perversion profligacy, sinfulness turpitude, vice viciousness, vulgarity wickedness, disgrace bad reputation, abasement abuse, baseness black eye, blemish blur, brand comedown,contempt contumely, corruption culpability, debasement debasing, defamation degradation, derision disbarment, discredit disesteem, disfavor dishonor, disrepute disrespect, humbling humiliation ignominy ill repute, infamy ingloriousness, meanness obloquy,odium opprobrium, pollution ,put-down reproach, scandal scorn, slander slight, slur spot, stain stigma, taint tarnish, turpitude venality, enormity Horribleness , abomination atrociousness, atrocity crime and the positive aspectionalities thereof .let there not be any incursions where evil mercenaries in the garb of mendicants thrive with adversely antagonistic, inexorably irreconcilable, antithetically antipodal, diametrically opposed, repugnantly retrogressive, inimically inverse, violatively unsimilar, diversely dissimilar, antipathetically anti thetical, conflictingly combative, obstructively pugnacious, inimically obstructive, repellently restrictive, disputatious gainsaying or conformingly pugnacious Manichaeian artifice, agathokakological malevolence, agley chicane fourberie, fraud, furtiveness, gambit, awry and bad ,flagitious mechanizations, cacodemonic gambit, hanky-panky, intrigue, machination, deprecatory and diabolic underhandedness and wiles, energuminal skulduggery, execrable and goetic chicanery, lenocinant stratagem, malefic and maleficent strategy,

malominous sell, sellout, sham subterfuge, peccable, humbug, imposture, misrepresentation peccant fraud, fraudulence, hoax, quod, scelerate pretense, ruse, sell, sellout, sham stratagems' of maladroitness manoeuvres, sinister, tortious and venal sophistry, contingency enterprise, endangerment enterprise, jeopardizing adventure, jejune jeremiad, avoidable inertia, latent passiveness. statement for Mephistophelean mercenaries philosophy merges with ontology; ontology merges with univocity of being; analogy has always a theological vision; not a philosophical vision; one becomes adapted to the forms of god; self and world; the univocity of being does not mean that there is one and the same being; on the contrary, beings are multiple and different they are always produced by disjunctive synthesis; and they themselves are disintegrated and disjoint and divergent; membra disjuncta. the univocity of being signifies that that being is a voice that is said and it is said in one and the same "consciousness". everything about which consciousness is spoken about being is the same for everything for which it is said like gravity; it occurs therefore as an unique event for everything; for everything for which it happens; eventum tantum; it is the ultimate form for all of the forms; and all these forms are disjointed; it brings about resonance and ramification of its disjunction; the univocity of being merges with the positive use of the disjunctive synthesis, and this is the highest affirmation of its univocity like gravity; it is the eternal resurrection or a return itself, the affirmation of all chance in a single moment, the unique cast for all throws; a simple rejoinder for Einstein's god does not play dice; one being, one consciousness, for all forms and all times. a single instance for all that exists, a single phantom for all the living; a single voice for very hum of voices; or a single silence for all the silences; a single vacuum for all the vacuums; consciousness should not be said without occurring; if consciousness is one unique event in which all the events communicate with each other; univocity refers both to what occurs to what it is said. This is attributable to all states of bodies and states of affairs and the expressible of every proposition. So univocity of consciousness means the identity of the noematic attribute and that which is expressed linguistically and sense fully. Univocity means that it does not allow consciousness to be subsisting in a quasi state and but expresses in all pervading reality. Negative energy stabilising wormholes are considered candidates for darkmatter. An analysis of negative energy might be an useful ingredient of the thoughts of travel at speeds more than the velocity of light. Peirce explains that an "imperfect continuum" possesses "topical singularities, or places of lower dimensionality" which break continuity, while "the parts of a perfect continuum have the same dimensionality as the whole." (CP 4.642) Circular continua without singularities are time and space. (CP 6.210-212) By using math as his lens, Peirce envisioned singularities or lower dimensional holes in the imperfect cosmologic continuum, while maintaining that a probabilistic law of chance exists that entropy decreases (**Wikipedia**). On the side of philosophy, quite explicitly, Deleuze provides a critique of Husserl's "sleight-of-hand" change from a static categorical phenomenology to one supposedly more dynamically constitutive of reality. This a clear cut **chagrined circumvention, contraventional curbing, defeated disgruntlement, dissatisfactional downer, dragged failure, fizzled foil, grievance hindrance, impedimental irritation**, letdown, nonfulfillment, nonsuccess, obstruction, old one-two, **resent mental setback**, unfulfillment of the earlier theories makes Sartre contend that human existence is a conundrum whereby each of us exists, for as long as we live, within an overall condition of nothingness (nothingness)—that ultimately allows for free consciousness. But simultaneously, within our being (in the physical world), we are constrained to make continuous, conscious choices. It is this dichotomy that causes anguish, because choice (subjectivity) represents a limit on freedom within an otherwise unbridled range of thoughts. Subsequently, humans seek to flee our anguish through action-oriented constructs such as escapes, visualizations, or visions (such as dreams) designed to lead us toward some meaningful end, such as necessity, destiny, determinism (God), etc. Thus, in living our lives, we often become unconscious actors—Bourgeois, Feminist, Worker, Party Member, Frenchman, Canadian or American—each doing as we must to fulfill our chosen characters' destinies. However, Sartre contends our conscious choices (leading to often unconscious actions) run counter to our intellectual freedom. Yet we are bound to the conditioned and physical world—in which some form of action is always required. This leads to failed dreams of completion, as Sartre described them, because inevitably we are unable to bridge the void between the purity and spontaneity of thought and all-too constraining action; between the being and the nothingness that inherently coincide in our self. Deleuze insists on the absolute non-resemblance between what conditions (metastable, pre-individual plane of singularities) and what is conditioned (actualized or the individuated 'thing'). Only then do "the conditions of the true genesis become apparent," Deleuze writes (Logic of Sense 105). Without addressing this issue, Deleuze's borrowings from

Simondon, Lautman and structuralism remain untethered. Because of Bowden's decision to completely sidestep Deleuze's overt negotiations with phenomenology, his otherwise sophisticated commentary suggests a pedagogical narrowness. If, the omission of phenomenology strengthens Bowden's focus on the ontological priority of the event over substantial ontologies via the genealogy of figures and movements Deleuze outlines, it also incompletely articulates the problem of the sense-event as Deleuze formulates it. We have lost any idea of whom Deleuze takes to be his primary adversary, against whom he stakes out his position. This worry is most clearly manifested in Bowden's description of Deleuze's ontological prioritizing of events as offering a "transcendental ontology." For Bowden Deleuze's transcendental ontology defines events that "are ontologically prior to substances, 'all the way down'" (82). But Bowden provides a second definition: it is that the priority of events over worldly individuals means that the world of events "is not something external to the conditions of knowledge" (69).

SEAN BOWDEN the Priority of Events: Deleuze's Logic of Sense Sean Bowden, *the Priority of Events: Deleuze's Logic of Sense*, Edinburgh University Press, 2011, 296pp., \$40.00 (hbk), ISBN 9780748643646. Reviewed by David Scott, Coppin State University

Weltschmerz, agony, apprehension, blues, depression, dread, mid-life crisis, misgiving, nervousness, uneasiness, angst were expressed with such effulgent words..... aggrandizement-amplification, caricature and crock, understatement- unembellishment, elocution-emphasis, enunciation-inflection, announcement, argument, articulation assertion, asseverational choice of words, commentary and communication, declaration and definition, delivery, diction, elucidation, emphasis, enunciation-execution, explanation, exposition-formulation, idiom, interpretation and intonation about the human predicament is brought out in Being and Nothingness. Being and Nothingness offers a critique of Sigmund Freud's theory of the unconscious, based on the claim that consciousness is essentially self-conscious. Sartre also argues that Freud's theory of repression is internally flawed. According to Sartre, in his clinical work, Freud encountered patients who seemed to embody a particular kind of paradox—they appeared to both know and not know the same thing. In response, Freud postulated the existence of the unconscious, which contains the "truth" of the traumas underlying the patients' behavior. This "truth" is actively repressed, which is made evident by the patients' resistance to its revelation during analysis. Yet what does the resisting if the patients are unaware of what they are repressing? Sartre finds the answer in what Freud calls the "censor". "The only level on which we can locate the refusal of the subject," Sartre writes, "is that of the censor. The Holographic Principle, as conceived in current physics, applies to fields, and perhaps, even to more elementary entities, called strings and branes. These more elementary entities remain hypothetical at the time of this writing. There are many layers between the level of fields and that of neurons. We know that neurons are surrounded by a surface phospholipid membrane, which supports an electrochemical process that is fundamentally necessary to human experience. As the information supported by the membrane surface according to the Holographic Principle is proportional to its area, we should expect to find that the neural correlates of consciousness most clearly associated with those parts of the neuron that have the have a high proportion of neuronal surface area, and, in particular, receptive surface area. Dendrites account for an average of about 90% of the neuron is receptive surface area (Wong, 2002), and so would be expected to be the most important neural structures in information processes. We find this to be true

The Holographic Principle Theory of Mind MARK GERMINE Institute for Psycho science (Wikipedia). Discordant-discrepant, dissonant-hard-line, implacable, incompatible, incongruous- inconsistent, inexorable-inflexible, inharmonious-intransigent, opposed, reluctant, unappeasable, uncompromising, unfriendly some of these expatiations and enumerations might sound, a more generalised interpretations is given by a Vedantin. Look at this arresting passage: The teacher continued to be silent. When addressed a second and third time he said: "I am teaching, but you do not follow. The Self is silence." The undetermined and unthinkable character of the Brahman is a consequence of the absolute's eternal and immutable nature. To concede the existence of a real universe is, from the Vedantin's point of view, to posit the existence of a reality apart from the Brahman. Nor can we simply identify a real universe with the absolute unless we are prepared to compromise its unchanging, absolute status. The criterion of authenticity is immutability. Reality never changes; only that which is less than real can appear to do so. Reality is constant in the midst of change. What this means essentially is that there is change although nothing changes. This impossible situation is reflected in the ultimate impossibility of change itself. That which does not exist prior to its changing and at the end, after it has changed,

must be equally non-existent between these two moments. Although the world of change appears to be real, it cannot be so. Change, according to the Vedantin, presupposes a loss of identity. Reality cannot suffer transformation; if it were to do so, it would become something else and the real would be deprived of its reality. The immortal can never become mortal, nor can the mortal become immortal. The ultimate nature of anything cannot change. Change of any sort is merely apparent (vivarta); the world of change and becoming is a false superimposition (adhyaropa, adhyasa) on the absolute. In cosmic terms, the mistake (bhranti) consists of the supposition that the real Brahman is the unreal universe and the unreal universe is the real Brahman. In microcosmic terms, it is the mistake of falsely conceiving the body, mind or even one's personality to be the Self. In the same way as the image of a snake is falsely superimposed on a rope, similarly the universe is falsely projected onto the real substratum, the Brahman. Ignorance is not merely a personal lack of knowledge, but a cosmic principle. As such it is called "Maya," the indefinable factor (anirvacaniya) that brings this mistake in identity about. The reality status of this cosmic illusion is also indefinable: on the one hand it is not Brahman, the sole reality; on the other hand it is not absolutely non-existent like a hare's horn or the son of a barren woman. Brahman is the source of world appearances only in the sense of being their unconditioned ground or essential nature. The universe is false not because it has no nature of its own but because it does have one. Just as the illusion of a snake disappears when one sees that it is nothing but a rope, similarly cancellation (badha) of the empirically real occurs when the absolute reality of the Brahman is realised. Thus, according to Vedanta, appearance implies the real, while the real need not imply appearance. To appear is essentially to appear in place of the real, but to be real is not necessarily to appear. All things -exist because the absolute exists. It is their Being. Thus the very existence of phenomena implies their non-existence as independent realities. When they are known to be as they are, in the fullest sense of their existence, their phenomenal nature disappears leaving the ground of Being naked and accessible. This approach was validated by a critique of experience. The Vedanta established that space, time and the other primary categories of our daily experience can have no absolute existence. It was therefore necessary to make a distinction between relative truths — that accepted by the precritical common man — and an absolute truth discovered at a higher level of consciousness. The Saiva absolutist 17 rejects any theory that maintains that the universe is less than real. From his point of view a doctrine of two truths, one absolute and the other relative, endangers the very foundation of monism. The Kashmiri Saiva approach is integral: everything is given a place in the economy of the whole. It is equally wrong to say that reality is either one or diverse. Those who do so fail to grasp the true nature of things which is neither as well as both. "We do not" says Abhinavagupta, "base our contention that [reality] is one because of the contradictions inherent in saying that it is dual. It is your approach {paksa} that accepts this [method] While, if [duality and oneness] were in fact [to contradict each other], they would clearly be two [distinct realities]." The Vedantin, who maintains that non-duality is the true nature of the absolute by rejecting duality as only provisionally real, is ultimately landed in a dualism between the real and illusory by the foolishness of his own excessive sophistry (vacafadurvidya). Oneness is better understood as the coextensive unity (ekarasa) of both duality and unity. They are equally expressions of the absolute. Gopinath Kaviraj says: According to Sarikara, Brahman is truth and Maya is inexplicable (anirvacaniya). Hence the [Advaitin's] endeavour to demonstrate the superiority of Advaita philosophy is turned against his own system. It tarnishes the picture of its philosophical perfection and profundity. He cannot accept Maya to be a reality, therefore his non-dualism is exclusive. The whole system is based on renunciation and elimination and thus is not all-embracing.... By accepting Maya to be Brahman (brahmamayi), eternal (nitya) and real (satyarupa), Brahman and Maya [in the Tantra] become one and coextensive. **The Doctrine of Vibration: An Analysis of the Doctrines and Practices of Kashmir Shaivism Mark S. G. Dyczkowski. Concept of being from Shaivite point of view:** Solutional behaviour and stability analysis of such systems is of paramount importance, notwithstanding the assumptions made which are not farfetched, in that it throws light on the fact whether they could be candidates for yet unsolved problems, providing Authenticative determination, unimpeachable validation, incontrovertible establishment, apodictic evidence, reliable roll out, unfailing cinquecento **quattrocento trecento**, incontrovertible indication for the wide amplitudinal ramificatory usage of the thesis propounded. The world is now the other person's world, a foreign world that no longer comes from the self, but from the other. The other person is a "threat to the order and arrangement of your whole world...Your world is suddenly haunted by the Other's values, over which you have no control"(Satre: being and Nothingness) Sartre states that many relationships

are created by people's attraction not to another person, but rather how that person makes them feel about themselves by how they look at them. This is a state of emotional alienation whereby a person avoids experiencing **their subjectivity by identifying themselves with "the look" of the other**. The consequence is **conflict**. In order to maintain the person's own being, the person must control the other, but must also control the **freedom of the other "as freedom"**. These relationships are a profound manifestation of "bad faith" as the for-itself is replaced with the other's freedom. The purpose of either participant is not to exist, but to maintain the other participant's looking at them. This system is often mistakenly called "love", but it is, in fact, nothing more than emotional alienation and denial of freedom. *In essence we are always simulating and selling ourselves at the cost of our freedom. Both as a script writer and participant consciousness, human beings experience complete subjective feelings and sometimes absolute subjective feelings, which if remains unchecked might lead to pathological ramifications.* (italics mine)

Perspicuous forbearance, Sophisticated seasoning, Participational observation, Background actuality, Existential worldliness, Existential strife, Predicational anteriorities, Character consonance, Ontological consonance, Primordial exactitude, Phenomenological correlates, Accolytish representation, Atrophied asseveration. Anamensial alienisms, Anchorite aperitif, Arcadian Atticism all delineated in the conditionalities and functionalities and orinetatilities of the system which incorporate the rules and regulations, axiomatic predications and postulation alcovishness of the foregoing state form the bastion, pillar, post, stylobate and sentinel of the classification scheme and doxa. appropriate and diagnostic, differentiating-discriminating, distinctive-distinguishing, emblematic- especial, essentially exclusive, fixed, idiosyncratic-inborn, inbred, indicative and individual, individualistic-inherent, individualizing and ingrained, inherent, innate, local, marked, native, normal, original, particularly- peculiar, personal, private, proper, regular, representative-singular, special, specific, symbolic, symptomatic- unique these form the classification and stratification measure that is suitable notwithstanding the constancy in many a case, as in gravity, which depends on the mass and distance for two given objects. Like Kant and Bergson, Deleuze considers traditional notions of space and time as unifying forms imposed by the subject. He therefore concludes that pure difference is non-spatio-temporal; it is an idea, what Deleuze calls "the virtual". (The coinage refers to Proust's definition of what is constant in both the past and the present: "real without being actual, ideal without being abstract.") Proust, *Le Temps Retrouvé*, ch. III: see the fourth line from the bottom of this page, or, in English translation, the thirteenth paragraph here: "I began to discover the cause by comparing those varying happy impressions which had the common quality of being felt simultaneously at the actual moment and at a distance in time, because of which common quality the noise of the spoon upon the plate, the unevenness of the paving-stones, the taste of the madeleine, imposed the past upon the present and made me hesitate as to which time I was existing in. Of a truth, the being within me which sensed this impression, sensed what it had in common in former days and now, sensed its extra-temporal character, a being which only appeared when through the medium of the identity of present and past, it found itself in the only setting in which it could exist and enjoy the essence of things, that is, outside Time. [...] **Nothing but a moment of the past Much more perhaps; something which being common to the past and the present, is more essential than both. [...] a marvellous expedient of nature had caused a sensation to flash to me—sound of a spoon and of a hammer, uneven paving-stones**—simultaneously in the past which permitted my imagination to grasp it and in the present in which the shock to my senses caused by the noise had effected a contact between the dreams of the imagination and that of which they are habitually deprived, namely, the idea of existence—and thanks to that stratagem had permitted that being within me to secure, to isolate and to render static for the duration of a lightning flash that which it can never wholly grasp, a fraction of Time in its pure essence. When, with such a shudder of happiness, I heard the sound common, at once, to the spoon touching the plate, to the hammer striking the wheel, to the unevenness of the paving-stones in the courtyard of the Guermantes' mansion and the Baptistry of St. Mark's, it was because that being within me can only be nourished on the essence of things and finds in them alone its subsistence and its delight. It languishes in the observation by the senses of the present sterilised by the intelligence awaiting a future constructed by the will out of fragments of the past and the present from which it removes still more reality, keeping that only which serves the narrow human aim of utilitarian purposes. But let a sound, a scent already heard and breathed in the past be heard and breathed anew, simultaneously in the present and in the past, real without being actual, ideal without being abstract, then instantly the permanent and characteristic essence hidden in things is freed and our true being which has for long seemed

dead but was not so in other ways awakes and revives, thanks to this celestial nourishment" While Deleuze's virtual ideas superficially resemble Plato's forms and Kant's ideas of pure reason, they are not originals or models, nor do they transcend possible experience; instead they are the conditions of actual experience, the internal difference in itself. "The concept they [the conditions] form is identical to its object." [Desert Islands, p. 36.] A Deleuzian idea or concept of difference is therefore not a wraith-like abstraction of an experienced thing, it is a real system of differential relations that creates actual spaces, times, and sensations. [See "The Method of Dramatization" in Desert Islands, and "Actual and Virtual" in Dialogues. In fact it not as if the style and substance of science and philosophers have been entirely different. look at the conformality and consonance in the Gaussian Theory and Deleuze's statements on identity and the loss of it in the following passage.....That is, identity is a continuous process that finds its limit in exhaustion. As Deleuze will write, "...the logical relation of causality is inseparable from a physical process of signalling, without which it would not be translated into action. By 'signal' we mean a system with orders of disparate size, endowed with elements of dissymmetry; by 'sign' we mean what happens within such a system, what flashes across the intervals when a communication takes place between disparates. The sign is indeed an effect, but an effect with two aspects: in one of these it expresses, qua sign, the productive dissymmetry; in the other it tends to cancel it" (dr, 20). An actualized entity just is such a signal-sign system, remaining in perpetual communication with the virtual multiplicity it actualizes, constituting its own elements and identity. Such is Deleuze's account of what Badiou refers to as the operation of the "count-for-one"... an operation that is immanent to being itself and energetic in character. yet how are we to discern the local nature of ontological situations, the manner in which there is no global, overarching global situation in which all local situations are embedded as if parts of a whole? in part the above passage already responds to this question. if series must be brought into relation to communicate in order for causality to occur, and if the resonance of series is always a matter of chance, of a throw of the dice, like Lacan's objet a where no series enjoys primacy of model over copy, then all we have are local situations without a total situation. Unlike Heidegger's analysis of Dasein where Dasein is always being-in-the-world, Deleuze's simulacra or actualizations are always divergent, enjoying only local relations. Heidegger argues that in order to understand the being of Dasein we must understand the manner in which Dasein is a being-in-the-world. By contrast, Deleuze's concept of multiplicity provides us with the principle of a new structuralism, of a new local ontology, that allows us to understand the immanent organization of a multiplicity without referring it to an embedding global space. as DeLanda so beautifully puts it in relation to the mathematician gauss, "...when gauss began to tap into these differential resources, a curved two-dimension surface was studied using the old Cartesian method: the surface was embedded in three-dimensional space complete with its own fixed set of axes; then, using those axes, coordinates would be assigned to every point of the surface; finally, the geometric links between points determining the form of the surface would be expressed as algebraic relations between the numbers. but gauss realized that the calculus, focusing as it does on infinitesimal points on the surface itself (that is, operating entirely with local information), allowed the study of the surface without any reference to a global embedding space. basically, gauss developed a method to implant the coordinate axes on the surface itself (that is, a method to implant the coordinate axes on the surface itself (that is, a method of 'coordinatizing' the surface) and, once points had been so translated into numbers, to use differential (not algebraic) equations to characterize their relations. as the mathematician and historian Morris Kline observes, by getting rid of the global embedding space and dealing with the surface through its own local properties, 'gauss advanced the totally new concept that a surface is a space in itself'" (intensive science and virtual philosophy, 11-12). according to the old Cartesian method, we can only outline the properties of a space by relativizing it to a global space in terms of which it is then mapped. by contrast, gauss is able to explore a space in terms of its intrinsic metric and organization as a local space, without referencing it to a whole of which it is conceptualized as a part. **The Image of Thought**

"Image of thought "permeates both popular and philosophical discourse. According to this image, thinking naturally gravitates towards truth. Thought is divided easily into categories of **truth and error**. The model for thought comes from the educational institution, in which a master sets a problem and the pupil produces a solution which is either true or false. This image of the subject supposes that there are different faculties, each of which ideally grasps the particular domain of reality to which it is most suited. In philosophy, this conception results in discourses predicated

on the argument that **"Everybody knows..."** the truth of some basic idea. Descartes, for example, appeals to the idea that everyone can at least think and therefore exists. Deleuze points out that philosophy of this type attempts to eliminate all objective presuppositions while maintaining subjective ones. Deleuze maintains, with Artaud, that real thinking is one of the most difficult challenges there is. Thinking requires a confrontation with stupidity, the state of being formlessly human without engaging any real problems. One discovers that the real path to truth is through the production of sense: the creation of a texture for thought that relates it to its object. Sense is the membrane that relates thought to its other. Accordingly, learning is not the memorization of facts but the coordination of thought with a reality. **"As a result, 'learning' always takes place in and through the unconscious, thereby establishing the bond of a profound complicity between nature and mind"**. Deleuze's alternate image of thought is based on difference, which creates a dynamism that traverses individual faculties and conceptions. This thought is fundamentally energetic and signifying: if it produces propositions, these are wholly secondary to its development. Philosophy merges with ontology; ontology merges with univocity of being; analogy has always a theological vision; not a philosophical vision; one becomes adapted to the forms of god; self and world; the univocity of being does not mean that there is one and the same being; on the contrary, beings are multiple and different they are always produced by disjunctive synthesis; and they themselves are disintegrated and disjoint and divergent; membra disjuncta. the univocity of being signifies that that being is a voice that is said and it is said in one and the same "consciousness". Everything about which consciousness is spoken about being is the same for everything for which it is said like gravity; it occurs therefore as an unique event for everything; for everything for which it happens; even tum tan tum; it is the ultimate form for all of the forms; and all these forms are disjointed; it brings about resonance and ramification of its disjunction; the vorticity of being merges with the positive use of the disjunctive synthesis, and this is the highest affirmation of its univocity like gravity; it is the eternal resurrection or a return itself, the affirmation of all chance in a single moment, the unique cast for all throws; a simple rejoin der for Einstein's god does not play dice; one being, one consciousness, for all forms and all times. a single instance for all that exists, a single phantom for all the living; a single voice for every hum of voices or a single silence for all the silences; a single vacuum for all the vacuousness consciousness should not be said without conjuring if consciousness is one unique event in which all the events communicate with each other; univocity refers both to what occurs to what it is said. The attributable to all states of bodies and states of affairs and the expressible of every proposition So univocity of consciousness means the identity of the noematic attribute and that which is expressed linguistically and sense-fully. Univocity. We mean that it does not allow consciousness to be subsisting in a quasi state and but expresses in all pervading reality. (Reference: LOS: Deleuze) Deleuze sometimes spoke of getting up behind an author and "creating a monster", as if his reading methodology somehow consisted of a distortion of the philosopher's or author's thought. Certainly there is a case to be made for this with regard to the process of deterritorialization, where something is wrested from a territory and deterritorialized upon a new territory, like the animal paw that is deterritorialized from the earth and reterritorialized on the branch. however, it seems to me that the more interesting aspect of Deleuze's approach to other philosophers and art is not so much his "monstrous creations" (i seldom find them particularly monstrous), but rather their Gaussian or Riemannian style, where he explores them in terms of their own internal organization and metric, without reducing them to something alien such as history, society, biology (reductivism), or the signifier. What we find in Deleuze's approach to phenomena is a Gaussian technique. for instance, take Deleuze's books on cinema, his book on Francis Bacon, or his study of Sacher-Masoch. In the first instance, Deleuze's carefully separates cinema from narrative and the signifier, studying it in terms of its specific organization pertaining to the production of images. in the case of Bacon, Deleuze doesn't look for an underlying narrative or "meaning", but instead studies the manner in which Bacon composes and organizes his images and lines, both in terms of their production and actuality, so as to liberate a logic of sensation. finally, in approaching Sacher-Masoch, Deleuze allows Sacher-Masoch's novels to speak for themselves in terms of their desire and relation to pain, stalwartly refusing to reduce masochism to the complement of sadism. in each case, we have a local exploration of a "space" of multiplicities that is extremely precise. Sean Bowden the priority of events: Deleuze's logic of sense Sean Bowden, the priority of events: Deleuze's logic of sense, Edinburgh university press, 2011, 296pp., \$40.00 (hbk), isbn 9780748643646 Reviewed by David Scott, Coppin state university and Gaussian spaces and multiplicities a note on Gaussian spaces and multiplicities posted by

larval subjects under uncategorized. What you "see" is not always what you "get." Many people mistakenly take their own visions literally without "seeing through" the various possibilities that are outside their belief system, knowledge or skill base. Deeper reality is not remote in the physical sense but in a psychological sense. The archetypes of the collective unconscious are arrayed behind the scenes of current worldwide conditions, of crisis and confusion. They mirror our own states back at us, whether we perceive them as such or interpret them plausibly or not. The noise of ordinary consciousness and beliefs drowns out the signal. Unconsciousness is the background of our ordinary awareness. Our organism is very much at the center of such effects. The organismic source is our human bodies and the focus of human consciousness. The fantasy principle dethrones reality, but can be dissociative or compensatory. The human mind is a meme-Scape. Pre-conceived concepts vie with structures, concepts with images. Like scientists who ignore assumed truths, we leapfrog over our beliefs and personality deficits, claiming idiosyncratic imagination is literal reality. It couldn't be further from the truth and symbolism is utterly lost. The metaphor that might heal us enslaves us. Perhaps images like the holographic universe have an implicate order. Can we have a sense of the cosmos in the world without projecting myriad fantasies on it that we embrace literally? Has the world become so horrible it is unreasonable to be realistic? We may need to look at our drives and wishes, rather than the fantasy content. Psyche constructs reality. Our experience of so-called reality is always mediated by our image of it. Even if all the contents of the psyche are real, that doesn't mean they are realistic. That psyche is real is still a radical proposition, but psychic politics certainly color the self-image and ideas of everyone. We observe and participate with images. It is not a question of nature or nurture (genes alone or experience alone). Rather, everything is both. We inherit the structures that make our experience what it is. But the structure itself is "empty," and each human culture "fills" it with its own specific adaptations. It is difficult to define an archetype and set boundaries that distinguish it from others. In a hologram each part contains all the information but in lower resolution. Archetypes have this holographic quality. There are patterns within patterns within patterns. Some overlap with others, and some are nested inside others. Archetypal realities, passed on through DNA, are expressed in distinctive neuronal tracts in the brain. They include customs and laws regarding property, incest, marriage, kinship, and social status or roles; myths and legends; beliefs about the supernatural; gambling, adultery, homicide, schizophrenia, and the therapies to deal with them. A mythic and visionary language of immediate experience encompasses themes of deepest, highest, and ultimate concern. Most fantasy-based individuals are at a complete loss to coherently explain their own conventional behavior much less anomalous events and their deep meaning, much less the cultural unconscious or mythological unconscious matrix. But they try, and become utterly entrenched in their belief that they are right about the nature of the world and reality. We have pseudo-memories about our personal lives. Why not more so for our collective life? The subject matter often revolves around catastrophe, creation and the mythopoeic forces of mankind. Ignorant of such dynamics, interpretive mistakes and displaced psychic contents proliferate into errors of fact. Propaganda, media distortions, memes, and disinformation compound the social problem of misapprehension further. Shameless self-promotion by personalities of such ideas leads to cults. They make up myths about the myths of by-gone eras. Roiling unconscious images can be fatally confusing. Thought illusions culminate in projections and projections of mythology. Jung suggested symbols live only as long as they are pregnant with meaning. Philosophy arose from criticism of myth, from discussing and challenging it. In science, we criticize, reject and eliminate theories. At the edge of the abyss of the unknown, new signs and symbols emerge. Credible theories and paradigms must include biology, physics, and neurophysiology. One of the reasons people "see God", or a guru, or anomalies may be because our brains are constructed to see reality through the eyes of others. There are heaps of mirror neurons which are there to make us feel the 'other'. Mirror neurons do for psychology what DNA did for biology. They provide a unifying framework and help explain a host of mental abilities. As in the psychochemical processes of empathy or falling in love, a complex feedback loop sustains a state of mind. But when we empathically transpose ourselves into someone else's position, we expose ourselves to that reality -- cognitively and emotionally. The unconscious complicates empathy, both ways. Mirror neurons might well play a role in bonding, language and self-awareness. Naively, we take too much as self-evident. But 'seeing' does not always 'believe', though many make this error or leap in logic and formulate their choices and future accordingly. Yet, there is only one way to learn what consciousness is. Experience. But we have no satisfactory explanatory edifice for consciousness. Would such a theory release in each of us our own inner knowledge of the

creativity of our own consciousness, and its infinite possibilities? The problem is trying to define a verb, a dynamic, as if it were a noun. But we do recognize the effect of consciousness. It functions to mediate states of consciousness, high and low psychobiological arousal. Consciousness is the subconscious lifted up by the physical body. When the body fails, the consciousness collapses back into the subconscious. All our thoughts come from the subconscious which can see our intentions but not our world. This relates somehow to intention being imaginary and not of the physical frictionized world (King). Gerald Edelman postulates that the flows of information in the brain are mediated through 're-entrant' feedback loops. As evolution provides new cognitive functions, new re-entrant loops are established. Even language itself is an archetype -- a chaotic field of dynamic associations. A subtle net of tropes, grammar, symbols, and meaning, the program language begins in limbic resonance. Some phenomena generate their own language patterns, nomenclature, and internal coherence of meaning and representation. In a holographic universe, even time and space could no longer be viewed as fundamentals. Because concepts such as location break down in a universe in which nothing is truly separate from anything else, time and three-dimensional space would also have to be viewed as projections of this deeper order. At its deeper level reality is a sort of super hologram in which the past, present, and future all exist simultaneously. This suggests that given the proper tools it might even be possible to someday reach into the super holographic level of reality and pluck out scenes from the long-forgotten past. Or not. A fantasy of such penetration or phenomenon inside the head is not the same as that penetration.

Jung in the 21st Century: Synchronicity and science By John Ryan Haule Mind Control Countermeasures. Adversely antagonistic, inexorably irreconcilable, antithetically antipodal, diametrically opposed, repugnantly retrograde, inimically inverse, violatively unsimilar, diversely dissimilar some statements in Vedanta, it is not superfluous or redundant. It is evolutionary. Authenticative determination, unimpeachable validation, incontrovertible establishment, apodictic evidence, reliable roll out, unfailing cinquecento **quattrocento trecento**, incontrovertible indication of the evolutionary process towards understanding is ever present in all, its thematic and discursive form. It is the relentless holding on to a theory this is despicable albeit the theory is not. Look at this paragraph. It is the 'I am the body/mind' belief that gives rise to the 'I am not (e) the world' belief. These two beliefs are (=) co-created."Consciousness projects (e&eb) the appearance of the mind, body and world by taking the shape of thinking, sensing and perceiving." "Attention is Consciousness with an object. When the object vanishes, attention simply remains what it always is, Consciousness." "There is (=) no purpose to (e) meditation. The purpose is already accomplished." "Everything that is experienced is experienced by, through, in and as Consciousness." "The seen cannot be separated from seeing and seeing cannot be separated from (e) Consciousness." "The Reality of any experience is not hidden in the appearance; it is expressed by the appearance." "Once we see that everything is Consciousness... Maya still dances, but it is a dance of love not seduction." There are some truly excellent sections such as a long one taking, as a starting point, the following quotation from the painter Cézanne: "Everything vanishes, falls apart, doesn't it? Nature is always the same but nothing in her that appears to us lasts. Our art must render the thrill of her performance, along with her elements, the appearance of all her changes. It must give us a taste of her Eternity." Rupert begins: "That statement must be one of the clearest and most profound expressions of the nature and purpose of art in our era." And he goes on for some 14 pages to elaborate on this claim, examining the nature of the 'elements' Existence, appearance and Consciousness and their relationship. "This Reality is the support or ground of the appearance. The appearance may be an illusion, but the illusion itself is real. There is an illusion. It has Reality." He refers to the rope-snake metaphor and says that: "We do not see anything new. We see in a new way." "So that we know that nature is real, that there is something present, that there is a reality to it, even if everything that appears to us is insubstantial and fleeting." **The Transparency of Things' by Rupert Spira Book Review by Dennis Waite The following is a review of Rupert's book: 'The Transparency of Things: Contemplating the Nature of Experience'. There is an essay from the book here and the essay on Cézanne may be read at the awakened eye website For models see Mitigation Of Beam-Induced Backgrounds, Multi-Particle Azimuthal Correlations, Search For Extra Dimensions, Multi-Higgs-Boson Cascade, Top Quark Pair (Tt) Production Charge Asymmetry, $\Delta Y \equiv Y_t - Y_{\bar{t}}$, Reflects The Asymmetry In Tt^- Production, Ultra-Relativistic Electrons And Other Letters Of Interest: A Synecdochal Syncretism: Et Lux In Tenebris Lucet: Light Shines In The Darkness Models.** What then is the relation of Higgs boson and consciousness? Functionalist approaches generally assume that conscious experience

appears as a novel property at a critical level of computational complexity. On the surface this would seem to deal with issues 1 and 3, however a conscious threshold has neither been identified nor predicted, and there are no apparent differences in electrophysiological activities between non-conscious and conscious activity. Regarding the nature of experience (why we are not unfeeling "zombies") functionalism offers no testable predictions. Problem 2) of 'binding' in vision and self is often attributed by functionalists to temporal correlation (e.g. coherent 40 Hz), but it is unclear why temporal correlation per se should bind experience without an explanation of experience. As functionalism is based on deterministic computation, it is also unable to account for Penrose's proposed non-computability (4), or free will (5). Something may be missing. To address these issues, various proposals have been suggested in which macroscopic quantum phenomena are connected to the brain's known neural activity. For the problem of unitary binding, Marshall (1989) suggested that coherent quantum states known as Bose-Einstein condensation occurred among neural proteins (c.f. Penrose, 1987; Bohm and Hiley, 1993; Jibu and Yasue, 1995). Pre-conscious to conscious transitions was identified by Stapp (1992) with collapse of a quantum wave function in pre-synaptic axon terminals (c.f. Beck and Eccles, 1992). In another proposal, protein assemblies called microtubules within the brain's neurons are viewed as self-organizing quantum computers ("orchestrated objective reduction - Orch OR" e.g. **Penrose and Hameroff, 1995; Hameroff and Penrose 1996a; 1996b; c.f. Hameroff 1997; 1998a; 1998b; 1998c; 1998d**). **Stability analysis of such theories would certainly provide a gateway for further bringing in the "reality" in space time.** String theory suggests that the graviton particle would not be detected from a particle collision because it would have "jumped" into another dimension. So, if the graviton exists and the theory is correct, there should be a "gap" in the energy profile of all the particles ejected from the collision. To understand this, think of breaking the pack of balls in a game of pool. The total energy of all the balls after the collision will depend on the force with which the cue ball was hit. So, after the collision, the energy of all the balls is measured and added-up to make sure it is the same as the energy of the cue ball. BUT - imagine that there is some energy missing, because one of the pool balls has "gone missing" - into a higher dimension. (POSTED BY THEREALJEFFHAL) **Talbot says** "If the universe is a hologram, in some sense it suggests that there may be two very drastically different levels of reality: the concrete reality that we see when we look at [things]... and at some deep level there's a level of reality where everything dissolves into an ocean of energy that is holographically interconnected.". A theory of everything must encompass all the variations however contradictory it may be. Nor need any theory be sent in to grand limbo of oblivion and hibernation or a differentiated groundless less ness because there are some irregularities which are like audit discrepancies need to be rectified. Although the language of religion and science seem different, both are talking about the central order of nature relative to DNA and information, as is Peirce, Petrus, Bacon, and pharaonic Egypt. This path-ordering, active information suggests that the central pattern or law of our holographic cosmos may be modeled on the biophysical battleground of gene regulation in bacteriophage Lambda, one of nature's most efficient and highly evolved mechanisms identified by experimental praxis. Egyptian texts support that the Lambda lifestyles of lysogeny and lysis can be understood as two paths to two living systems, photosynthesis and its reversal chemiluminescence, the cosmic key for evolution of mind and continuity. The texts support that the Lambda genome is the world-heart of two ways. The viral lifestyle of lysogeny controlled by cI protein results in matter, classical spacetime, our world of projected shadow, photosynthesis, sexual genesis or vertical gene transfer, mind in the human body, and the imperfect continuum. In contrast, the viral lifestyle of lysis controlled by cro protein and lactose metabolism results in transformation to energy, the quantum, chemiluminescence, asexual genesis or HGT, mind in the cosmos, and the perfect continuum. Some scientists believe our cosmos is entangled with another (Chown 2007). On the holographic quantum level, perhaps this is just the competitive entanglement of cI and cro proteins cycling in perpetual motion—an inherent law of nature elucidating the assembly of things. In light of modern experiments, physicist John Wheeler (1988) speculated that the act of conscious observation functions with quantum mechanics, catalyzing a probability or outcome. So, if an observing consciousness has knowledge of the classical path and its magnetic field orientation, an act of measurement or choice might entangle space to reify the holographic quantum paths to a perfect continuum. Peirce guesses that to be drawn into a new system, a particle must have the right mass, the right velocity in the right direction, the right attraction, and it must present itself at the right point (EP 1. 270). Similarly, a finite probability

exists for any particle that approaches the black hole event horizon to bounce back, dependent on the incoming particle's energy, its charge, and its projection of the orbital momentum on the axis of rotation of the black hole (Kuchiev 2003; 2004; 2004a), the same conditions in Egyptian texts (King 2004; 2006). Only an approximate conclusion remains: chance may be knowledge of the cosmos' holographic mode of operation and magnetic fields; habit breaking may result in the transformation of energy by a violation of the second kind; and evolution of mind or energy might be possible through HGT via a cosmic viral code of Lambda-genesis. "Order is simply thought embodied in arrangement;" (CP 6.490). Accordingly, the original meaning of the signs may relate to a viral biophysics of crystallized mind for human evolution via a natural chemical pathway to a perfect continuum. Again, this semiotics of evolvability and order recurs in time with the chaotic dream of reason that projects a pervasive falsification of our perceptual world. As Wallace Stevens envisions in "Connoisseur of Chaos", perhaps "the pensive man may see." **Peircean Process Metaphysics Origin of Science Quantum and Classical Laws Ancient Egyptian Evolutionary Biophysics** Adversely antagonistic, inexorably irreconcilable, antithetically antipodal, diametrically opposed, repugnantly retrograde, inimically inverse, violatively unsimilar, diversely dissimilar are the views of epistemologists in the analysis of *raison d'être* of knowledge. Epistemology is the study of knowledge and those things closely related to it: justification, what it takes for you to be justified, the relation between knowledge and justification, whether you can have any justified beliefs at all, and if so, how you come to know (or justifiably believe) things, how you can use what you know (or justifiably believe) to come to know (or justifiably believe) other things, and whether and why it's valuable to know instead of merely having justified and true beliefs. The literature on epistemology is vast. Here's a very brief summary of some epistemological discussions. Concerning knowledge, many epistemologists think knowledge is justified true belief, where the justification you have is linked to the truth of the matter in the right kind of way, though what this way is a matter of debate; some epistemologists think knowledge can't be analyzed this way. Concerning justification's relation to knowledge, some epistemologists think we don't need to be justified to know, and some think we do need to be. Concerning what it takes to be justified, some epistemologists think that what it takes for you to be justified are only factors internal to the believer (Internalists). Others think it takes an external factor, like reliable or well-functioning cognitive faculties (Externalists). Skeptics argue that we can't have any justified beliefs at all, and many epistemologists reply to the skeptic's arguments. Concerning how we use what we know (justifiably believe) to come to know (justifiably believe) other things, some epistemologists (Foundationalists) argue that there are bedrock propositions that we know (justifiably believe), and we build our knowledge (justified beliefs) on those. Others (Coherentists) argue that there aren't bedrock propositions; rather, a set of beliefs is justified as a whole, and several beliefs can be mutually supporting. Concerning the value of knowledge, some argue that knowledge is intrinsically valuable. Others have argued that knowledge is valuable only because of the role it plays in practical reasoning, and others argue that knowledge isn't more valuable than justified and true belief, but there are other epistemic states such as understanding, that do have value above their proper subparts. **Edited y Matthew McGrath (University of Missouri, Columbia). It is not as if there is an anathema or misnomer in the philosophy or physics. Infact most of the pnysicists have used philosophical ideas towards the end of consummation of their their and their concomitant justification thereof.** The eternal return produces becoming-active. It is sufficient to relate the will to nothingness to the eternal return in order to realize that reactive forces do not return. However far they go, however deep the becoming-reactive of forces, reactive forces will not return. The small, petty, reactive man will not return. Affirmation alone returns, this that can be affirmed alone returns, joy alone returns. Everything that can be denied, everything that is negation, is expelled due to the very movement of the eternal return. We were entitled to dread that the combinations of nihilism and reactivity would eternally return too. The eternal return must be compared to a wheel; yet, the movement of the wheel is endowed with centrifugal powers that drive away the entire negative. Because Being imposes itself on becoming, it expels from itself everything that contradicts affirmation, all forms of nihilism and reactivity: bad conscience, resentment..., we shall witness them only once. [...] The eternal return is the Repetition, but the Repetition that selects, the Repetition that saves. Here is the marvelous secret of a selective and liberating repetition. There is no need to remind the reader that neither the image of a centrifugal movement nor the concept of a negativity-rejecting repetition appears anywhere in Nietzsche's writings, and indeed Deleuze does not refer to any text in support of this interpretation. Further, one could highlight that Nietzsche never formulates the

opposition between active and reactive forces, which constitutes the broader framework of Deleuze's interpretation. For some years, Marco Brusotti has called attention to the fact that Deleuze introduced a dualism that does not exist in Nietzsche's writings. To be sure, the German philosopher describes a certain number of "reactive" phenomena (for example, in the second essay of the *Genealogy of Morality*, § 11, he talks about "reactive affects" [reaktive Affekte], "reactive feelings" [reaktive Gefühlen], reactive men [reaktive Menschen]); but these are nonetheless the result of complex ensembles of configurations of centers of forces that remain in themselves active. Neither the word nor the concept of "reactive forces" ever appears in Nietzsche's philosophy. **After the discovery of the two principles of thermodynamics began a debate about the dissipation of energy and the thermal death of the universe which framed the modern renewal of the debate between the linear and circular conceptions of time.** Scientists such as Thomson, Helmholtz, Clausius, and Boltzmann and-by way of Kant, Hegel and Schopenhauer-philosophers such as Dühring, Hartmann, Engels, Wundt and Nietzsche have tried to address this problem by using the force of scientific argumentation and of philosophical discussion. Whoever believed in an origin and a final end to the motion of the universe (be it in the physical form of the gradual loss of heat, or in the metaphysical form of a final state of the "world process"), relied on the second principle of thermodynamics or on the demonstration of the thesis of Kant's first cosmological antinomy. On the contrary, those who refused to admit a final state to the universe used Schopenhauer's argument of infinity a parte ante-according to which if a final state were possible, it should already have established itself in the infinity of time past-to propose henceforth a number of alternative solutions. Scientists would propose the hypothesis that energy could have re-concentrated after a cosmic conflagration, thus reversing the tendency towards dissipation. Those belonging to the monistic and materialistic tradition relied on the first principle of thermodynamics and on the infinity of matter, space and time, and regarded the universe as an eternal succession of new forms. A certain critical agnosticism was widespread among scientists and philosophers, oftentimes through a reaffirmation of the validity of Kant's antonymic conflict, this movement avoided to take a stand on specifically speculative issues. Other German philosophers, like Otto Caspari, or Johann Carl Friedrich Zöllner, had reintroduced an organicist and pan-psychical conception of the universe, investing atoms with the ability to escape any state of balance. Indeed, it is probably one of Otto Caspari's works, *The Correlation of Things* (*Der Zusammenhang der Dinge. Gesammelte philosophische Aufsätze* (Breslau: Trewendt, 1881)), which awakened Nietzsche's interest for all things cosmological, in that summer of 1881, in Sils-Maria. addressing Schopenhauer and Eduard von Hartmann's mystical pessimism according to which the world is the creation of a stupid and blind essence (which, after having created the world by mistake, comes to the realization that it had made a mistake and strives to return it to nothingness) Caspari stresses that it is nothing short of mystical to imagine that the world may have been borne out of an originary and undifferentiated state. Where would it have drawn the first impulse? But, continues Caspari, even if the world had received this first impulse from some *deus ex machina*, there is no doubt that, in the temporal infinity of past time thus far, it would have either attained the end of the process (but this is impossible because the world would then have ended), or it would be necessarily bound to repeat indefinitely this original mistake, and the entire process that accompanies it. But then, what is the process of the world? We must now take one more step back and understand further the process of the world according to von Hartmann. Hartmann objects that the regressive movement postulated by Schopenhauer is possible only in thought: it remains nothing more than an "ideal postulate" with no real object and which "does not teach us anything about the real process of the world that unfolds in a movement contrary to this backwards movement of thought" (Hartmann, *Philosophie des Unbewussten*, third edition (1871): 772). Hartmann affirms that if unlike Schopenhauer one admits the reality of time and of the world process, one must also admit that the process must be limited in the past and therefore that there must be an absolute beginning. In Hartmann's mind, failure to do so would result in positing the contradictory **concept of an accomplished infinity**: "The infinity that from the point of view of regressive thinking, remains an ideal postulate, which no reality may correspond to, must, for the world, whose process is, on the contrary, a progressive movement, open up to a determinate result; and here the contradiction comes to light" (Hartmann (1871): 772). What really "comes to light" in this passage is the fact that Hartmann does not provide a demonstration but a *petitio principii*. Indeed, the concept of the world process analytically contains the concept of a beginning of the world. In all rigor, it is therefore impossible to demonstrate these concepts with reference to each other. Secondly, Hartmann's view that one is bound to accept the reality of the world process even if one rejects the

ideality of Schopenhauer's time is mistaken. Hartmann believes that if time is real there must be a world process with both an **absolute beginning and an absolute end. Without any justification, Hartmann jumps from Schopenhauer's negated time to oriented time.** Hartmann's view is that the world process leads into a final state absolutely identical to the initial state. However, it follows from this that even as the cosmic adventures of the unconscious come to a close, we are still haunted by the specter of a new will and of another beginning of the world process. This exposes a serious internal flaw of Hartmann's system insofar as it jeopardizes the possibility of a final liberation from existence and suffering. This is why in the last pages of his work, "On the Last Principles," he painstakingly calculates the degree of probability of a reawakening of the volitional faculty of the unconscious. Insofar as the will is entirely free, unconditioned and a-temporal, the possibility of a new volition is left to pure mathematical chance and is therefore $\frac{1}{2}$. Hartmann further stresses that if the will were embedded in time, the probability of the repetition would amount to 1 and the process of the world would be bound to begin again, in an eternal return which would completely preclude the possibility of a final liberation. Fortunately, this is not the case since-according to Hartmann's remarkable logic-the world-process develops through time, but the original will is outside of time. In fact, one may even affirm, along the lines of Hartmann's peculiar theory of probability, that every new beginning gradually reduces the probability of the next beginning: let n be the number of times that the will is realized, the probability of any new realization is $\frac{1}{2} n$. "But it is clear that the probability $\frac{1}{2} n$ diminishes as n increases, in a way that suffices to reassure us in practice" (Hartmann (1869): 663). In his famous work entitled **On the Conservation of Force (1847)**, Hermann von Helmholtz had divided the totality of the energy in the universe between potential energy and kinetic energy and affirmed the reciprocal convertibility of the two. In 1852, William Thomson pointed out that there exists a sub-ensemble within kinetic energy, heat, which, once it has been generated, is no longer entirely convertible into potential energy-or into any other form of kinetic energy. Considering that the (partial) reconversion of heat into labor is possible only in situations that present a disparity in temperature, and that heat tends to pass from warmer to cooler bodies by spreading on an even temperature level through space, Thomson concluded that the universe tends towards a final state where any energetic transformations, every movement and every form of life will cease. Caspari's atoms (which bring to mind those in Leibniz's Monadology) resemble some sort of biological monads, endowed with internal states. For Caspari, every atom obeys the ethical imperative to participate in the conservation of the general organism and its movement does not only follow the simple physical kind of interaction but also an a priori law ensuring. If a balance of forces had been attained at any moment, this moment would still be going on: therefore, it never happened. The present state contradicts this proposition. Supposing that a certain state rigorously identical with the present state had, one day, existed, this supposition is not refuted by the present state. As one of the infinite possibilities, it is necessary that the present state had been given anyway, since until now an infinite period of time has already unfolded. **If equilibrium were possible, it must have occurred;** and if the present state has already taken place, then so too the one that preceded it as well as the one preceding that one. Therefore it has already taken place a second time, a third time and so on. And likewise it shall take place again a second time, a third time. Innumerable times forwards and backwards. This amounts to saying that all becoming occurs within a repetition of an innumerable number of absolutely identical states. [...] The immovability of forces, their equilibrium is a conceivable case, but it has not occurred. As a result the number of possibilities is greater than the number of realities. -The fact that nothing identical recurs may be explained not thanks to chance, but only thanks to an intention infiltrated within the essence of force. Indeed, supposing an enormous amount of cases, the random occurrence of the same combination is more probable than the same combination never recurring. The quantum of force in the universe is determinate and not "infinite": let us beware [hüten wir uns] from such conceptual extravaganza! Therefore the number of situations, modifications, combinations and developments of this force is doubtless enormous and practically "immeasurable," but in any case this number is determinate and not infinite. On the other hand, the time in which the universe exerts its force is infinite. That is to say, that force is eternally identical and eternally active: -until the present instant an infinity has already taken place, that is to say that all possible developments must have already taken place. Consequently, the present development must be a repetition and therefore both this that was born from it and this that shall be born from it and so on both forwards and backwards. Everything has taken place an innumerable number of times because the overall situation of all forces always recurs (FP 11 [202] of 1881). Nietzsche regards the mechanistic

vision as more plausible and less anthropomorphic than organicism. However, faced with the two major **cosmological models of his time, the mechanistic model and the organic model, Nietzsche wishes to return its polymorphous, proteiform, unstructured** and chaotic character to nature of which the perfectly non-theological and non-teleological theory of eternal return is the strongest seal. This is the first of the "new battles" which come to whoever is aware of the consequences of the death of God: take any antropomorphism away from nature. In the preparatory papers, the third book of the Gay Science is entitled "Gedanke eines Gottlosen / Thoughts of a Godless One." Aphorism 109 of this book, which immediately follows the famous aphorism against the shadows of God, **Boltzmann's theory belongs to the third phase of the debate on thermodynamics and cosmology. The publication of Thomson's brief paper "On a Universal Tendency in Nature to the Dissipation of Mechanical Energy" of 1852 signals the beginning of the scientific controversies on the problem of the dissipation of energy and opened the first phase of the debate, which was announced in the Reflexions on the Motor Powers of Fire by Sadi Carnot and whose conclusion is represented by Clausius's recapitulative article on the concept of entropy in 1865.[44] The second phase started in 1867 when, at the forty-first congress of German scientists and doctors, Clausius gave a lecture on "The Second Principle of the Mechanistic Theory of Heat," where he applied the results of his research on thermodynamics to the universe.** It is true that in his famous lecture of 1854, Helmholtz had already presented the cosmic consequences of the second principle, but Clausius' contribution had a strong impact on German culture. This is because in this lecture he robustly rejected the possibility to consider the universe as an eternal and self-renewing circle, an ewiger Kreislauf in which force and matter are in constant transformation, as was heretofore affirmed by the materialism of the scientists and philosophers, and he did so in the name of the second principle of thermodynamics. In this way, the debate on the principles of thermodynamics gained great importance in European Culture starting in 1867. In the two first phases, it is Thomson's mechanism that predicts the thermal death of the universe. In the third phase, on the contrary, the meaning of the term mechanism changes radically.[46] In accordance with the apocalyptic climate of this period dominated by the "rebirth of idealism," the "overcoming of scientific materialism" and the "bankruptcy of science", the mechanistic paradigm which had accompanied the birth of modern science became challenged on **account of the second principle of thermodynamics**. According to the theorem of the quasi-periodicity of the motions of mechanical systems demonstrated by Poincaré as part of the problem of the three bodies (1890), a mechanical system must evolve according to a quasi-periodical movement and consequently it must always return-sooner or later-to the initial state. **An easily established theorem informs** us that a limited world obeying solely the laws of mechanics shall always pass through a state closely similar to its initial state. On the contrary, according to established experimental laws (supposing we grant them absolute value and wish to push their consequences to the end), the universe is directed towards a final state, which once it is attained, it shall not be able to escape. In this final state, which shall be like a sort of death, all material bodies shall be at rest at the same temperature. **Poincaré's theorem seems** therefore incompatible with the second principle of thermodynamics, which predicts a unidirectional movement of all natural phenomena until the whole universe is brought to a total standstill. Wilhelm Ostwald and the entire energeticist school of thought contended that the principles of thermodynamics were fundamentally new, and could not be re-incorporated to traditional physics and that they should serve as a basis for a new science that regards the qualitative diversity of energy and its tendency to degradation as its axioms. Against energeticism and in an effort to bring entropic phenomena back into the theoretical framework of mechanism, Ludwig Boltzmann introduced the concept of probability in physics, not as an instrument of calculation, but as an explicative principle. In Boltzmann's statistical thermodynamics, the increase of entropy assumed by Clausius is re-interpreted as an increase in molecular chaos. As a result, it becomes possible to explain mechanistically the evolution of closed systems endowed with increasing entropic value, without it committing us to granting absolute value to the second principle of thermodynamics. **Moreover, one no longer needs to fear the thermal death of the universe insofar as the state of equilibrium will in principle** never be complete, but rather will be attained only statistically, leaving open the possibility of fluctuations towards less probable states. Boltzmann's critics remarked that this hypothesis involved two paradoxes called the objection of reversibility (Umkehrinwand), and that of repetition (Wiederkehrinwand). I shall only address here the second one since it coincides with the theory of the eternal return. Based on Poincaré's theorem quoted above, Ernst Zermelo objected to Boltzmann that his model of

the universe suggested that after a finite (if admittedly very long) time the system would return to its initial position. In his first response to Zermelo, Boltzmann avoids committing himself directly to cosmological questions and he only observes that, in the case of concrete thermodynamic systems, the time of recurrence may be extremely long. For example, in normal conditions of pressure and temperature, one-centimeter cube of gas requires 10¹⁰ years to reach a molecular configuration identical to the original one! However, following a response by Zermelo, Boltzmann wrote a new article where he outlines a cosmological picture that he will re-use later in his conclusion to his famous Lectures on Gas Theory. In this cosmological picture, Boltzmann considers the universe as a closed system with constant entropy, within which some fluctuations occur, creating islands of negative entropy. Our solar system originates in one of these fluctuations. As Clausius correctly pointed out, the entropy of our solar system increases constantly as the solar system gets closer to the state of chaos and of the thermal death of the rest of the cosmos. However, in other zones of the universe, some new fluctuations and new islands appear, so that thermal death is never generalized. Here we are given a grand cosmic image, in which the solar system and the sparkle of life that was lit on planet earth are only a fluctuation of order from within a dominant entropic tendency. Life, and the order on which it is based are exceptions, transitory forms taking place in the realm of the shapeless, they are islands of the cosmos that will soon be re-absorbed into chaos. According to Poincaré's theorem, our island will have to be reborn, to develop and die innumerable times in a strictly identical fashion. This happened an infinite number of times during the past eternity and it shall take place again an infinite number of times in the eternity to come.

Boltzmann accepts the "paradox" of recurrence—that is the eternal return of the same—as a legitimate consequence of the probabilistic conception of thermodynamics. It may be rejected for ethical reasons, it may be stored away as an abstract speculation or dismissed along with other cosmic fantasies, but it cannot be rejected on the basis of any rigorously scientific viewpoint. **The Eternal Return: Genesis and Interpretation BY PAUL D'IORIO(excerpts).** Nietzsche found the template of a materialistic cosmology based upon the first principle of thermodynamics. During the same year, he could have found a model of an organic solution to the problem of thermal death of the universe as well as a discussion on the conformation of space in Zöllner's book *On the Nature of Comets* (Johann Carl Friedrich Zöllner, *Über die Natur der Kometen. Beiträge zur Geschichte und Theorie der Erkenntnis* (Leipzig, Engelmann, 1872), 299 f. and 313 f.); Assumed Association In Space Time Of Particles Or Men Leads To Detrimental Ramification Be In Terms Of Progress Of "Subject Matter" Or In The Pathological Disequilibrium Of Gratification Deprivation Complex Which Again Has Mental Turbulence In Case Collective Consciousness Approve It (Raaga) Or Pernicious Implications In Case The Theory Is Not Accepted; After All What Is A Theory: Set Of Correlations And Causations Which Are "Tested" Against A Collective Consciousness Background, Which Yields "Results" Which Are Themselves In "Question" Due To The Very Perception And The Simulation Prospective. The Definition Of Quantum Theorists' Terms, Such As Wavefunctions And Matrix Mechanics, Progressed Through Many Stages. For Instance, Erwin Schrödinger Originally Viewed The Electron's Wavefunction As Its Charge Density Smeared Across The Field, Whereas Max Born Reinterpreted It As The Electron's Probability Density Distributed Across The Field. Although The Copenhagen Interpretation Was Originally Most Popular, Quantum Decoherence Has Gained Popularity. Thus The Many-Worlds Interpretation Has Been Gaining Acceptance Moreover, The Strictly Formalist Position, Shunning Interpretation, Has Been Challenged By Proposals For Falsifiable Experiments That Might One Day Distinguish Among Interpretations, As By Measuring An AI Consciousness Or Via Quantum Computing. More Or Less, All Interpretations Of Quantum Mechanics Share Two Qualities: They Interpret Formalism—A Set Of Equations And Principles To Generate Predictions Via Input Of Initial Conditions They Interpret A Phenomenology—A Set Of Observations, Including Those Obtained By Empirical Research And Those Obtained Informally, Such As Humans' Experience Of An Unequivocal World Two Qualities Vary Among Interpretations: Ontology—Claims About What Things, Such As Categories And Entities, Exist In The World Epistemology—Claims About The Possibility, Scope, And Means Toward Relevant Knowledge Of The World. Now How Does A Person In A Crowd Behave? They He Would Due To Herd Mentality; Ok; That Is One Theory; Simulation By Terrestrial Brahman Anti Brahman Will Make Them Behave In A Highly Organised And Coordinated Manner; With Association Of Things And Individuals Already Established Resultant Orientation Of The Actions Hall Be Only Oriented Towards The Figure Of Imagination And Product Of Puerile Prognostication Formally Established By Terrestrial Brahman Anti Brahman

Before Such A Coordinated And Organised Execution Of Exercise Is Conducted; Analogically, If A Simulation Is Going On In A Bunch Of Particles Or Electrons, And If Some Agency Say Celestial Brahman Anti Brahman Agency Is Simulating It, There Shall Be Coordination And Organisation So That You See “Reality” While The Truth Shall Remain Hidden. Result Is The Suppression Of Information, Delineation Of Disinformation, And The Concomitant Development Of Mushrooms Of Concocted And Fabricated Theories To That “Reality”; In The Case Of Terrestrial Brahman Anti Brahman Agency Intention May Be To Bulldoze And Bowdlerize You, Baffle And Befuddle You In To Submission; In The Case Of Celestial Brahman Anti Brahman Agency Only Reason Could Be To Shield The Truth Being Disseminated; But Why?; May Be You Should Never Become God Which You Are Trying To; May Be God Is Really Playing Dice; What Does That Result In To ; You Become A Szeizophreniac Associating Again And Again What You Had Been Until Now Of Particles And Other Concomitants Again And Again Leading To Flawed Inferences. Confusing The Epistemic With The Ontic, Like If One Were To Presume That A General Law Actually “Governs” Outcomes—And That The Statement Of A Regularity Has The Role Of A Causal Mechanism—Is A Category Mistake. In A Broad Sense, Scientific Theory Can Be Viewed As (=) Offering Scientific Realism—Approximately True Description Or Explanation Of The Natural World—Or Might Be Perceived With Antirealism. A Realist Stance Seeks The Epistemic And The Ontic, Whereas An Antirealist Stance Seeks Epistemic But Not The Ontic. In The 20th Century's First Half, Antirealism Was Mainly Logical Positivism, Which Sought To Exclude Unobservable Aspects Of Reality From Scientific Theory. Since The 1950s, Antirealism Is More Modest, Usually Instrumentalism, Permitting Talk Of Unobservable Aspects, But Ultimately Discarding The Very Question Of Realism And Posing Scientific Theory As A Tool To Help Humans Make Predictions, Not To Attain Metaphysical Understanding Of The World. The Instrumentalist View Is Carried By The Famous Quote Of David Mermin, “Shut Up And Calculate”, Often Misattributed To Richard Feynman. Quantum Mechanical Wave Function (Absolutely Squared) Describes The Completed Interference Pattern, It Must Describe An Ensemble Interpretations Of Quantum Mechanics From Wikipedia. Philosophy Of Science, The Distinction Of Knowledge Versus Reality Is Termed Epistemic Versus Ontic. A General Law Is A Regularity Of Outcomes (Epistemic), Whereas A Causal Mechanism May Regulate The Outcomes (Ontic). A Phenomenon Can Receive Interpretation Either Ontic Or Epistemic. For Instance, Indeterminism May Be Attributed To Limitations Of Human Observation And Perception (Epistemic), Or May Be Explained As A Real Existing Maybe Encoded In The Universe (Ontic). Confusing The Epistemic With The Ontic, Like If One Were To Presume That A General Law Actually “Governs” Outcomes—And That The Statement Of A Regularity Has The Role Of A Causal Mechanism—Is A Category Mistake. Interpretations Of Quantum Mechanics From Wikipedia. Reality Is Constant In The Midst Of Change. Note The Conservation Of Individual, Collective And Cosmic General Ledgers Are What Are Being Spoken Of. Not The Changing Transactions, But The Immutable Refractory Zero. (Italics Mine) What This Means Essentially Is That There Is Change Although Nothing Changes. Like There Are Changing Transactions And Still The Assets And Liabilities Will Be Equal. This Impossible Situation Is Reflected (E&Eb) In The Ultimate Impossibility Of Change Itself. That Which Does Not Exist Prior To Its Changing And At The End, After It Has Changed, Must Be Equally Non-Existent Between These Two Moments. Although The World Of Change Appears To Be Real, It Cannot Be So. (Real)Change, According To The Vedantin, Presupposes A Loss Of Identity. Reality Cannot Suffer Transformation If It Were To Do So, It Would Become Something Else And The Real Would Be Deprived Of Its Reality. The Immortal Can Never Become Mortal, Nor Can The Mortal Become Immortal The Doctrine Of Vibration: An Analysis Of The Doctrines And Practices Of Kashmir Shaivism Mark S. G. Dyczkowski. Concept Of Being From Shaivite Point Of View: In His Famous Work Entitled On The Conservation Of Force (1847), Hermann Von Helmholtz Had Divided The Totality Of The Energy In The Universe Between Potential Energy And Kinetic Energy And Affirmed The Reciprocal Convertibility Of The Two. In 1852, William Thomson Pointed Out That There Exists A Sub-Ensemble Within Kinetic Energy, Heat, Which, Once It Has Been Generated, Is No Longer Entirely Convertible Into Potential Energy-Or Into Any Other Form Of Kinetic Energy. Considering That The (Partial) Reconversion Of Heat Into Labor Is Possible Only In Situations That Present A Disparity In Temperature, And That Heat Tends To Pass From Warmer To Cooler Bodies By Spreading On An Even Temperature Level Through Space, Thomson Concluded That The Universe Tends Towards A Final State Where Any Energetic Transformations, Every Movement And Every Form Of Life Will Cease: The Eternal Return: Genesis And

Interpretation BY PAUL D'IORIO: The Ultimate Nature Of Anything Cannot Change. Change Of Any Sort Is Merely Apparent (Vivarta). The World Of Change And Becoming Is A False Super- Imposition (Adhyaropa, Adhyasa) On The Absolute. It Appears So On The Screen Of Individual, Collective Consciousness, Albeit One Cannot Be So Sure Of Cosmic, For There Are Many Reasons That Cosmic Transactions Also Had To Be Subjected To Change By ParaBrahman. In Cosmic Terms, The Mistake (Bhramanti) Consists Of The Supposition That The Real Brahman Is The Unreal Universe And The Unreal Universe Is The Real Brahman. In Microcosmic Terms, It Is The Mistake Of Falsely Conceiving The Body, Mind Or Even One's Personality To Be The Self. Note We Have Taken Self As Witness Consciousness. Notwithstanding There Is Nothing Wrong In The Conception Of Self As A Dynamical Presupposition, In That Individual Consciousness And Collective Consciousness And Even Cosmic Consciousness Evolutes In To ParaBrahman. In The Same Way As The Image Of A Snake Is Falsely Superimposed On A Rope, Similarly The Universe Is Falsely Projected Onto The Real Substratum, The Brahman. Ignorance Is Not Merely A Personal Lack Of Knowledge, But A Cosmic Principle. As Such It Is Called "Maya," The Indefinable Factor (Anirvacaniya) That Brings This Mistake In Identity About. The Reality Status Of This Cosmic Illusion Is Also Indefinable: On The One Hand It Is Not Brahman, The Sole Reality; On The Other Hand It Is Not Absolutely Non-Existent Like A Hare's Horn Or The Son Of A Barren Woman. Note Here The Vedantin Avoids Talking About Zero Which We Have Avidly And Assiduously Stressed About. Nothing Is The Factor That Exists. Also See the Remarks about Satre's Being And Nothingness. Brahman Is The Source Of World Appearances Only In The Sense Of Being Their Unconditioned Ground Or Essential Nature. The Universe Is False Not Because It Has No Nature Of Its Own But Because It Does Have One. The Doctrine Of Vibration: An Analysis Of The Doctrines And Practices Of Kashmir Shaivism Mark S. G. Dyczkowski. Concept Of Being From Shaivite Point Of View: It Is The 'I Am The Body/Mind' Belief That Gives Rise To The 'I Am Not The World' Belief. These Two Beliefs Are (=) Co-Created."Consciousness Projects The Appearance Of The Mind, Body And World By Taking (Eb) The Shape Of Thinking, Sensing And Perceiving." "Attention Is Consciousness With An Object. When The Object Vanishes, Attention Simply Remains What It Always Is Consciousness." "There Is No Purpose To Meditation. The Purpose Is Already Accomplished." "Everything That Is Experienced Is Experienced By, Through, In And As Consciousness." "The Seen Cannot Be Separated From Seeing And Seeing Cannot Be Separated From Consciousness." "The Reality Of Any Experience Is Not Hidden In The Appearance; It Is Expressed By The Appearance The Transparency Of Things' By Rupert Spira Book Review By Dennis Waite The Following Is A Review Of Rupert's Book: 'The Transparency Of Things: Contemplating The Nature Of Experience'. There Is An Essay From The Book Here And The Essay On Cézanne May Be Read At The Awakened Eye Website [Nature Rejects The Naiveté That Seeks Absolute Truth. We Are Beginning To Realize, Individually And Culturally, That "Realities" Are All Human Constructions.](#) The Task Becomes One Of "Catching Ourselves In The Act" Of Creating Our Own "Reality" From The Flow Of Events. Human Truth Is Always An Engagement Of Mind With Experience. The Challenge Of The Therapist In These Times Of Chaotic Change Is To Validate The Concept That We Don't Have To Fear The Collapse Of What We Think We Are. Strange Attractors: Transference, Holography, And An Archetype Burke, J. (2003). **Strange Attractors: Transference, Holography, And An Archetype (Doctoral Dissertation, Pacifica Graduate Institute, 2003).** When we get to the topic and understanding of the Transcendental Logic, the brilliant and baffling Transcendental Deduction (TD) Recall the two movements just discussed, the one from experience to its conditions and the one from the forms of valid inference to the concepts that we must use in all judging (the Categories). This duality led Kant to his famous question of right (quid juris) (A84=B116): with what right do we apply the Categories, which are not acquired from experience, to the contents of experience? (A85=B117). Kant's problem here is not as arcane as it might seem. It reflects an important question: How is it that the world as we experience it conforms to our logic? In briefest form, Kant thought that the trick to showing how it is possible for the Categories to apply to experience is to show that it is necessary that they apply (A97). TD has two sides, though Kant never treats them separately John Marlado, accounts, attributes, and ascribes the philosophy of Nishida Kitarō is lucid, insightful, and deeply informative despite the highly questionable argument that gives the book its structure. In comparison with other, technically more accurate accounts of Nishida's philosophy, it is broader in scope than most but more single-minded in its approach. The analysis is for the most part limited to four books that have been translated into English, out of approximately 16 volumes of philosophical

essays in Japanese, but it aims to capture the contours and most important details of Nishida's entire philosophy. To appreciate the achievement of this account, one would need only to read, in English translation or in the Japanese original, any of the volumes with their streams of repetitive, puzzling expressions and circuitous, often dead- end argumentation. Wilkinson's elucidation of details in the passages he samples makes it easier to understand why Nishida is considered Japan's greatest academic philosopher. A critical look at the frame of the book and some of its own claims reveals where, to my mind, it has gone astray. The argument that frames this book is repeated throughout: Nishida's philosophy is an attempt to articulate, in a characteristically Western philosophical manner, the Zen experience he had as a young man. Nishida first made the attempt by using the conceptual frameworks of Western philosophers, and after they failed him he developed his own, largely by recasting Buddhist insights. In the end, a philosophical articulation of the Zen worldview in Western terms or even in Eastern terms recast by conceptual, Western philosophical methods appeared impossible. What Nishida attempted is incommensurable with Western philosophy. While the author first offers these statements as hypotheses (p. 2), one premise of his argument that there is such a thing as "the Zen world-view" or "the Zen experience" remains unquestioned; and another that Nishida's pivotal experience was the Zen experience is asserted as "beyond question" (p. 151; see also pp. 28, 48, 10, 114). Wilkinson's presentation of Zen in the first chapter is drawn largely from D. T. Suzuki, with help from Suzuki's portage Abe Masao; some comments on time, birth, and death in; and a few other sources. The result is a caricature that imagines Zen as an invariable, unitary experience. In fact, Dogen's teachings on practice as the manifestation of enlightenment do not sit well with D. T. Suzuki's emphasis on a satori experience, and Suzuki does not enlighten us about what Zen in its various practices has historically been. The author would have benefited from acquaintance with other presentations by Zen teachers, such as that of Shunryū Suzuki's *Zen Mind Beginner's Mind*, with historical accounts of Zen traditions, and with contrasts between Dōgen's view of time and Sam Āśāra and that of other Buddhist philosophers. A little research in contemporary Zen scholarship would have shown how implausible is the talk of "nirvana or awareness of mu" (p. 24), "the basic Zen position [that] never varies" (p. 3), "the Zen vision" (pp. 5, 7), "Zen epistemology", "Zen experience" (p. 59), "the conception of time involved in Zen" (p. 15), and "the fundamental Zen assertion that there is no absolute time" (p. 16). Even if Dōgen's notion that time and being are nondual (p. 16) epitomized most of Zen, Nishida's early notion of "a transcendent, unchanging reality apart from time" (p. 51) would contrast sharply with it. Equally questionable is Wilkinson's contention that Nishida's "pure experience" is equivalent to Zen satori. The notion of pure experience Nishida developed in his first major work may be pivotal for his entire philosophy, but he never claims or implies that any part of his philosophy presupposes an exceptional experience like satori that grounds his convictions and renders them unverifiable to those who have not had the experience, as Wilkinson intimates (pp. 55, 159, and 161). Nishida presents his notions of pure experience and intellectual intuition, the grasp of its unity, and later... Perennialism and Constructivism is studied by Randolph T Dible : experiencer or the observer and the transcendental subjectivity itself Pure objectivity is essentially a deconstruction, apophatic and negativa The encountering of the experiencer or observer—transcendental subjectivity itself—at the foundation of the world leads inevitably to the recognition of pure objectivity as ultimate reality (which can be taken as its ultimate deconstruction, analogous to the apophatic or via negativa), from which objects derive their value, weight, significance, meaning or objectivity. In this way, pure objectivity can be seen as the supra-self-evident Axiological Axiom, so to speak, even Unconditional Love, in romantic terms. This axiology (value theory) has a structure inverse to the relationship between transcendental subjectivity as the radical unity of pure self-reference and on the other hand, the world of forms, as mere traces (representations, indications) of the unique, original "first distinction" Spencer-Brown speaks of at the foundation of his calculus. That is, all forms (i.e., distinctions, differences) would reduce to being the first distinction, also known as the marked state, which can be called penultimate reality (pure self-reference or transcendental subjectivity: the Spirit which animates us), except that forms are complementary to their content, which is their objectivity or value, which would reduce to the unmarked state or ultimate reality. It is the incongruity of form (thoughts; Whitehead's "negative prehensions") and value (feelings; Whitehead's "positive prehensions," or my notion of objectivity, meaning and qualia; in short, the non-formal aspects of experience) that holds forms open and keeps them from absolute reduction. This accounts for the brute, concrete persistence of the "functional illusion"-- to use a term from Dzogchen Buddhism-- of the world. Thus this system has an axiology of metaphysical objectivity grounded on the

ideal of pure objectivity as the source of all value, meaning and significance, itself the very fecundity of profundity, which is the motive of drawing the distinction in the first place **(Wikipedia, Kant's writings and Logic Of Sense)**

Nano rings were created by accident while intending to make quantum dots. They have interesting optical properties associated with excitons and the Aharonov–Bohm effect. Application of these rings used as light capacitors or a buffer includes photonic computing and communications technology. Analysis and measurement of geometric phases in mesoscopic rings is ongoing. Several experiments, including some reported in 2012, show Aharonov-Bohm oscillations in charge density wave (CDW) current versus magnetic flux, of dominant period $h/2e$ through CDW rings up to 85 μm in circumference above 77 K. This behavior is similar to that of the superconducting quantum interference devices (seeSQUID). **Aharonov–Bohm nano rings (Wikipedia) Stability analysis of nano rings would help progressively accentuate the research about the materials division that uses nano rings.**

From the profound contemplation and wisdom of the Buddha and Mahabodhisattvas, we know there is no such reality. Instead, egolessness (non-self) is the only path to understand the reality of the deluded life. All existences are subject to the law of causes and conditions. These include the smallest particles, the relationship between the particles, the planets, and the relationship between them, up to and including the whole universe! From the smallest particles to the biggest matter, there exists no absolute independent identity. Egolessness (non-self) implies the void characteristics of all existence. Egolessness (non-self) signifies the non-existence of permanent identity for self and existence (dharma). Sunyata stresses the voidness characteristic of self and existence (dharma). Sunyata and egolessness possess similar attributes. As we have discussed before, we can observe the profound significance of sunyata from the perspective of inter-dependent relationships. Considering dharma-nature and the condition of nirvana, all existences are immaterial and of a void-nature. Then we see each existence as independent of each other. But then we cannot find any material that does exist independent of everything else. So egolessness also implies void-nature! From the observation of all existences, we can infer the theory of nirvana and the complete cessation of all phenomena. From the viewpoint of phenomena, all existences are so different from each other, that they may contradict each other. They are so chaotic. In reality, their existence is illusory and arises from conditional causation. They seem to exist on one hand, and yet do not exist on the other. They seem to be united, but yet they are so different to one another. They seem to exist and yet they do cease! Ultimately everything will return to harmony and complete calmness. This is the nature of all existence. It is the final resting place for all. If we can understand this reality and remove our illusions, we can find this state of harmony and complete calmness.

Teachings in Chinese Buddhism (6) sunyata (emptiness) in the Mahayana context (Wikipedia) Many models have been given concatenating Buddhist doctrines and Vacuum energy. These models essentially throw light on the essence of self(witness consciousness) and the cosmic general ledger (cosmic consciousness) while the imbalance in the gratification deprivation complex always remain unbalanced when it comes to individual general ledger , advocating the evolutionary theory of individual consciousness to cosmic consciousness. It is to be noted that it is not the question of knowing all the transactions but the wisdom of what lies behind the debit- Credit, Debit-Debit, or Credit-Credit principle that is most important for assimilation. Look at this simple thesis to prove the point stated in the foregoing. Thesis: 'The World Has A Beginning In Time, And In Space It Is Also Enclosed In Boundaries.' Proof: 'For If One Assumes That The World Has No Beginning In Time, Then Up To Every Given Moment In Time An Eternity Is Elapsed, And Hence An Infinite Series Of States Of Things In The World, Each Following Another, Has Passed Away. But Now The Infinity Of A Series Consists Precisely In The Fact That It Can Never Be Completed Through A Successive Synthesis. Therefore An Infinitely Elapsed World-Series Is Impossible, So A Beginning Of The World Is A Necessary Condition Of Its Existence, Which Was The First Point To Be Proved (Kant, Critique Of Pure Reason, Tr. By R. Guyer And A. W. Wood (Cambridge: Cambridge University Press, 1998): B 454, P. 470). As Quoted In The Eternal Return: Genesis And Interpretation BY PAUL D'ORIO" The Cyclical Hypothesis, So Heavily Criticized By Nietzsche (VP II 325 And 334), Arises In This Way." In Fact, Nietzsche Was Not Criticizing The Cyclical Hypothesis But Only The Particular Form Of That Hypothesis Presented In Vogt's Work. All Of Nietzsche's Texts Without Exception Speak Of The Eternal Return As The Repetition Of The Same Events Within A Cycle Which Repeats Itself Eternally. If Deleuze's Interpretation Holds That The Eternal Return Is Not A Circle, Then What Is It? A Wheel Moving Centrifugally, Operating A "Creative Selection," "Nietzsche's Secret Is That The Eternal Return Is Selective" Says Deleuze: The Eternal Return

Produces Becoming-Active. It Is Sufficient To Relate The Will To Nothingness To The Eternal Return In Order To Realize That Reactive Forces Do Not Return. However Far They Go, However Deep The Becoming-Reactive Of Forces, Reactive Forces Will Not Return. The Small, Petty, Reactive Man Will Not Return. Affirmation Alone Returns, This That Can Be Affirmed Alone Returns, Joy Alone Returns. Everything That Can Be Denied, Everything That Is Negation, Is Expelled Due To The Very Movement Of The Eternal Return. We Were Entitled To Dread That The Combinations Of Nihilism And Reactivity Would Eternally Return Too. The Eternal Return Must Be Compared To A Wheel; Yet, The Movement Of The Wheel Is Endowed With Centrifugal Powers That Drive Away The Entire Negative. Because Being Imposes Itself On Becoming, It Expels From Itself Everything That Contradicts Affirmation, All Forms Of Nihilism And Reactivity: Bad Conscience, Ressentiment..., We Shall Witness Them Only Once. [...] The Eternal Return Is The Repetition, But **The Repetition That Selects, The Repetition That Saves. Here Is The Marvelous Secret Of A Selective And Liberating Repetition.** There Is No Need To Remind The Reader That Neither The Image Of A Centrifugal Movement Nor The Concept Of A Negativity-Rejecting Repetition Appears Anywhere In Nietzsche's Writings, And Indeed Deleuze Does Not Refer To Any Text In Support Of This Interpretation. Further, One Could Highlight That Nietzsche Never Formulates The Opposition Between Active And **Reactive Forces, Which Constitutes The Broader Framework Of Deleuze's Interpretation. For Some Years, Marco Brusotti Has Called Attention To The Fact That Deleuze Introduced A Dualism That Does Not Exist In Nietzsche's Writings. To Be Sure, The German Philosopher Describes A Certain Number Of "Reactive" Phenomena** (For Example, In The Second Essay Of The Genealogy Of Morality, § 11, He Talks About "Reactive Affects" [Reaktive Affekte], "Reactive Feelings" [Reaktive Gefühlen], Reactive Men [Reaktive Menschen]); But These Are Nonetheless The Result Of Complex Ensembles Of Configurations Of Centers Of Forces That Remain In Themselves Active. Neither The Word Nor The Concept Of "Reactive Forces" Ever Appears In Nietzsche's Philosophy. The Eternal Return: Genesis And Interpretation BY PAUL D'ORIO Some Parts Are Deleted Due To Spatial Constraints. In the midst of thoughts about the eternal return we find at least two other thematic axes On the one hand, the view of the **world as a constant flux of forces without any goal, law, or rules of becoming. A chaos sive natura de-divinized and de-anthropomorphized which constitutes the "ontological substratum"** of the whole of Nietzsche's reflexions On the other hand, an ensemble of fragments of an anthropologico-sociological character, designing a path of liberation leading to the creation of superior individuals by way of a profound transformation of their instinctual structure. This transformation must be achieved by a practice of solitude and internal struggle towards the liberation from the ancient representations of the world and from the incorporated herd values. For an analysis of these thematic perspectives, see Paolo D'Iorio, *La linea e il circolo. Cosmologia e filosofia dell'eterno ritorno in Nietzsche* (Genova: Pantograf, 1995): 233-322. With most of the parameters quantified, with assumptions there must be picturesque view of the self and non self, space and time, science and philosophy with the concatenational consummations. After The Chapter "On Redemption," Where Zarathustra Dares Not Expose His Doctrine, The Eternal Return Begins To Be Enunciated In Part Three Of The Work. In The First Place, It Is The Dwarf Who Formulates It In The Chapter "On The Vision And The Riddle." Facing The "Gate Of The Instant" Which Symbolizes The Two Infinities That Stretch Towards The Past And The Future, The Dwarf Whispers: "All Truth Is Crooked, Time Itself Is A Circle" The Dwarf Represents The Spirit Of Gravity, And He Embodies The Herd Morality, "The Belittling Virtue" Which Is The Title Of Another Chapter From Part III. The Dwarf Can Endure The Eternal Return Without Great Difficulties Because He Has No Aspirations; Unlike Zarathustra He Does Not Wish To Climb The Mountains That Symbolize Elevation And Solitude. In Two Unpublished Notes, From The Summer And The Fall Of 1883, Nietzsche Writes: The Doctrine Is At First Favored By The Rabble, Before It Gets To The Superior Men. The Doctrine Of Recurrence Will First Smile To The Rabble, Which Is Cold And Without Any Strong Internal Need. It Is The Most Ordinary Of Life Instincts, Which Gives Its Agreement First. Hence, The Content Of The Doctrine Is The Same, But Whereas The Dwarf Can Endure It (Because He Interprets It According To The Pessimistic Tradition For Which "Nothing Is New Under The Sun"), Zarathustra, Who Is The "Advocate Of Life" Regards The Eternal Return As The Strongest Objection To Existence, And As The Rest Of The Dream Suggests, He Does Not Yet Succeed In Accepting It After The Vision At The Gate Of The Instant, The Chapter Is Brought To An End By The Enigma Of The Shepherd. The Eternal Return: Genesis And Interpretation BY PAUL D'ORIO .How does the memory affect the GD complex or for that matter the gratification deprivation of an

individual or a collective identity? Non-Markovian local-in-time master equations give a relatively simple way to describe the dynamics of open quantum systems with memory effects. Despite their simple form, there are still many misunderstandings related to the physical applicability and interpretation of these equations. Here, we clarify these issues both in the cases of quantum and classical master equations. Further introduction of the concept of a classical non-Markov chain signified through negative jump rates in the chain configuration. **E-M Laine et al 2012 J. Phys. B: At. Mol. Opt. Phys. 45 154004 doi:10.1088/0953-4075/45/15/154004 Local-in-time master equations with memory effects: applicability and interpretation.** Supposing That There Were Indeed An "Energy Of Contraction" Constant In All Centers Of Force Of The Universe, It Remains To Be Explained Where Any Difference Would Ever Originate. It Would Be Necessary For The Whole To Dissolve Into An Infinite Number Of Perfectly Identical Existential Rings And Spheres, And We Would Therefore Behold Innumerable And Perfectly Identical Worlds COEXISTING [Nietzsche Underlines This Word Twice] Alongside Each Other. Is It Necessary For Me To Admit This? Is It Necessary To Posit An Eternal Coexistence On Top Of The Eternal Succession Of Identical Worlds? The Eternal Return: Genesis And Interpretation BY PAUL D'IORIO. Science is usually portrayed as dehumanizing. Brave New World epitomizes this fear. "The more we understand the world, the more it seems completely pointless" (Steven Weinberg). Certainly science can seem chilling when conceived in the abstract as a metaphysical world-picture. We may seem to find ourselves living in a universe with all the human meaning stripped out: Participants in a soulless dance of molecules, or harmonics of pointlessly wagging superstrings and their braneworld cousins. Nature seems loveless and indifferent to our lives. What rights have we to be happy? (**Brave New World: Aldous Huxley**). Religion can afford to be dogmatic but science cannot. Look at this passage which counters both scientists and philosophers alike: However, Hartmann Objects That The Regressive Movement Postulated By Schopenhauer Is Possible Only In Thought: It Remains Nothing More Than An "Ideal Postulate" With No Real Object And Which "Does Not Teach Us Anything About The Real Process Of The World That Unfolds In A Movement Contrary To This Backwards Movement Of Thought" (Hartmann, Philosophie Des Unbewussten, Third Edition (1871): 772). Hartmann Affirms That If Unlike Schopenhauer One Admits The Reality Of Time And Of The World Process, One Must Also Admit That The Process Must Be Limited In The Past And Therefore That There Must Be An Absolute Beginning. In Hartmann's Mind, Failure To Do So Would Result In Positing The Contradictory Concept Of An Accomplished Infinity: "The Infinity That From The Point Of View Of Regressive Thinking, Remains An Ideal Postulate, Which No Reality May Correspond To, Must, For The World, Whose Process Is, On The Contrary, A Progressive Movement, Open Up To A Determinate Result; And Here The Contradiction Comes To Light" (Hartmann (1871): 772). What Really "Comes To Light" In This Passage Is The Fact That Hartmann Does Not Provide A Demonstration But A Petitio Principii. Indeed, The Concept Of The World Process Analytically Contains The Concept Of A Beginning Of The World. In All Rigors, It Is Therefore Impossible To Demonstrate These Concepts With Reference To Each Other. Secondly, Hartmann's View That One Is Bound To Accept The Reality Of The World Process Even If One Rejects The Ideality Of Schopenhauer's Time Is Mistaken. Hartmann Believes That If Time Is Real There Must Be A World Process With Both An Absolute Beginning And An Absolute End. Without Any Justification, Hartmann Jumps From Schopenhauer's Negated Time To Oriented Time. The Eternal Return: Genesis And Interpretation BY PAUL D'IORIO. In the last years several theoretical papers discussed if time can be an emergent property deriving from) quantum correlations. Here, to provide an insight into how this phenomenon can occur, authors present an experiment that illustrates Page and Wootters' mechanism of "static" time, and Gambini et al. subsequent refinements. A static, entangled state between a clock system and the rest of the universe is perceived as evolving by internal observers that test the correlations between the two subsystems. They implement this mechanism using an entangled state of the polarization of two photons, one of which is used as a clock to gauge the evolution of the second: an "internal" observer that becomes correlated with the clock photon sees the other system evolve, while an "external" observer that only observes global properties of the two photons can prove it is static. Phys. Rev. A 89, 052122 (2014) DOI:10.1103/PhysRevA.89.052122 arXiv: 1310.4691 [quant-ph] **Time from quantum entanglement: an experimental illustration Ekaterina Moreva.** Time is an entanglement. When the new ideas of quantum mechanics spread through science like wildfire in the first half of the 20th century, one of the first things physicists did was to apply them to gravity and general relativity. The results were not pretty. It immediately became clear that these two foundations of modern physics were entirely

incompatible. When physicists attempted to meld the approaches, the resulting equations were bedeviled with infinities making it impossible to make sense of the results. Then in the mid-1960s, there was a breakthrough. The physicists John Wheeler and Bryce DeWitt successfully combined the previously incompatible ideas in a key result that has since become known as the Wheeler-DeWitt equation. This is important because it avoids the troublesome infinities—a huge advance. But it didn't take physicists long to realise that while the Wheeler-DeWitt equation solved one significant problem, it introduced another. The new problem was that time played no role in this equation. In effect, it says that nothing ever happens in the universe, a prediction that is clearly at odds with the observational evidence. This conundrum, which physicists call **'the problem of time'**, has proved to be thorn in flesh of modern physicists, who have tried to ignore it but with little success. Then in 1983, the theorists Don Page and William Wootters came up with a novel solution based on the quantum phenomenon of entanglement. This is the exotic property in which two quantum particles share the same existence, even though they are physically separated. Entanglement is a deep and powerful link and Page and Wootters showed how it can be used (e) to measure time. Their idea was that the way pair of entangled particles evolves is a kind of clock that can be used to measure change. But the results depend on how the observation is made. One way to do this is to compare the change in the entangled particles with an external clock that is entirely independent of the universe. This is equivalent to god-like observer outside the universe measuring the evolution of the particles using an external clock. In this case, Page and Wootters showed that the particles would appear entirely unchanging—that time would not exist in this scenario. But there is another way to do it that gives a different result. This is for an observer inside the universe to compare the evolution of the particles with the rest of the universe. In this case, the internal observer would see a change and this difference in the evolution of entangled particles compared with everything else is an important a measure of time. This is an elegant and powerful idea. It suggests that time is an emergent phenomenon that comes about because of the nature of entanglement. And it exists only for observers inside the universe. **Time is an entanglement (Wikipedia)**. Danish religious philosopher Soren Kierkegaard carried out a systematic critique of the pretensions of reason and an abstract rationalism which he believed that the modern age was nurturing. Condemning reflection as a "danger" that ensnares people in logical delays and machinations, Kierkegaard compared it to a prison. Reflection is for him a form of captivity, a bondage which "can only be broken by [passionate] religious inwardness" (1978: 81). **Reflection seduces individuals into thinking its possibilities are "much more magnificent than a paltry decision"** (1978: 82). It leads them to act **"on principle,"** to dwell on the deliberation of the context of their actions and the calculation of their worth or outcome. Kierkegaard argues that this drives away feeling, inspiration, and spontaneity, all of which are crucial for true inner being and a vital relation to God. For Kierkegaard, as Nietzsche would later agree, genuine inner being (and culture) is characterized by the tautness and tension of the soul which characterizes passionate existence. But the "coiled springs of life relationships ... lose their resilience" in reflection (1978: 78) and "everything becomes meaningless externality, devoid of [internal] character" (1978: 62). Kierkegaard thus contributes to the development of an irrationalist tradition that has echoes in some later postmodern thought. Kierkegaard might well have agreed with his contemporary Fyodor Dostoyevsky, who wrote: "An intelligent [reflective] man cannot seriously become anything ... excessive consciousness is a disease" (1974: 3, 5). In an age overtaken by rules and regulations, genuine action -- which **Kierkegaard assumes to be subjective and spontaneous -- is frustrated at every turn**. Complaining that we are too "sober and serious" (1978: 71) even at banquets, Kierkegaard bemoans the fact that even suicides are premeditated (1978: 68)! "That a person stands or falls on his actions is becoming obsolete; instead, everybody sits around and does a brilliant job of bungling through with the aid of some reflection and also by declaring that they all know very well what has to be done" (1978: 73). Thus, it is passion, not reflection, that guarantees "a decent modesty between man and man [and] prevents crude aggressiveness" (1978: 62). "Take away the passion and the propriety also disappears" (1978: 64). The ambiguity in the word "passion" may cause some confusion here. To say that the age and its individuals are "passionless" is not to say there are no emotions whatsoever, but rather that there is no true spiritual inwardness and depth, no intensively motivated action and commitment. It suggests that passion exists only in a simulated, pseudo-form, "the rebirth of passion" through "talkativeness" (1978: 64). "Chattering" for Kierkegaard gets in the way of "essential speaking" and merely "reflects" inconsequential events (1978: 89-99). Hence, in "the present age," emotions -- which in fact are all too pronounced -- have been transformed into negative forces. Anticipating Nietzsche's genealogy of the

"slave revolt" in morality, Kierkegaard claims that the "enthusiasm" of the prior age of Revolution, a "positively unifying principle," has become a vicious "envy," a "negatively unifying principle" (1978: 81), a leveling force in its own right insofar as those lacking in talent and resources want to tear down those who have them. **THE POSTMODERN TURN IN PHILOSOPHY: THEORETICAL PROVOCATIONS AND NORMATIVE DEFICITS** By Steven Best and Douglas Kelner <http://www.gseis.ucla.edu/faculty/kellner/kellner.html>. In November 2011, the United States government argued before the US Supreme Court that it wants to continue utilizing GPS tracking of individuals without first seeking a warrant. In response, Justice Stephen Breyer questioned what this means for a democratic society by referencing Nineteen Eighty-Four. Justice Breyer asked, "If you win this case, then there is nothing to prevent the police or the government from monitoring 24 hours a day the public movement of every citizen of the United States. So if you win, you suddenly produce what sounds like 1984..." [65]. What is to be said in unmistakable terms is that when it comes to national security and innocent citizens there is question of compromise even it be an infringement and encroachment of so called privacy which has been made a word to subterfuge the activities that are far said to be fair. Quantum theory has provoked intense discussions about its interpretation since its pioneer days. One of the few scientists who have been continuously engaged in this development from both physical and philosophical perspectives is Carl Friedrich von Weizsäcker. The questions he posed were and are inspiring for many, including the authors of this contribution. Weizsäcker developed Bohr's view of quantum theory as a theory of knowledge. **Harald Atmanspacher, Hans Primas** show that such an epistemic perspective can be consistently complemented by Einstein's ontically oriented position. **Time, Quantum and Information 2003, pp 301-321 Epistemic and Ontic Quantum Realities Harald Atmanspacher, Hans Primas Robert W. Spekkens** present a toy theory that is based on a simple principle: the number of questions about the physical state of a system that are answered must always be equal to the number that are unanswered in a state of maximal knowledge. Many quantum phenomena are found to have analogues within this toy theory. These include the noncommutativity of measurements, interference, the multiplicity of convex decompositions of a mixed state, the impossibility of discriminating nonorthogonal states, the impossibility of a universal state inverter, the distinction between bipartite and tripartite entanglement, the monogamy of pure entanglement, no cloning, no broadcasting, remote steering, teleportation, entanglement swapping, dense coding, mutually unbiased bases, and many others. The diversity and quality of these analogies is taken as evidence for the view that quantum states are states of **incomplete knowledge** rather than states of reality. A consideration of the phenomena that the toy theory **fails to reproduce, notably, violations of Bell inequalities** and the existence of a Kochen-Specker theorem, provides clues for how to proceed with this research program. DOI: <http://dx.doi.org/10.1103/PhysRevA.75.032110> **Evidence for the epistemic view of quantum states A toy theory Phys Rev A 75, 032110 – Published 19 March 2007 Robert W. Spekkens** In such specialized and autonomous areas, it is to the eminent and erudite authors to interpret the contradistinctions and contradictions' in the dove tailing explaining. This I feel would be yeoman service in the creation of Nature's general ledger. **M. S. Leifer and O. J. E. Maroney** examine the relationship between quantum contextuality (in both the standard Kochen-Specker sense and in the generalized sense proposed by Spekkens) and models of quantum theory in which the quantum state is maximally epistemic. We find that preparation noncontextual models must be maximally epistemic, and these in turn must be Kochen-Specker noncontextual. This implies that the Kochen-Specker theorem is sufficient to establish both the impossibility of maximally epistemic models and the impossibility of preparation noncontextual models. The implication from preparation noncontextual to maximally epistemic then also yields a proof of Bell's theorem from an Einstein-Podolsky-Rosen-like argument. DOI: <http://dx.doi.org/10.1103/PhysRevLett.110.120401> Received 25 August 2012 Published 20 March 2013 © 2013 American Physical Society **Maximally Epistemic Interpretations of the Quantum State and Contextuality Phys. Rev. Lett 110, 120401 – Published 20 March 2013 M. S. Leifer and O. J. E. Maroney.** Epistemological, ontic and ontological aspects particularly of quantum mechanics have invited profound thinkers to **contribute** their mite to the growing literature on the subject, which includes the contributions from the subterranean realm and ceratoid dualism of philosophy like that Chaos theory of Deleuze, Derrida, Kierkegaard and others. **Andrei Khrennikov** shows that the Dirac-von Neumann formalism for quantum mechanics can be obtained as an approximation of classical statistical field theory. This approximation is based on the Taylor expansion (up to terms of the second order) of classical physical variables – maps $f: \Omega \rightarrow \mathbb{R}$, where Ω is the infinite-

dimensional Hilbert space. The space of classical statistical states consists of Gaussian measures ρ on Ω having zero mean value and dispersion $\sigma^2(\rho) \approx \hbar$. This viewpoint to the conventional quantum formalism gives the possibility to create generalized quantum formalisms based on expansions of classical physical variables in the Taylor series up to terms of n th order and considering statistical states ρ having dispersion $\sigma^2(\rho) = \hbar/n$ (for $n = 2$ we obtain the conventional quantum formalism). **December 2005, Volume 18, Issue 7, pp 637-650 Date: 28 Nov 2005**

Generalizations of Quantum Mechanics Induced by Classical Statistical Field Theory Andrei Khrennikov It is widely accepted that consciousness or, in other words, mental activity is in some way correlated to the behavior of the brain or, in other words, material brain activity. Since quantum theory is the most fundamental theory of matter that is currently available, it is a legitimate question to ask whether quantum theory can help us to understand consciousness. Several approaches answering this question affirmatively, proposed in recent decades, will be surveyed. It will be pointed out that they make different epistemological assumptions, refer to different neurophysiological levels of description, and adopt quantum theory in different ways. For each of the approaches discussed, these imply both problematic and promising features which will be indicated. **Discrete Dynamics in Nature and Society Volume 2004 (2004), Issue 1, Pages 51-73 <http://dx.doi.org/10.1155/S102602260440106X>**

Quantum theory and consciousness: an overview with selected examples Harald Atmanspacher Copyright © 2004 Hindawi Publishing Corporation. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Another thought provoking and investigatory article is by Peter G. Lewis et al. Perhaps the quantum state represents information about reality, and not reality directly. Wave function collapse is then possibly no more mysterious than a Bayesian update of a probability distribution given new data. **Peter G. Lewis et al** consider models for quantum systems with measurement outcomes determined by an underlying physical state of the system but where several quantum states are consistent with a single underlying state—i.e., probability distributions for distinct quantum states overlap. Significantly, they demonstrate by example that additional assumptions are always **necessary to rule out** such a model. DOI: <http://dx.doi.org/10.1103/PhysRevLett.109.150404> **Distinct Quantum States Can Be Compatible with a Single State of Reality Phys. Rev. Lett. 109, 150404 – Published 9 October 2012 Peter G. Lewis et al** Communication complexity of a quantum channel is the minimal amount of classical communication required for classically simulating a process of state preparation, transmission through the channel and subsequent measurement. It establishes and perpetuates a limit on the power of quantum communication in terms of classical resources. **Alberto Montina** shows that classical simulations employing a finite amount of communication can be derived from a special class of hidden variable theories where quantum states represent statistical knowledge about the classical state and not an element of reality. This special class has attracted strong interest very recently. The communication cost of each derived simulation is given by the mutual information between the quantum state and the classical state of the parent hidden variable theory. Finally, find that the communication complexity for single qubits is smaller than 1.28 bits. The previous known upper bound was 1.85 bits. DOI: <http://dx.doi.org/10.1103/PhysRevLett.109.110501> **Epistemic View of Quantum States and Communication Complexity of Quantum Channels Phys. Rev. Lett 109, 110501 – Published 12 September 2012 Alberto Montina for models see elsewhere .** Stability analysis of what? Quantum state representing reality or our knowledge of reality? Does the quantum state represent reality or our knowledge of reality? In making this distinction precise, **Nicholas Harrigan, Robert W. Spekkens** are led to a novel classification of hidden variable models of quantum theory. They show that representatives of each class can be found among existing constructions for two-dimensional Hilbert spaces. Approach also provides a fruitful new perspective on arguments for the nonlocality and incompleteness of quantum theory. Specifically, **Nicholas Harrigan, Robert W. Spekkens** show that for models wherein the quantum state has the status of something real, the failure of locality can be established through an argument considerably more straightforward than Bell's theorem. The **historical significance of this result** becomes evident when one recognizes that the same reasoning is present in **Einstein's preferred argument for incompleteness**, which dates back to 1935. This fact suggests that Einstein was seeking not just any completion of quantum theory, but one wherein quantum states are solely representative of our knowledge. Hypothesis is supported by an analysis of Einstein's attempts to clarify his views on quantum theory and the circumstance of his

otherwise puzzling abandonment of an even simpler argument for incompleteness from 1927. **February 2010, Volume 40, Issue 2, pp 125-157 Date: 09 Jan 2010 Einstein, Incompleteness, and the Epistemic View of Quantum States Nicholas Harrigan, Robert W. Spekkens** statistical model of the probabilistic description of physical reality has been studied by many authors. **Andrei Khrennikov** posits a contextual statistical model of the probabilistic description of physical reality. Here contexts (complexes of physical conditions) are considered as basic elements of reality. There is discussed the relation with QM. He proposes a realistic analogue of Bohr's principle of complementarity. In the opposite of the Bohr's principle, **Andrei Khrennikov** principle has no direct relation with mutual exclusivity for observables. To distinguish his principle from the Bohr's principle and to give better characterization, he changes the terminology and speaks about supplementarity, instead of complementarity. Supplementarity is based on the interference of probabilities. It has quantitative expression through a coefficient which can be easily calculated from experimental statistical data. There is need not appeal to the Hilbert space formalism and noncommutativity of operators representing observables. Moreover, in our model there exists a pair of supplementary observables which cannot be represented in the complex Hilbert space. There are discussed applications of the principle of supplementarity outside quantum physics. **October 2005, Volume 35, Issue 10, pp 1655-1693 The Principle of Supplementarity: A Contextual Probabilistic Viewpoint to Complementarity, the Interference of Probabilities and Incompatibility of Variables in Quantum Mechanics Andrei Khrennikov Foundations of Physics** Description of the vacuum in Yang-Mills theory and arrive at a physical interpretation of the pseudoparticle solution and the attendant violation of symmetries has been done by **R. Jackiw and C. Rebbi**. It is here vacuum stability forms the bastion, pillar, post and stylobate of the mechanisms of Yang-Mills Theory and action of a particle. The existence of topologically inequivalent classical gauge fields gives rise to a family of quantum mechanical vacua, parametrized by a CP-nonconserving angle. **The requirement of vacuum stability against gauge transformations renders the vacua chirally noninvariant.** DOI: <http://dx.doi.org/10.1103/PhysRevLett.37.172> **Vacuum Periodicity in a Yang-Mills Quantum Theory Phys. Rev. Lett 37, 172 – Published 19 July 1976 R. Jackiw and C. Rebbi.** Three-dimensional Yang-Mills and gravity theories augmented by gauge-invariant mass terms are analyzed. These topologically nontrivial additions profoundly alter the particle content of the models and lead to quantization of a dimensionless mass-coupling-constant ratio. The vector field excitations become massive, with spin 1 (rather than massless with spin 0), and the mass provides an infrared cutoff. The gravitation acquires mass, mediates finite-range interactions, and has spin 2 (rather than being absent altogether); although its mass term is of third derivative order, there are no ghosts or acausalities. DOI: <http://dx.doi.org/10.1103/PhysRevLett.48.975> © 1982 The American Physical Society **Three-Dimensional Massive Gauge Theories Phys. Rev. Lett 48, 975 – Published 12 April 1982 S. Deser et al** All scattering amplitudes in the maximally supersymmetric N=4 super-Yang-Mills theory possess a new, dual superconformal symmetry which extends the previously discovered dual conformal symmetry of MHV amplitudes. To reveal this property we formulate the scattering amplitudes as functions on the appropriate dual superspace. Rewritten in this form, all tree-level MHV and next-to-MHV amplitudes exhibit manifest dual superconformal symmetry. We propose a new, compact and Lorentz covariant formula for the tree-level NMHV amplitudes for arbitrary numbers and types of external particles. The dual superconformal symmetry is broken at loop level by infrared divergences. However, we provide evidence that the dual conformal anomaly of the MHV and NMHV superamplitudes is the same and, therefore, their ratio is dual conformally invariant. We show this explicitly for the six-particle amplitudes at one loop. Authors conjecture that these properties hold for all, MHV and non-MHV, superamplitudes in N=4 SYM both at weak and at strong coupling. **Nuclear Physics B Volume 828, Issues 1–2, 21 March 2010, Pages 317–374 Dual superconformal symmetry of scattering amplitudes in N=4 super-Yang-Mills theory J.M. Drummond DOI: 10.1016/j.nuclphysb.2009.11.022.** By comparison with numerical results in the **maximal Abelian projection of lattice Yang-Mills theory**, it is argued that the nonperturbative dynamics of Yang-Mills theory can be described by a set of fields that take their values in the coset space $SU(2)/U(1)$. The Yang-Mills connection is parameterized in a special way to separate the **dependence on the coset field**. The coset field is then regarded as a collective variable, and a method to obtain its effective action is developed. It is argued that the physical excitations of the effective action may be knot solitons. A procedure to calculate the mass scale of **knot solitons** is discussed for lattice gauge theories in the maximal Abelian projection. The approach is extended to the $SU(N)$ Yang-Mills theory. A relation

between the large N limit and the monopole dominance is pointed out. **Physics Letters B Volume 458, Issues 2–3, 8 July 1999, Pages 322–330 An effective action for monopoles and knot solitons in Yang–Mills theory Sergei V. Shabanov DOI: 10.1016/S0370-2693(99)00612-7.** It is here the quintessential formulation of consciousness occurs. Like a man standing on the threshold of infinity trying to ponder what lies beyond the veil, which separates the seen from unseen a true seeker tries to lift the individual general ledger to the cosmic general ledger. Like in Darwinian evolution, mind also has to evolve from **individual consciousness to cosmic consciousness**. In Kaishmiri Shaivism one withdraws from the finite to the infinite, but one also goes on an outward journey from the infinite to the finite, because both the finite and the infinite have an intimate connection. The finite is not seen as unreal, but as a symbol of the infinite. There is no real distinction between them. Those two movements constitute Spanda, a key concept in Kaishmiri Shaivism. Spanda is the pulsation of the Absolute in different phases of being. There are no opposites like subject and object, unity or duality, absolute or relative. They are just different phases of the universal vibration of the Absolute. The goal is to realize or be at once infinite and finite. One does not turn away from appearances (like in Advaita Vedanta), but one realizes that the Absolute manifests all things. Spanda, the eternal pulsation of the Absolute, oscillates between a passion to create and dispassion from the created. Through it the Absolute transform itself into all things and then returns back into the emptiness of its undifferentiated nature. In Kaishmiri Shaivism the Absolute is seen as pure consciousness (=being). The Absolute is an eternal all-pervasive principle, the highest reality, the nature of all entities eternally and blissfully at rest within its own nature. The Absolute is the nature of the Self (and thus of us all). The Absolute is divine, it is Shiva, the Lord **of the Universe. It is full of conscious activity through which it generates the universe, and reabsorbs it into itself at the end of each cycle of creation. Thus we speak of monism, as everything resides within this one absolute consciousness**. It sustains all things, it embraces all things, and it pervades all things. All things are appearances within the absolute consciousness, but nevertheless real (in contrast to Advaita Vedanta where appearances are seen as unreal or illusionary). All things appear external (out there, outside ourselves), but they do not have a being on their own. They do not exist as separate entities on their own. Everything is contained within consciousness. What we see as objects are manifestations of consciousness. The event which constitutes the universe is always internal events happening within consciousness because their essential nature is consciousness itself. If a physical object were totally material, and independent or external to consciousness, it could never be experienced. The universe and consciousness are two aspects of a whole. The universe is an attribute of consciousness which bears consciousness as its substance. Doctrine of Vibration: S.G.Dyczkowski What you "see" is not always what you "get." Many people mistakenly take their own visions literally without "seeing through" the various possibilities that are outside their belief system, knowledge or skill base. Deeper reality is not remote in the physical sense but in a psychological sense. The archetypes of the collective unconscious are arrayed behind the scenes of current worldwide conditions, of crisis and confusion. They mirror our own states back at us, whether we perceive them as such or interpret them plausibly or not. The noise of ordinary consciousness and beliefs drowns out the signal. Unconsciousness is the background of our ordinary awareness. Our organism is very much at the center of such effects. The organismic source is our human bodies and the focus of human consciousness. The fantasy principle dethrones reality, but can be dissociative or compensatory. The human mind is a meme-Scape. Pre-conceived concepts vie with structures, concepts with images. Like scientists who ignore assumed truths, we leapfrog over our beliefs and personality deficits, claiming idiosyncratic imagination is literal reality. It couldn't be further from the truth and symbolism is utterly lost. The metaphor that might heal us enslaves us. Perhaps images like the holographic universe have an implicate order. Can we have a sense of the cosmos in the world without projecting myriad fantasies on it that we embrace literally? Has the world become so horrible it is unreasonable to be realistic? We may need to look at our drives and wishes, rather than the fantasy content. Psyche constructs reality. Our experience of so-called reality is always mediated by our image of it. Even if all the contents of the psyche are real, that doesn't mean they are realistic. That psyche is real is still a radical proposition, but psychic politics certainly color the self-image and ideas of everyone. We observe and participate with images It is not a question of nature or nurture (genes alone or experience alone). Rather, everything is both. We inherit the structures that make our experience what it is. But the structure itself is "empty," and each human culture "fills" it with its own specific adaptations. It is difficult to define an archetype and set boundaries that distinguish it from others. In a hologram each part contains all the information

but in lower resolution. Archetypes have this holographic quality. There are patterns within patterns within patterns. Some overlap with others, and some are nested inside others. Archetypal realities, passed on through DNA, are expressed in distinctive neuronal tracts in the brain. They include customs and laws regarding property, incest, marriage, kinship, and social status or roles; myths and legends; beliefs about the supernatural; gambling, adultery, homicide, schizophrenia, and the therapies to deal with them. A mythic and visionary language of immediate experience encompasses themes of deepest, highest, and ultimate concern. Most fantasy-based individuals are at a complete loss to coherently explain their own conventional behavior much less anomalous events and their deep meaning, much less the cultural unconscious or mythological unconscious matrix. But they try, and become utterly entrenched in their belief that they are right about the nature of the world and reality. We have pseudo-memories about our personal lives. Why not more so for our collective life? The subject matter often revolves around catastrophe, creation and the mythopoeic forces of mankind. Ignorant of such dynamics, interpretive mistakes and displaced psychic contents proliferate into errors of fact. Propaganda, media distortions, memes, and disinformation compound the social problem of misapprehension further. Shameless self-promotion by personalities of such ideas leads to cults. They make up myths about the myths of by-gone eras. Roiling unconscious images can be fatally confusing. Thought illusions culminate in projections and projections of mythology. Jung suggested symbols live only as long as they are pregnant with meaning. Philosophy arose from criticism of myth, from discussing and challenging it. In science, we criticize, reject and eliminate theories. At the edge of the abyss of the unknown, new signs and symbols emerge. Credible theories and paradigms must include biology, physics, and neurophysiology. One of the reasons people "see God", or a guru, or anomalies may be because our brains are constructed to see reality through the eyes of others. There are heaps of mirror neurons which are there to make us feel the 'other'. Mirror neurons do for psychology what DNA did for biology. They provide a unifying framework and help explain a host of mental abilities. As in the psychochemical processes of empathy or falling in love, a complex feedback loop sustains a state of mind. But when we empathically transpose ourselves into someone else's position, we expose ourselves to that reality -- cognitively and emotionally. The unconscious complicates empathy, both ways. Mirror neurons might well play a role in bonding, language and self-awareness. Naively, we take too much as self-evident. But 'seeing' does not always 'believe', though many make this error or leap in logic and formulate their choices and future accordingly. Yet, there is only one way to learn what consciousness is. Experience. But we have no satisfactory explanatory edifice for consciousness. Would such a theory release in each of us our own inner knowledge of the creativity of our own consciousness, and its infinite possibilities? The problem is trying to define a verb, a dynamic, as if it were a noun. But we do recognize the effect of consciousness. It functions to mediate states of consciousness, high and low psychobiological arousal. Consciousness is the subconscious lifted up by the physical body. When the body fails, the consciousness collapses back into the subconscious. All our thoughts come from the subconscious which can see our intentions but not our world. This relates somehow to intention being imaginary and not of the physical frictionized world (King). Gerald Edelman postulates that the flows of information in the brain are mediated through 're-entrant' feedback loops. As evolution provides new cognitive functions, new re-entrant loops are established. Even language itself is an archetype -- a chaotic field of dynamic associations. A subtle net of tropes, grammar, symbols, and meaning, the program language begins in limbic resonance. Some phenomena generate their own language patterns, nomenclature, and internal coherence of meaning and representation. In a holographic universe, even time and space could no longer be viewed as fundamentals. Because concepts such as location break down in a universe in which nothing is truly separate from anything else, time and three-dimensional space would also have to be viewed as projections of this deeper order. At its deeper level reality is a sort of super hologram in which the past, present, and future all exist simultaneously. This suggests that given the proper tools it might even be possible to someday reach into the super holographic level of reality and pluck out scenes from the long-forgotten past. Or not. A fantasy of such penetration or phenomenon inside the head is not the same as that penetration. **Jung in the 21st Century: Synchronicity and science By John Ryan Haule Mind Control Countermeasures.** The lower domain of experience is what we do not directly observe, the quantum realm, while the higher level is what we ordinarily observe, the classical realm. Based purely on experiment, the formulation of quantum theory initially placed the great divide between the quantum and classical domains between the measuring apparatus and the particle (Herbert, 1985). However, in the early 1930s, John von Neumann, in his

rigorous mathematical treatise on quantum mechanics, found no support for such a division, compelling him to conclude that the wave function was collapsed by consciousness. Von Neumann's treatise has been called the "quantum bible" and "the most influential book on quantum theory ever written" (Herbert, 1985). The von Neumann formulation leads us to conclude that the lower realms of quantum reality form a virtually limitless array of potentials that are inherently incapable of realization without the observation of a knowing entity, which we identify with consciousness (Kafatos and Nadeau, 1990). We thus have a correspondence between quantum theory, the hierarchical structure implied by the Holographic Principle, and the higher orders of experience, leading to the full expression of the Conscious Universe. David Bohm's theory of the implicate and explicate orders involves a holographic Principle that is fully consistent with the Holographic Principle discussed here (Bohm 1980; 1986; Bohm and Hiley, 1993). According to Bohm, there is an implicate order that represents the universal, holographic subtext of reality, and which unfolds in every moment to produce the explicate order that we all observe. Thomas Germinario (2004) has equated the implicate order with the unconscious process, and the explicate order with conscious process. He emphasizes the nature of dreams within the implicate order, and the importance of the dream work for maintaining a healthy mind through integration of the implicate order and our daily lives. Allan Combs and Mark Holland (1990) connected the implicate order or holomovement (Bohm, 1980) with Carl Jung's theory of synchronicity (Jung and Pauli, 1955), with the implicate order providing a holographic medium through which apparently disconnected individuals become connected. The principle of synchronicity, the instantaneous connection of people and events beyond the senses, has been equated with the quantum-physical principle of non-locality (Combs and Holland, 1990; Germinario, 1991), and has been proposed to be the fundamental mechanism of conscious process (Germinario, 1991). Applying the Holographic Principle theory of mind, subjects may be connected synchronously in the manner of two individuals conversing through cellular phones, where the radio signals are transmitted through a distant satellite. The difference here is that the radio signals travel at the speed of light, and thus there is some minuscule time lapse between the sending and receiving ends. The Holographic Principle implies a much more distant and instantaneous communication pathway, with the analogue of the satellite being the holographic boundary of the Universe, as well as a much richer communication, potentially involving a transmission of thoughts and feelings. Our ego-consciousness seems to mask the universal relatedness implied by the Holographic Principle, and it is perhaps only through transcendence of the ego-consciousness that the higher orders of experience can become conscious. In the phenomenon of synchronicity, there seems to be a meaningful connection between individuals that breaks through the barrier of ego-consciousness. Such a connection is reported by many individuals in the course of dreams, when the ego-consciousness has been suspended, at times informing the dreamer of something that has happened in the life of a meaningfully-connected individual. **The Holographic Principle Theory of Mind MARK GERMINE Institute for Psycho science** Impassioned-intense Connectionist neuroscience, in an effort to attain relevance to clinical practice, has recently given us a theory of the relationship of the unconscious to consciousness which is dualistic (Viamontes and Beitman, 2007), an ad hoc theory of conclusions based on assumptions, the most erroneous of which are that representations of information are information, and that such representations elaborate themselves through a process of **representations of representations in neuronal circuits (Bechtel and Abrahamsen, 1991)**. As theory, the connectionist models are lacking heuristic value and **are inimical to the progress** of basic and clinical neuroscience. This is not to denigrate or minimize the importance of neuronal connections, but these connections are rather like the wires in a radio that plays a symphony. The radio, in and of itself, is incapable of playing the symphony. The radio doesn't do not write the symphony, nor does it broadcast the Passionate-raging symphony over the airways. Yet, if we loosen one wire, the symphony is not manifested through the radio. The brain is not subject to the same vulnerability, in that it has **parallel and redundant networks of processing**, and, it is in this sense that the connectivity of the brain becomes important. Connectionism, to the extent that it accepts consciousness and the unconscious at all, attributes them to separate groups of "circuits" in the brain, with consciousness-processing circuits having a limited but detailed ability to analyze information, and **unconscious-processing processing** more information in less detail (Viamontes and Beitman, 2007). There is an element of truth in this view in that information that has importance and meaning resonates within the brain at higher levels of experience and at higher levels of holographic recursion. This resonance involves what we know, what we feel, and what we are capable and willing to consciously realize. The

Universe, as William Blake noted, can be seen in a grain of sand. Indeed, the grain of sand requires the Universe to exist. The human organism requires the Self to exist. The Self is not a simple amalgam of sensory experiences, as suggested in the connectionist model (Blinder, 2007). This amalgam view of the Self can be damaging when applied to psychotherapy, as the connectionists suggest (Blinder, 2007). The Self is ours forever and ever, from eternity to eternity. As Milton suggested, the mind is its own place. Our deepest purpose is for our minds to resonate with the supra-conscious levels of experience, much as it is purpose of the radio to play the symphony. If the symphony is not to our liking, we can turn the radio off. So, perhaps, the transpersonal consciousness can turn our own individual and collective radios off, or perhaps turn the volume down, if we are not in harmony with the Universal Mind. Physics has given us many enigmas. Quantum theory itself is an enigma. But some of the enigmas seem to involve the limitation of our own abilities of discernment. The thought experiment of "Schrödinger's Cat," as it is often interpreted (e.g. Gribbin, 1984), assumes that the cat is not conscious, and is incapable of observation in the context of collapse of the wave function. This is due, in large part, to the false dichotomy of information and experience. Our observation of other animals indicates, to those that are attuned to the animal mind, who can "feel" the animal mind, that, at the very least, all mammals are conscious. It would seem that the great neuropsychiatrist, Stanley Cobb (1948), may have been correct in his attribution of levels of consciousness to a wide variety of organisms: "...lower animals with no cerebrum appear to be conscious...even plants such as the sunflower that turns towards strong light may have a vague awareness or warmth and comfort. There are many degrees of consciousness and it is my contention that it is integrated at many levels like other important functions of the central nervous system." With respect to the theory of mind, recent mainstream thinking remains classical and mechanistic. Such views of mind are advocated by our most prominent neuroscientists (e.g. Changeux, 1985; Gazzaniga, 1985) as neuroscience research attempts to explain all mental function on the basis of pure mechanism and the localized function of specific areas of the brain. Francis Crick, the distinguished Nobel Laureate, had a very influential second career as a neuroscientist. He concluded that we are basically soulless creatures, that all mental processes could be reduced to identifiable, neural correlates in the brain, and that, in fact, we are nothing but a collection of neurons (Crick, 1995). Crick even went so far as to say that what we call the soul is a group of neurons located in the prefrontal lobes of the brain. Another Nobel Laureate, Gerald Edelman (1987), has written a critically acclaimed book on "neural Darwinism," which purports that the brain is "circuitry" is constructed through a developmental process of "survival of the fittest" neurons and neuronal connections. At the same time the total purposelessness and lack of direction in evolution is expounded, again to critical acclaim, by Richard Dawkins (1986). The idea that there essentially no self or soul imbued with agency and self determination was expounded by Gilbert Ryle (1949/2002) with his parody of the "ghost in the machine," and this work continues to be influential among many scholars of the mind. Philosophers such as Patricia Churchland (1986), who coined the term "neurophilosophy," take an "eliminative materialist" view of belief, free will, and consciousness. Other philosophers, such as Daniel Dennett (1991) would make experience "epiphenomenal," a by-product of brain processes with no effect whatsoever on the function of mind. However, unlike neuroscience, which is still in its infancy, the philosophy of mind has had a long history, including such notable figures as Plato, Aristotle, Descartes, Kant, Locke, Hume, Kierkegaard, and James, to name just a few. Perhaps, as noted by Ervin Laszlo (1974): "in today's world, most of the traditional functions of cognitive synthesis have atrophied and are ignored and neglected." Hopefully, what we have presented here will be a step in the right direction for what Laszlo calls a "conceptual synthesis" to help fill the need for meaningful engagement is such a world. The fundamental elements of such a synthesis have been described by Laszlo as follows: Conceptual synthesis performs at least five basic functions in the guidance of affairs. They are the mystical, the cosmological, the sociological, the pedagogical or psychological, and the editorial functions. The mystical function inspires in man a sense of mystery and profound meaning related to the universe and of himself in it. The cosmological function forms images of the universe in accord with local knowledge and experience, enabling men to describe and identify the structure of the universe and the forces of nature. The sociological function validates supports and enforces social order, representing it in accord with the nature of the universe, or as the natural or right form of social organization. The pedagogical or psychological function guides individuals through stages of life, teaching ways of understanding themselves and others and presenting desirable responses to life is challenges and trials. Finally, the editorial function of conceptual is to define

some aspects of reality as important and credible and hence to be attended to, and other aspects unworthy of serious attention. **The Holographic Principle Theory of Mind MARK GERMINE Institute for Psycho science** Kant sums up this inversion and its spirit early in the Critique of Pure Reason: Up to now it has been assumed that all our cognition must conform to the objects; but all attempts to find something about them a priori through concepts that would extend our cognition have, on this presupposition, come to nothing. Hence let us once try whether we do not get farther with the problems of metaphysics by assuming that the objects must conform to our cognition, which would agree better with the requested possibility of an a priori cognition of them, which is to establish something about objects before they are given to us. This would be just like the first thoughts of Copernicus, who, when he did not make good progress in the explanation of the celestial motions if he assumed that the entire celestial host revolves around the observer, tried to see if he might not have greater success if he made the observer revolve and left the stars at rest. (B xvi) Onticology, like all variations of object-oriented ontology, is realist in its orientation. In defending a realist ontology onticology holds that the vast majorities of objects, actants, beings, or entities are independent of humans and are what they are regardless of whether any humans regard them or register them. In short, onticology rejects any anthropomorphic, idealist, or anti-realist thesis to the effect that to be is to be the correlate of mind, spirit, the body, the human, and language or otherwise. While it is certainly the case that knowledge is necessarily dependent on the object to which it relates, the converse does not hold true. Objects are not dependent on being known, regarded, perceived, or spoken about. As such, and to put it in Aristotelean terms, knowledge is an accident of objects, not objects an accident of knowledge. As Althusser so nicely puts it, “[n]o doubt there is a relation between thought-about-the-real and thisreal, but it is a relation of knowledge, a relation of adequacy or inadequacy of knowledge, not a real relation, meaning by this a relation inscribed in that real of which the thought is the (adequate or inadequate) knowledge” (Reading Capital, 96). Althusser goes on to remark that “[t]he distinction between a relation of knowledge and a relation of the real is a fundamental one: if we did not respect it we should fall irreversibly into either speculative or empiricist idealism” (ibid.). Onticology categorically endorses Althusser’s verdict. It is a fundamental necessity to distinguish between those relations that belong to the object and those that belong to knowledge. Contemporary philosophy continuously confuses these two very different sorts of relations. Naturally the question arises of how it is possible to surmount our relation to the object so as to determine whether objects themselves possess the properties we encounter in relating to objects. In other words, given that we can only ever relate to the object in relating to the object how is it possible to surmount this relation to get at the being of the object itself? Much more will have to be said about this later— and the answers will be surprising with respect to standard prejudices about realism —however, for the moment it can be said that onticology takes its epistemological inspiration from the transcendental realism of Roy Bhaskar. Among other things, Bhaskar sought to provide a transcendental grounding for the sciences. Insofar as onticology defends the thesis that the field of being is much more vast than the field of objects investigated by the natural sciences, it parts way with the thesis that the domain of being is exhausted by the domain of natural objects. However, the general form of Bhaskar’s argument holds for our realist purposes. A transcendental argument seeks to elucidate the conditions under which certain acknowledged practices and forms of cognition are possible. Kant, for example, asked what must be the case for mathematical judgments to be possible. How is it both that we are able to extend our knowledge, as if by magic, through mathematical judgments and, more significantly, that these judgments are able to provide genuine knowledge of the world despite the fact that these forms of reasoning are not based on experience? Part of Kant’s argument consisted in claiming that mind imposes the forms of space and time on the data of experience. In other words, space and time are not attributes of being itself but rather of the mind that regards being. Insofar, Kant argues, as mathematics is ultimately a rumination on the nature of space and time taken in their most abstract form and insofar as the mind imposes space and time on the manifold of sensation, it thus follows that a priori judgments about the nature of spatio-temporal relations are possible that anticipate the structure of actual-space times without directly experiencing these space-times. Why? Because any manifold of sensation must necessarily be structured by these forms imposed by intuition. **Onticology– A Manifesto for Object-Oriented Ontology Part I Posted by larval subjects under Object-Oriented Philosophy My Question Is This: You Say, “Granted There Are Deeper And Deeper And More Primitive Forms Of “Knowledge” Or Abstraction Into Mechanism, But I Wouldn’t Say That The Atomic Level Brings A New Epistemic World Into View.” But Doesn’t Any Kind Of**

Knowledge, However Primitive, Imply Bringing Some Kind Of Epistemic World Into View Or Proto-View?

Joel: David, Yes Exactly. My Question Is, Just How Far Down The Complexity Gradient Can The Meaning Of ‘Knowledge,’ Or ‘Semiotics, Or Specifically Representation (Maya) Be Stretched Without Breaking? I Can Comfortably Stretch It To The Genetic Level, As The Proto-Epistemic, Where Genetic Codes ‘Represent’ Phenotypic Structures Or Modes, And The Whole Of Evolution Learns From Its Mistakes And Successes As That Knowledge And History Is Encoded Into And As The Recapitulation Of The Embryogenesis Of The Organism. But An Atom, In My Model, While Certainly Reacting To Its Environment From Its Own Attractor, Essence, And ‘Interiority’ (Prehension) Isn’t Making Representations Of It To Do So. But The Key To The Power And ‘Essence’ Of Representation (And Simultaneously Its Limitation) Is The Capacity Of Abstraction And Choice. Evolution Does This At A Rudimentary Level (And Anthropomorphizing A Bit) With Its Sending Into The World Its ‘Random’ Variations, Organismic Ideas, In A Sense. Its ‘Mind’ Is Literally On The Outside As Our Living World And Biosphere. A ‘Choice’ Is Made In The Biospheric Interiority, And Knowledge For The Evolutionary/Embryogenetic Trajectory Is Gained By The “Differential Reproductive Success” Of These Individuals Lives Through Time, As The Increasing Intelligence, Representation, And Capacity For Choice Is Directly Injected Into The Flow Through Sexual Selection. But Where Is The Encoded/Abstracted (Enfolded) Representation Of Options And A Choice Among Them Made At The Atomic Level? I Am Working With The Most Cutting-Edge And Coherent Models For The Atom (Including My Own) That I Can Find Among The Heterodoxy, And I See No Evidence Of Representation And Choice At This Level, Or Rather No Real Way To Further Stretch The Meaning Of The Terms To This Level. Sure There Is Uncertainty From Infinite Difference And Immanent Causation (See Bohm On Infinite Causation), But Without Representation There Really Is No Choice (Although Crudely We Could Call A ‘Bifurcation’ A ‘Choice’ By The System As A Whole). So I Am Simply Saying That Representation At The Atomic Level Is Entirely Enfolded And These Are Really Just Ontic Level Phenomena. The Atomic Level, In My View, Is Pre- Not Yet Proto-Epistemic, Which I Place At The Genetic Level Where We Find The First Codes And Primitive Representation. This Just Means That There Is A Real Distinction Between Ontic And Epistemic (And A Gradient Between Them) And We Can’t Collapse The Two Into Just The Ontic-Epistemic. And Yet, With The Distinction Intact, And The Gradient To Explore, They Remain Nondual, Univocal, ONE Or Ontic. What I Would Say Exists At All Levels (My Own Self-Aware Myth About The Given) Is What I Call The “Symbiogenesis Of Subject And Object”. This Is The Spinozan Essence Of Dynamic Stability And Growth, And Leibnizian Prehension. Also Called The “Nucleation Of Observability” In Spinbitz. It Is From This Nucleus Of The Observer (Fuller) In A “Point-Free Geometry” (Whitehead) Of Pointless “Points Of View” As Conditions Of Boundary That Wilber’s “All Is Perspective” Finds Its Ontic Grounding. And This Symbiogenesis Is Also A Rudimentary Or Deep-Level Native Or Primitive Intelligence, Evolution At The Involution Of The Subject-Object Interface. It Is An Exploration Of Creative Learning And Direct Immanent Awareness, Unmediated (Enfolded) By Representational Forms (Unfolded). It Just Is, In Waves (Cosmic Vertebrae) Before It Abstracts, Represents And Incarnates The Flesh Of What Could Be. Matter Is Enfolded Seed Of The Flower Of Its Abstraction. Maya Is This Depth And Recursion Of Brahma Into And Through The Boundary Conditions Of Abstract Relation. And Brahma Is The Span Of The Emergent Rigor Of The Cosmic Ergodic Spine That Opens The Recursions Into The Field Of Infinite Difference. In Immanent-Transcendent Waves, Literally And Empirically A Limit-Cycle In Ergodicity, It Enfolds And Unfolds From Anatom To Anatomy, From Infinite Intelligence Or Omnirelational Awareness Of Self, Through The Representational Depths Of Self As Self-Consciousness. Omni-Evolution, Trans-Dynamics, And The Anatom Posted On February 21, 2013A Discussion With Tom Huston, Part II The Following Is Part 2 Of An Excerpt From A Discussion Initiated By Tom Huston On His Facebook Page, Edited And Embellished For Clarity. Find Part One, Here. In This Part We Dig Into The Nature Of The “Anatom” And The “Ergodic Spine” Of An Electro-Fractal Cosmos, And Into The Trans-Dynamic Integration Between Being And Becoming, And The X-Interface—The Crossroads Of The Ontic-Epistemic (Brahma And Maya) And The Subject-Object Polarities. Univocal_Dynamics_V1 Tom Huston Is A Founding Member Of Integral Institute And A Former Editor Of Enlighten Next Magazine. His Webpage Is Tomhuston.Com. David Marshall Is A Writer And Editor Living In Chicago. No meaning, value, or normativity to be found in nature, that there is nothing **natural beings ought to be**, but that, rather, these judgments **arise from us** (See last post of larval subjects) which has led some to raise valuable

questions about the coherence of these claims. The problem arises when the following three propositions are taken together: 1. there is nothing outside of nature. 2. **Beings have no intrinsic meaning, purpose, or value** (in and of themselves, there's nothing they ought to be). 3. Value judgments about what beings and being ought to be arise from us and beings like us (bonobo apes, dolphins, institutions, birds of paradise, etc). The problem arises between proposition 1 and 3. How can it both be true that there is nothing outside of being and those normative judgments belong to us and other beings capable of making normative judgments, not nature? The problem arises from restricting these judgments to humans and beings capable of making these judgments. In making such a claim it seems as if we're saying that there's something outside of nature, something that is beyond nature, thereby **violating** the first thesis and potentially **reintroducing the nature/culture distinction**. By contrast, if we say that normative judgments are the special domain of those living beings with the proper degree of sentience to make such judgments, then we seem to **reintroduce the nature/culture distinction** and fall back into the sorts of problems that I outlined in my last post (and that are so nicely critiqued by thinkers such as Latour). Is there a way out of this? I don't know.

Great_Chain_of_Being_2_ (lighter) ASIDE: It seems to me that this is really what the debate between realism and anti-realism, realism and socio-linguistic constructivism surrounding the new materialisms and speculative realists has really been all about. It's very easy to treat this as an abstract, academic debate: "Are you a realist or are you an anti-realist?", as if it were just a matter of what happens to be true. But it seems to me that this debate has, in reality, always been **about politics**. As I outlined in my last post, we have perpetually seen how appeals to the real and natural have been used in the name of oppressive power, inscribing both the exploitation of nature and the oppression of various people **in the very fabric of being itself**. **Theistic theology and realist ontology have perpetually been used in the name of what Deleuze called "State Philosophies" or philosophies that ontologize** contingent orders of power and privilege (e.g., "the great chain of being" used to justify patriarchy, monarchy, serfdom, poverty, etc, and appeals to nature used to justify poverty and racial inequality (The Bell Curve), patriarchy (evolutionary psychology), heteronormativity, capitalism, etc). Because arrangements of power and inequality are always contingent in the sense that there's no marked difference in the capacities of peoples, power always looks for a **transcendent supplement** that would provide justification through ontological necessity. Antirealism— from the Greek atomists to present —became the radical and emancipatory gesture because it revealed the lie behind all of these forms of social organization or their inherent contingency or arbitrariness. Realism, by contrast, has all too often functioned as an **apologetics for arbitrary power and social organizations**. Here it's worth recalling what Foucault said about science: "...Even before we know to what extent something like Marxism or psychoanalysis is analogous to a scientific practice in its day-to-day operations, in its rules of construction, in the concepts it uses, we should be asking the question, asking ourselves about the aspiration to power that is inherent in the claim to being a science. The question or questions that have to be asked are: "What types of knowledge are you **trying to disqualify** when you say that you are a science? What speaking subject, what discursive subject, what subject of experience and knowledge are you trying to minorize when you say 'I speak this discourse, I am speaking a scientific discourse, and I am a scientist.' What theoretico-political vanguard are you trying to put on the throne in order to detach it from the massive, circulating, and discontinuous forms that knowledge can take?" (Society Must Be Defended, 10). All of these questions hold equally for claims to something being real. What is one trying to minorize when claiming something is real? What becomes privileged? What is excluded? It is these questions that have been at the heart of the realism debates, for as Spencer-Brown taught us, **every distinction has a marked and unmarked space**, draws attention to something to be included and pushes something into the unconscious or the domain of the invisible, hidden, or veiled. This is above all the case with evocations of the real. However, as I've tried to show antirealism leads to its own problems. First, so long as we exclude **real beings from our ontological inventory, we are unable to fully understand how power functions** ("such and such a set of cultural formations have been thoroughly debunked, yet people still live as if they believed them"). Not only do we not fully understand the sources of the problems due to too much focus on the discursive and semiotic, but we deny ourselves valuable sites of political intervention at the level of infrastructure. Second, we are prevented from addressing things such as cultural racism, such as that found in Heidegger's privileging of the West and the Greeks and Germans in particular. We do a good job addressing biological and theological racism, heteronormativity, and sexism by showing how it is a cultural construction, but when faced with racism such as Heidegger's where he argues that there's **something**

“unique” about the Greek event, it’s language, and about the German language, or Badiou/Zizek’s racism with respect to the “Pauline Event”, we really have no response. Here someone like Jared Diamond or Fernand Braudel is needed to explain global-geographical inequalities. Third, the tools of the cultural turn really do not provide us with the means of thinking the ecological as a site of the political. For this reason, I’ve tried to formulate a third way— which might be called “**constructivist realism**” or “**constructivist naturalism**”—that retains the insights of the cultural turn, while also allowing a robust place for the material. I don’t claim to be original in this. I think that many such as Manuel DeLanda, Deleuze and Guattari, Stacy Alaimo, Karen Barad, Jane Bennett, etc., are up to something similar. It’s a vast project that requires the work of a multitude of voices, especially given the way in which the **culturalism pervades humanities**. **The Relational and the Non-Relational: Notes towards an Immanent and Pluralist Theory of Meaning Posted by larval subjects under uncategorized k-bigpic I have deleted or reformulated some sentences for which I beg pardon attributable to spatial constraints** What is correlation, what is causation? What is coordination? What is coincidence? These thoughts have bothered me since long. I have been able to excavate some reliable material from Wikipedia which I would like to share. “All views are partial and contingent – that’s the lesson **of pluralism**, and you say as much yourself when you say that we must be attentive to ways our own knowledge might contain superstitions. But it’s not discourse that determines whether one view works and another doesn’t – it’s a confrontation with **non-discursive (non-human)** agencies.” I wonder if part of the issue here is that we understand pluralism differently. For me this doesn’t sound like pluralism at all. Rather, it just sounds like our **epistemological condition**. We [hopefully] come to understand those portions of existence we question or investigate. The reason we come to investigate them is largely contingent. Finally, the accounts of these features of existence we give can be mistaken. All of this is perfectly consistent with a **monism**. I’m not sure why what he outlines above is the lesson of pluralism. It seems to me that every realist knows this. It seems to me that ontological pluralism is something quite different. Ontological pluralism is the thesis that there are many different worlds inhabited by many different entities. Thus, for example, you would have one world; say that of Lucretius, that’s only inhabited by atoms and their combinations. You would have another world; say the world of the Mongolian shaman, that’s inhabited by spirits that do all sorts of things. The ontological pluralist is saying that these spirits are, that they exist, that they’re real. This is something quite different than merely saying that there are partial and contingent points of view on the world. It’s this saying that these entities are rather than that some person or groups of people believe that they are that is the nub of the issue. Now, I think part of the issue here is that there’s an ambiguity in the term ontology. Ontology can be one of two things. On the one hand, ontology is a group or persons set of beliefs as to what is. Here it’s trivially true that there are pluralities of ontologies and the realist readily recognizes this. This is the whole reason there are debates over ontology. **Mongolian shamans** have their ontology, Europeans theirs, Christian fundamentalists theirs, materialists theirs, etc. When striving to understand and communicate with others it’s vital to understand these ontologies because, as rhetoricians like Burke point out, our **beliefs about what is** are among the things that motivate our action. Despite having never seen bacteria I was my hands because I believe there are bacteria and viruses on door handles and whatnot and don’t want to get sick. My belief about a particular thing existing is what motivates my action. And who knows, perhaps this belief is as superstitious as the belief that the crops failed because God was displeased with my community. On the other hand, ontology is a theory **about what is**. It is making a claim that something exists. This is where the rubber hits the road. The ontological pluralist seems committed to the thesis that every group set of beliefs about what exists is sufficient for granting the existence of those entities. This is what I find objectionable in Latour. Obviously I’m not bothered by Latour’s suggestion that we should take into account the role that nonhumans like speed bumps, rivers, microbes, etc., play in the form that social assemblages take. But this is not all that Latour claims. This is precisely where the philosopher might balk. Unlike the ethnographer, the philosopher is not interested in what people believe exists, but rather philosophers— at least of the realist variant —are trying to figure out what is. In other words, the realist philosopher **begins with the premise** that not all of these beliefs about what is are true. So for the philosopher, recognizing that Mongolian shaman’s believe in the existence of shamans would only be the first step. The next step would consist in determining whether there’s good reason for thinking such entities really do exist, i.e., whether there’s good reason for believing these entities have mind and culture independent reality. Lest readers think that I’m just picking on the supernatural here, we can ask similar questions about strings, subatomic particles, galaxies,

etc., etc., etc. I guess the question really comes down to what exactly we mean by **ontological pluralism**. When we talk about ontological pluralism are we defending the thesis that people have different theories of being? If so, then ontological pluralism is trivially true. If this is what is meant, then I certainly share Jeremy's view that it's valuable to understand the different world's people believe in. Certainly when I was practicing as an analyst I didn't get in ontological debates with my patients and it was necessary to understand their theory of being or their ontology to properly attend to them. Or, when we talk about ontological pluralism are we defending the thesis that all these ontological theories are true and refer to really existing entities? That's quite a different claim and is not one I would defend or endorse. Now someone might object that "in both The Democracy of Objects and Onto-Cartography you defend the thesis that there are **multiple worlds**." This is true. Because I hold that not everything is related I'm led to the conclusion that there are diverse worlds. However, I also hold that however many worlds there might be, these worlds are nonetheless composed solely of material entities. Within the framework I propose I wouldn't suggest that there's one world where there are spirits and another world where there are souls and yet another where there are only material entities. My view is that there aren't spirits or souls in any of these worlds. One of the things I keep hearing in these discussions is that somehow the realist adopts a view from nowhere. I honestly don't understand this criticism. Investigation always occurs somewhere and requires all sorts of mediations involving technologies, experiments, etc. It's that labor of gathering evidence, conducting experiments, using technologies to observe the world, etc., that gradually gives us a body of data that allows us to say there's good reason to believe that such and such a thing exists and has these powers. Another charge seems to be that the realist refuses to recognize that their claims about the world are fallible. I find this charge particularly strange because it's precisely because the realist recognizes the difference between our theories of the world or what we say about the world and the world itself that fallibility is built into the core of his position. Realism doesn't mean one holds they have special access to the world, that they know all truths, or that they have the truth in hand, only that there are truths to be known and that we can be mistaken about things. Circling **Squares Philip** has a post responding to my quandaries about how to mesh realism and pluralism. He writes: Ontologically and metaphysically the idea of realist pluralism is no longer an issue. There are (appropriately) numerous variants but the basic idea that reality is itself pluralistic is well established. The question is political-discursive. It's what Stengers and Latour are getting at with their concepts of diplomacy and cosmopolitics. They grant, first, that all entities exist and, second, that to say that someone's cherished idol (or whatever disputed entity they hold dear) is non-existent is a '**declaration of war**' – 'this means war,' as Stengers often says. They thus shunt onto-political discourse off of the terrain of knowledge/belief in the sense of existence/non-existence. Their basic claim seems to be that 'respect for otherness,' i.e. political pluralism, can only come from granting the entities that others hold dear an ontology, even if you don't 'believe' in them. You are thus permitted to say 'I do not follow that god, he has no hold over me' but you are not permitted to say 'your god is an inane, infantile, non-existent fantasy, grow up.' And it's not just a question of politeness (although there's that too). The point is to grant others' idols and deities an existence – one needn't agree over what that existence entails, over what capacities that entity has or what obligations it impresses upon you as someone in its partial presence but to deny it existence entirely is to 'declare war' – to deny the possibility of civil discourse, of pluralistic co-existence. I believe that it was Richard Rorty who once quipped something like claims to reconcile realism and idealism always seem to end with a triumph of idealism. We don't, in fact, get realism through such approaches, but rather just get a **pervasive anti-realism**. I think this is also the problem with the "non-controversial pluralism" advocated by Stengers and Latour that Phillip defends here. Such pluralism is not realism but is, in fact, a thoroughgoing social constructivism. I think this is the central problem with Latour's argument in Irreductions (these days I regret having ever defended it). In rejecting both Enlightenment critique and what he calls "reduction" he wants to say something like "The Pentacostal really is filled with the Holy Spirit", that for the 19th working scientist **heat really is a fluid** and phlogiston really is what allows things to burn, and that for the Greek lightning really is an expression of Zeus's anger. Latour tells us that we aren't to reduce or explain away the entities posited by another group's "ontology" but are to develop explanations from within that ontology. It might not sound particularly sexy– and it certainly doesn't tell us what is worth thinking –but I can't help but believe that philosophy is the critical and reflective investigation of basic concepts that guide our investigation of the world about us, how we ought to live our lives, and what form of governance might be best. Compare two figures. A scientist might ask,

what causes depression? We can very well imagine a philosopher turning around and asking the scientist, what is **causality**? The scientist presupposes a concept of **causality** in her investigations. She uses this concept in her inquiry. Now she might have a sophisticated concept of causality or she might never have thought much about causality at all, using it in the sort of colloquial and unreflective way that Plato decried when, for example, people like Euthyphro talked about piety. A whole cascade of questions arise when we raise a question like “what is causality?” We can ask whether or not causality exists at all. We can ask how we distinguish between correlation and two events that merely accompany one another from genuine causation. This, for example, was **Hume’s question**. But perhaps most importantly we can ask whether there is only one form of causality or many forms of causality. Is there only one-to-one causation; one cause and one effect? Is there many-to-one causation; or many events conspiring to produce an effect? Is there one-to-many causation; or one event producing a variety of different effects? We can even ask whether causality necessarily moves from past to present or whether there aren’t forms of causality that move from future to past! **January 23, 2014 Pluralism and Realism:** multiple-worlds Over at Struggles Forever, Jeremy Trombley has an interesting post up on “the ontological turn” in anthropology or ethnography. I’ve been meaning to have a discussion with him about this as I think it’s an issue many of us are struggling with. For example, the core project of The Democracy of Objects— a project which I think many have missed —is to somehow reconcile some version of **social constructivism with a realist ontology** capable of making room for ecology (which requires realist and materialist positions as there’s a fact of the matter where global warming is concerned) as well as the role played by objective agencies in social assemblages such as technologies, infrastructure, features of geography, local climates, the growth cycles of plants and animals, waste, etc. Maybe we can try to organize some cross-blog event to discuss these issues. I certainly think they’re close to the heart of Jeremy, Michael of Archive Fire, Arran James, and a host of others. As an aside, I’m beginning to realize how the different sites of the political I’ve been outlining— semiopolitics, thermopolitics, oikopolitics (political economy), geopolitics, eropolitics (the politics of sex and desire), biopolitics, and chronopolitics (and I’m sure there are other political sites!) —are drawing me away from traditional Marxism. Assuming that classical Marxism holds that economics or the conditions and relations of production are determinative of all other sites of the political, the various sites of the political that I’ve been outlining would lead to the conclusion that there is not one determinative base of the political. This would not require committing Marx to flames, but rather of recognizing the phenomenon of overdetermination, or of a variety of different entangled sites of the political. But I digress. First, I find myself wondering what the ontological turn means in ethnography. Is it 1) the investigation of the different ontologies held/proposed by different cultures? E.g., the Aztecs believed that reality was structured in this way, while the Greeks in that way, and the ancient Chinese this way, etc? Or 2) is it an investigation of how real entities— independent of cultural beliefs —influence cultural formations? Or is it a combination of both? A position that I would favor. January 20, 2014 Thoughts on the Social and Political Implications of Correlationism .Lion Mirror Truth be told, as my thought has evolved the issue of correlationism had fallen off the radar for me. Somehow the debate had come to seem too “philosophical” to me, too “scholastic”, too remote from what interests me: understanding why social assemblages are organized as they are, how power functions in social assemblages, and what we might do to address that power and change things. Somehow the question of whether or not we can get out of the correlation between thinking and being just came to seem remote from these sorts of issues. Somehow it seemed too epistemological. **ASIDE:** Numerous discussions over the years have led me to believe that the debate over correlationism is poorly understood (or maybe I just don’t understand it). On countless occasions I’ve heard people say “of course we must relate to things in order to know them.” Well yeah, of course! I don’t think this is what the critic of correlationism is getting at. It seems to me that correlationism is something more robust than the theses that we must relate to something to know it. **Correlationism** instead seems to require the theses that thought and being are indiscernible. Put more concretely, the correlationist is someone who argues that we either a) can never tell whether being is merely a construction of our thought (weak correlationism), or b) who argues that thought actually constructs being (strong correlationism). In other words, correlationism is another name for idealism. One can hold that we must relate to something in order to know it without being a correlationist. As an aside I should also add that I am a correlationist about some things. For example, I think money is something constructed by society and is therefore a strong correlationist when it comes to money. At any rate, for a long time

I'd become rather indifferent to debates about correlationism and philosophies of access. I had learned the lessons of speculative realism— which I could have also learned, I think, from Deleuze and Guattari, the new materialist feminists, actor-network theorists such as Latour –and had moved on. However, occasionally you come across a tone of phrase that pitches something in a different light. In *The Cut of the Real*, Katerina Kolozova writes, ...the political problem of contemporary philosophy identified by the 'new realists' is, in fact, the product of a more fundamental epistemic problem. In his book *After Finitude*, Quentin Meillassoux calls this problem 'correlationism' and identifies it as an essentially post-Kantian legacy, which continues to dominate and limit philosophy. As a matter of fact, correlationism lies at the heart of postmodern theory and consists in the premise that thought can only 'think itself,' that the real is inaccessible to knowledge and human subjectivity, and that there is nothing but discursive constructs that fully determine thinking and that are methodologically accounted for all the way down. (1 – 2) Thought thinking only itself: Thought only encountering itself. In the jargon of postmodern and poststructuralist lingo, this would be the thesis of infinite semiosis, where signs ("thoughts") only ever relate to other signs. Within this framework, discursivity comes to be the hegemonic framework defining all of being. At the level of politics and social theory more generally, if the correlationist thesis is true the consequences are clear: all social phenomena are discursive and all solutions to social and political problems will be discursive. The sole sphere of the political will be the discursive and all questions of politics will be questions of speech-acts and interpretation. The problem here is not that many theorists recognize that the discursive and semiotic plays an important role in the social and the political. It does and I've repeated this tirelessly. The problem is with what happens when thought or the semiotic becomes a hegemon, an "all", foreclosing our ability to recognize other forms of power. What I've wanted to say is that not all power functions discursively. In my last post and elsewhere I spoke of some other forms of politics: **Thermopolitics**: The politics surrounding **energy in the form of calories and fuels such as gasoline** and coal, and how our life and our very bodies are structured by energy dependencies and by being trapped in particular distributive networks that render these forms of energy available. I'm being quite literal when I speak of energy, talking about the effects, for example, of the absence of food in certain educational environments on cognition, for example; and am generally hostile to metaphorical extensions of the concept of energy which I see as erasing the dimension of real materiality. **Geopolitics**: The role that features of natural and built geography such as mountain ranges, rivers, oceans, soil conditions, roads, housing design, etc., play in the form that social relations take and how they impact individual bodies. **Chronopolitics**: The way in which the structurations of time organize what is possible for us. For example, the structuration of the working day, how much we can say and comprehend at any given time, the impact of things like the invention of the clock, etc. **Oikopolitics**: **This would be the domain of political economy described so well by Marxists**. So five different types of politics: Semiopolitics (or what currently dominates critical theory), thermopolitics, geopolitics, chronopolitics, and oikopolitics. No doubt there are other sites of the political or political struggle that we could speak of, but this is a good start. Also, it should be obvious that these aren't exclusive domains, but are entangled in all sorts of important ways. For example, something might take place at the level of semiopolitics (speech, law, rhetoric, norms, and communication) that has all sorts of impact at the level of thermopolitics. Congress might decide to cut programs that fund school meal programs. This, in turn, will have a **thermodynamic impact on** those students that go without the calories they need developmentally and cognitively to function in a particular way. There is an entanglement here of semiopolitical and thermopolitical domains. The young student here has been constrained both at the level of **semiotic phenomena and thermodynamic structures**. The point is that if true, semiotic intervention (speech-acts, protests, interpretations, deconstructions, etc) will not be an appropriate response to all political problems because social formations are not entirely structured by the semiotic. The child in that school does not suffer from a lack of the right signs, but from a lack of calories needed to run the engine of his thought and body. Certainly semiotic interventions might be needed to render that energy available, but it is the energy itself that is at issue and the absence of that energy that forms the spider web entangling him in his position. A correlationist perspective tends to erase this as even being a site of the political. **Larval Subjects January 28, 2014 Ontological Anarché Posted by larval subjects under Uncategorized January 27, 2014 Roden on Pluralism Over at enemy industry Roden has an excellent post up on the pluralism discussion. Check it out here. January 25, 2014 Different Senses of Pluralism and Ontology Posted by larval subjects under Uncategorized Responding to one of my comments**

over at his blog, Jeremy Trombley: I have deleted restricted, reformulated some sentences .Kindly bear with me. Intention is to make the discursive enucleation consistent with the subject in question. Recourse to epistemology of the quantum mechanics generic and fervor of string theory is a sine qua non for the further comprehension of the theory and ontology .How would the world appear to us if its ontology was that of classical mechanics but every agent faced a restriction on how much they could come to know about the classical state? We show that in most respects it would appear to us as quantum. The statistical theory of classical mechanics, which specifies how probability distributions over **phase space evolve** under Hamiltonian evolution and under measurements, is typically called Liouville mechanics, so the theory we explore here is Liouville mechanics with an epistemic restriction. The particular epistemic restriction we posit as our foundational postulate specifies two constraints. The first constraint is a classical analog of Heisenberg's uncertainty principle; the second-order moments of position and momentum defined by the phase-space distribution that characterizes an agent's knowledge are required to satisfy the same constraints as are satisfied by the moments of position and momentum observables for a quantum state. The second constraint is that the distribution should have maximal entropy for the given moments. Starting from this postulate, authors derive the allowed preparations, measurements, and transformations and demonstrate that they are **isomorphic to those allowed in Gaussian quantum mechanics** and generate the same experimental statistics. They argue that this reconstruction of Gaussian quantum mechanics constitutes additional evidence in favor of a research program wherein quantum states are interpreted as states of incomplete knowledge and that the phenomena that do not arise in Gaussian quantum mechanics provide the best clues for how one might reconstruct the full quantum theory. I: <http://dx.doi.org/10.1103/PhysRevA.86.012103> **Reconstruction of Gaussian quantum mechanics from Liouville mechanics with an epistemic restriction Phys. Rev. A 86, 012103 – Published 10 July 2012 Stephen D. Bartlett, Terry Rudolph, and Robert W. Spekkens.** Deleuze in fact characterizes modes of existence, with their powers and capacities. The answer is this: Deleuze approaches modes of existence, ethically speaking, not in terms of their will, or their conscious **decision making power (as in Kant)**, nor in terms of their interests (as in Marx, for example), but rather in terms of their drives. For Deleuze, conscious will and preconscious interest are both subsequent to our unconscious drives, and it is at the level of the drives that we have to aim our ethical analysis. Here, I would like to focus on two sets of texts on the drives taken, not from Nietzsche and Spinoza, but rather from Nietzsche and Leibniz (Leibniz being one of the first philosophers in the history of philosophy to have developed a theory of the unconscious). The first set of texts comes from Nietzsche's great early book entitled Daybreak, published in July 1881. Nietzsche first approaches the question of the drives by giving us an everyday scenario: "Suppose we were in the market place one day," he writes, "and we noticed someone laughing at us as we went by: this event will signify this or that to us according to whether this or that **drive happens at that moment to be at its height in us**—and it will be a quite different event according to the kind of person we are. One person will absorb it like a drop of rain, another will shake it from him like an insect, another will try to pick a quarrel, another will examine his clothing to see if there is anything about it that might give rise to laughter, another will be led to reflect on the nature of laughter as such, another will be glad to have involuntarily augmented the amount of cheerfulness and sunshine in the world—and in each case, a drive has gratified itself, whether it be the drive to annoyance, or to combativeness or to reflection or to benevolence. This drive seized the event as its prey. Why precisely this one? Because, thirsty and hungry, it was lying in wait" (D 119). This is the source of Nietzsche's **doctrine of perspectivism** ("there are no facts, only interpretations"), but what is often overlooked is that, for Nietzsche, it is our drives that interpret the world, that are perspectival—and not our egos, not our conscious opinions. It is not so much that I have a different perspective on the world than you; it is rather that each of us has multiple perspectives on the world because of the multiplicity of our drives—drives that are often contradictory among themselves. "Within ourselves," Nietzsche writes, "**we can be egoistic or altruistic, hard-hearted, magnanimous, just, lenient, insincere, can cause pain or give pleasure**" (Parkes, pp. 291-292). We all contain such "a vast confusion of contradictory drives" (WP 259) that we are, as Nietzsche liked to say, multiplicities, and not unities. Moreover, these drives are in a constant struggle or combat with each other: my drive to smoke and get my nicotine rush is in combat with (but also coexistent with) my drive to quit. This is where Nietzsche first developed his concept of the **will to power**—at the level of the drives. "Every drive is a kind of lust to rule," he writes, "each one has its perspective that it would like to compel all the other drives to accept as a norm" (WP 481). **DELEUZE AND THE QUESTION OF DESIRE:**

TOWARD AN IMMANENT THEORY OF ETHICS Daniel W. Smith. **In on a Possible Physical Metatheory of Consciousness** Miroljub Dugic et al show that the modern quantum mechanics, and particularly the theory of decoherence, allows formulating a sort of a physical metatheory of consciousness. Particularly, the analysis of the necessary conditions for the occurrence of decoherence, along with the hypothesis that consciousness bears (more-or-less) well definable physical origin, leads to a wider physical picture naturally involving consciousness. This can be considered as a sort of a psycho-physical parallelism, but on very wide scales bearing some cosmological relevance. Open Systems & Information Dynamics Volume 9, Number 2, 153 (2002) arXiv: quant-ph/0212128. On a broader perspective and background, the theoretical computation of the universe is justified. IN physics and cosmology, digital physics is a collection of theoretical perspectives based on the premise that the universe is, at heart, describable by information, and is therefore computable. Therefore, according to this theory, the universe can be conceived of as either the output of a deterministic or probabilistic computer program, a vast, digital computation device, or mathematically isomorphic to such a device. Digital physics is grounded in one or more of the following hypotheses; listed in order of decreasing strength. The universe or reality: is essentially informational (although not every informational ontology needs to be digital); is essentially computable (the pancomputationalist position); can be described digitally; is in essence digital; is itself a computer (pancomputationalism); is the output of a simulated reality exercise. Following Jaynes and Weizsäcker, the physicist John Archibald Wheeler wrote the following: [...] it is not unreasonable to imagine that information sits at the core of physics, just as it sits at the core of a computer. (John Archibald Wheeler 1998: 340) It from bit. Otherwise put, every 'it'—every particle, every field of force, even the space-time continuum itself—derives its function, its meaning, its very existence entirely—even if in some contexts indirectly—from the apparatus-elicited answers to yes-or-no questions, binary choices, bits. 'It from bit' symbolizes the idea that every item of the physical world has at bottom—a very deep bottom, in most instances—an immaterial source and explanation; that which we call reality arises in the last analysis from the posing of yes-no questions and the registering of equipment-evoked responses; in short, that all things physical are information-theoretic in origin and that this is a participatory universe. (John Archibald Wheeler 1990: 5) David Chalmers of the Australian National University summarised Wheeler's views as follows: Wheeler (1990) has suggested that information is fundamental to the physics of the universe. According to this 'it from bit' doctrine, the laws of physics can be cast in terms of information, postulating different states that give rise to different effects without actually saying what those states are. It is only their position in an information space that counts. If so, then information is a natural candidate to also play a role in a fundamental theory of consciousness. We are led to a conception of the world on which information is truly fundamental, and on which it has two basic aspects, corresponding to the physical and the phenomenal features of the world. (Wikipedia). **Participatory consciousness forms the bastion, pillar, post, stylobate and fulcrum of the redressal measures and corrective steps taken for circumvention of many problems, identified and unidentified, spread widest commonalty. Anne Baring's sententious and pithy article will be good to have a timely outlook.** I would like to open my contribution to this meeting with the words of Ken Wilber that I quoted in the first Millennium Symposium: 'The secret impulse of life is towards greater consciousness. Maybe the evolutionary sequence is from matter to body, to mind, to soul, to spirit - each transcending and including, each with a greater depth and greater consciousness and wider embrace' Ken Wilber, and with these from Rupert Sheldrake's contribution to the same meeting: "It is possible to make quite a good case that the sun could be conscious and if the sun, then why not the stars? Why not the galaxies? Why should there not be galactic minds? And if they communicate with each other then the distances are far too great for light to be the medium of transmission. We are into inter-galactic telepathy." After many decades of study, I know of only one diagram in all religious traditions that can adequately explain these statements; that can give us both a blue-print of the many levels of consciousness and a route-map to help us reach beyond our present level of understanding. I would like to introduce a diagram of the four worlds of Kabbalah to illustrate the theme that spirit has brought these four worlds into being and that we, living in the fourth and manifest world know nothing of the existence of the other three invisible ones. It is the clearest diagram I know of that can explain to us the multi-levelled structure of consciousness and our place in the great chain of being. We are indeed, as Wilbur suggests, evolving from matter to body, to mind, to soul, to spirit but where have we come from? It is possible that, as participants in the emanation of spirit into manifest life, we have also come from spirit to soul, to mind, to body, to matter. Everything, from the

immense galactic energies to the minute forms of matter participates in one life, is generated from one creative spirit. There is nothing outside this life, this spirit. This diagram suggests that truth is not an article of faith but the experience of participating in the undiscovered reality of what has brought us into being. We cannot know truth until we have entered into communion with that reality. Continuing in similar vein **Michel Bitbol dilates upon the dialectic deliberation and polemical argumentation between** hermeneutists and eliminativists. When he formulated the program of Neurophenomenology, Francisco Varela suggested a balanced methodological dissolution of the “hard problem” of consciousness. He shows that his dissolution is a paradigm which **imposes itself onto seemingly opposite views**, including materialist approaches. I also point out that Varela's revolutionary epistemological ideas are gaining wider acceptance as a side effect of a recent controversy between hermeneutists and eliminativists. Finally, he emphasizes a structural parallel between the science of consciousness and the **distinctive features of quantum mechanics**. This parallel, together with the former convergences, point towards the common origin of the main puzzles of both quantum mechanics and the philosophy of mind: neglect of the constitutive blindspot of objective knowledge. Phenomenology and the Cognitive Sciences Phenomenology and the Cognitive Sciences **Phenomenology and the Cognitive Sciences 2002, Volume 1, Issue 2, pp 181-224 Science as if situation mattered Michel Bitbol**. Caroline Williams investigated affective density of the political and its effect on our understanding of political subjectivity. Taking up Spinoza's challenge to think about affect beyond corporeal embodiment, he argues that there is a modality of affectivity that cannot simply be inscribed within the borders of subjectivity. theorising affect as an impersonal force anchored in a relational ontology that gives due recognition to the circulation of affects, as well as to their ambivalent structure in creating sites of identification, and he utilises this ontology to reflect on the dynamic of the political and the shape of political subjectivity. Williams argues that Spinoza's philosophy (through ideas of conatus and imagination) offers the conceptual resources to reconfigure the composition of affective subjectivity as a transindividual social bond and as an unconscious dynamic of ethico-political existence. **Subjectivity (2010) 3, 245–262. doi:10.1057/sub.2010.15 Affective processes without a subject: Rethinking the relation between subjectivity and affect with Spinoza Caroline Williams** Phenomenological Time bears ample evidence and infallible observatory and apodictic evidence albeit with a sense of chagrined circumvention and defeated disgruntlement circularity of both psychological and phenomenological interpretations of art which are sometimes insightful, but most often work on the logic of cyclical time and unavoidable drives and representations played out as a ‘natural’ order: where ‘men can’t avoid being boys’ a circularity which valorizes a helplessness/truthfulness conundrum. One begins with a classical phenomenological fallacy with Duchamp’s Fountain, does the urinal keep returning or do we keep returning to the urinal? Put in this way, it is a male question, something that phenomenology conveniently conceals when positing the “I”, or indeed, the “we”. But we must be careful not to simply assume that the phenomenological first person is white, heterosexual, able-bodied and male, providing a foundation upon which knowledge should be based as a universal condition. Deleuze’s second critique of time is linked to Kant. This model of time extends not only to phenomenology but to the phenomenological interpretation of art and is seen as a straight line. In the Critique of Pure Reason, **Kant cuts up circular time and reassembles it as a series of sensory experiences which memory processes and which constitute subjectivity**. Whereas the circular model of time (and interpretation) inscribes fatalism into the subject and the world, the phenomenological method is in danger of solipsism, where the subject constitutes the world. Much of embodied philosophy and enactivist theories of cognition go down this line with a plethora of art historical analyses by the likes of Rosalind Krauss, Michael Fried and others taking the lead from Merleau-Ponty. These approaches successfully challenge the **homunculus model of a Cartesian mind-body** dualism where there is a supervisor in the mind who processes experience and even consciousness and to whom all representations are directed. But phenomenology is mainly concerned with **perceptual consciousness**, the processing of colors, shapes, forms, motion in art and the world. The senses and sensorimotor processes are the fundamental pathways by which the interior is connected to the exterior in a chiasm of self-constitution in the world. The problem with this is that we explore the art through our sense perceptions, and what does art reveal to us?—our sense perceptions! Such solipsistic approaches have real problems relating complex and abstract conceptual production to bodily and sensorimotor contingencies, and they nearly always posit the subject as the site and object of discovery in the world. To be fair, phenomenology has moved on to rather more sophisticated models of

intersubjectivity, taking into account the specificities of different kinds of bodies and ways of being beyond the white, male heterosexual as the common denominator, yet many naïve approaches to art perpetuate the notion that perceptual, rather than conceptual processes and evolutionary neural patterns are primary, fundamental ways to understand the art experience. In fact, these aspects are only part of art production and reception, aspects which are often filtered out or made known to us in order for us to parody such trigger responses, as in much of Duchamp's work, which holds in contempt such retinal approaches to art. If we take the **subject of abstract expressionism** in the 1950s or later minimalist art, art history, teaching and interpretation still insists on putting the act of perceiving as fundamental to the constitution not only of the phenomenological first person but also of art itself. The male act of usually standing up, which consists of directing a stream with the hand(s) into a container or on a flat surface, sand or snow in order to leave patterns and traces (the usual child's prank) underlies the logic of macho 1950s Abstract Expressionist culture, especially that of Jackson Pollock, so effectively parodied by Carolee Schneeman's *Vagina Scroll*, 1975. Here, painting or writing is a kind of a discharge which invests the **object and space with** the phenomenological "I", disciplined by existential struggle and reflected back by the traces, signatures and gestures left behind in the work of art, rediscoverable as an element in the making of other subjectivities. It is precisely this kind of solipsism which is blind to the art event as something well in excess of re-establishing habits of subjective self-identity through the production and interpretation of art. The attempt to reinforce a phenomenological identity upon the event by personalizing sense experience is unable to see art as something which remodels the phenomenological "I" and neural plasticity itself. We learn new technologies and redraw our neural pathways through such new and spontaneous experiences which draw us into intersubjective worlds. Duchamp's *Fountain* and Deleuze's *Repetition and Difference* Grégory Minis sale (*I have made some changes*) The relation of different groups to deconstruction is settled by its relation to groups. An American enthusiast--one who has had a role in the development of our affection for Derrida--notices that the motivational chains are phobically driven by concerns with social scale. Richard Rorty: "He wants to figure out how to break with the temptation to identify himself with something big. . . ." (10) And: "So I take Derrida's importance to lie in his having had the courage to give up the attempt to unite the private and the public, to stop trying to bring together a quest for private autonomy and an attempt at public resonance and utility. He privatizes the sublime, having learned from the fate of his predecessors that the public can never be more than beautiful." (11) Notice is made here of what is indeed the telling feature. Levinas is praised, for example, for his alarm when "the social will [is] sought in an ideal of fusion . . . the subject losing himself in a collective representation, in a common ideal. . . . It is the collectivity which says 'us,' and which, turned toward the intelligible sun, toward the truth, experience, the other at his side and not face to face with him. The tellingly titled *D'un ton apocalyptique* contains passages that are particularly excited: These people situate themselves outside the ordinary, but they have in common this: they describe themselves as having an immediate and intuitive relation with mystery. And they want to attract, to seduce, and lead others to the mystery, through mystery. This agogic function of the leader of men, of the duce, of the Führer, of the leader, places him above the crowd that he manipulates with the aid of a small numbers of adepts joined together in a sect with a secret language, a clique or a small party with its ritualized practices. The mystifiers pretend to have exclusive access to the privilege of a secret mystery. . . . The revelation or the unveiling of the secret is something that they jealously reserve for themselves. Jealousy is here a major characteristic. There is not narcissism and non-narcissism; there are narcissisms that are more or less comprehensive, generous, open, extended. What is called non-narcissism is in general but the economy of a much more welcoming, hospitable narcissism, one that is much more open to the experience of the other as other. I believe that without a movement of narcissistic reappropriation, the relation to the other would be absolutely destroyed; it would be destroyed in advance. The relation to the other--even if it remains asymmetrical, open, without reappropriation--must trace a movement of reappropriation in the image of oneself for love to be possible, for example. Love is narcissistic. Beyond that, there are little narcissisms, there are big narcissisms. . . ." (23)

Justice of the Pieces Deconstruction as a Social Psychology Douglas Collins Sabine Wilke cogently argues for the rise of the aesthetic dimension in philosophical writing by showing how the metaphorical and poetic style of Heidegger, the principles of phrase configurations with Adorno, or the topographic writing of Derrida replace through their very form something what traditionally would have been subject to discursive exposure. The concern shifts from telling to showing, showing first and foremost that something remains untold. An exemplary book in this

aesthetic drive is Avital Ronell's Telephone Book where the playful layout and explicit break with traditional design interrupt the sequential logic of the paragraphs thereby performing the very argument at stake: open up new circuits of meaning and signification. The engagement with form and publishing formats is only consequential since media theory as exposed by Walter J. Ong, Eric A. Havelock, Parry Miman or Marshall McLuhan besides many others made it clear **that different communication technologies condition the mode of thought of a given culture**. The fractals of Benoit Mandelbrot or the butterfly shaped Lorenz attractor are the popular icons of this turn in the sciences. For communication studies and philosophy, Vilém Flusser was one of the pioneer thinkers reflecting the rise of what he called "techno-images." The photograph is the paradigmatic case which condenses the whole crises of Western thought and culture on its grainy surface. It subverts the linear logic of writing.⁶ Flusser's central diagnosis detects a shift from the grand narratives associated with the literate mind and Modernity towards creating and envisioning the world through image based practices. Writing himself in a succinct, visually inspired style, Flusser contents that the web of meaning, spun throughout the centuries by our textual practices, has become transparent. We discover the void between the letters through which the world escapes: "We are alienated from the world of text, precisely because we see through them as our own product."⁷ As a consequence we are confronted with a world of dissociated facts and information bits. Flusser hails this fragmented world as the advent of the zero dimensional point-universe that will provide the new raw material for our image based practices. Techno-images are nothing else than a conscious, inherently artistic drawing-together of point-elements as exemplified first and foremost by the silver-grains of the photograph but also by more recent pixel based techno-images. Similar to blueprints, designs or statistical curves, the weather map for example visualizes and contracts a tremendous amount of meteorological data from widely distributed stations into a single image of high- and low pressure

areas, temperatures, wind speeds, etc. Techno-images "compute" in this sense the universe of unimaginable, abstract information-bits into a concrete surface – the image. Echoing Paul Feyerabend, these image practices have to be understood as creative practices where the truth of scientific models has been replaced by its aesthetic quality

Aesthetics, Writing, Networked Computers Jörg Muller Dissertation Submitted to the Division of Media and Communications of The European Graduate School in Candidacy for the Degree of Doctor of Philosophy

Lothar Schäfer (1) and Sisir Roy (2) in Quantum Reality, the Importance of Consciousness in the Universe, the Discovery of a Non-Empirical Realm of Physical Reality, and the Convergence with Ancient Traditions of Indian and Western Philosophy investigate metaphysical aspects that include 1) the discovery of a non-empirical part of physical reality in a realm of potentiality; 2) the **emanation of the empirical world out of a realm of non-material forms**; 3) the discovery that the nature of physical reality is that of an indivisible Wholeness – the One; and 4) panpsychism: the possibility that the One is aware of its processes like a Cosmic Consciousness. The convergence of powerful traditions of seemingly disparate cultures is particularly important to point out in the present process of globalization, when a unifying view is needed to avoid controversy and conflict. To quote them in extenso: This is the nonlocality of the quantum world. Menas Kafatos and Robert Nadeau (1990) have drawn a remarkable conclusion from this phenomenon: If reality is nonlocal, the nature of the universe is that of an undivided wholeness. Because our consciousness has emerged from this wholeness and is part of it, it is possible to conclude that an element of consciousness is active in the universe: a Cosmic Consciousness. Pre-reflective self-consciousness is pre-reflective in the sense that (1) it is an awareness we have before we do any reflecting on our experience; (2) it is an implicit and first-order awareness rather than an explicit or higher-order form of self-consciousness. Indeed, an explicit reflective self-consciousness is possible only because there is a pre-reflective self-awareness that is an ongoing and more primary self-consciousness. Although phenomenologists do not always agree on important questions about method, focus, or even whether there is an ego or self, they are in close to unanimous agreement about the idea that the experiential dimension always involves such an implicit pre-reflective self-awareness. In line with Edmund Husserl (1959, 189, 412), who maintains that consciousness always involves a self-appearance (Für-sich-selbst-erscheinens), and in agreement with Michel Henry (1963, 1965), who notes that experience is always self-manifesting, and with Maurice Merleau-Ponty who states that consciousness is always given to itself and that the word 'consciousness' has no meaning independently of this self-givenness (Merleau-Ponty 1945, 488), Jean-Paul Sartre writes that pre-reflective self-consciousness is not simply a quality added to the experience, an accessory; rather, it constitutes the very mode of being of the experience: This self-consciousness we ought to

consider not as a new consciousness, but as the only mode of existence which is possible for a consciousness of something (Sartre 1943, 20 [1956, liv]). The notion of pre-reflective self-awareness is related to the idea that experiences have a subjective 'feel' to them, a certain (phenomenal) quality of 'what it is like' or what it 'feels' like to have them. As it is usually expressed outside of phenomenological texts, to undergo a conscious experience necessarily means that there is something it is like for the subject to have that experience (Nagel 1974; Searle 1992). This is obviously true of bodily sensations like pain. But it is also the case for perceptual experiences, experiences of desiring, feeling, and thinking. There is something it is like to taste chocolate, and this is different from what it is like to remember what it is like to taste chocolate, or to smell vanilla, to run, to stand still, to feel envious, nervous, depressed or happy, or to entertain an abstract belief. Yet, at the same time, as I live through these differences, there is something experiential that is, in some sense, the same, namely, their distinct first-personal character. All the experiences are characterized by a quality of mineness or for-me-ness, the fact that it is I who am having these experiences. All the experiences are given (at least tacitly) as my experiences, as experiences I am undergoing or living through. All of this suggests that first-person experience presents me with an immediate and non-observational access to myself, and that consequently (phenomenal) consciousness consequently entails a (minimal) form of self-consciousness. To put it differently, unless a mental process is pre-reflectively self-conscious there will be nothing it is like to undergo the process, and it therefore cannot be a phenomenally conscious process. The mineness in question is not a quality like being scarlet, sour or soft. It doesn't refer to a specific experiential content, to a specific what; nor does it refer to the diachronic or synchronic sum of such content, or to some other relation that might obtain between the contents in question. Rather, it refers to the distinct givenness or the how it feels of experience. It refers to the first-personal presence or character of experience. It refers to the fact that the experiences I am living through are given differently (but not necessarily better) to me than to anybody else. It could consequently be claimed that anybody who denies the for-me-ness of experience simply fails to recognize an essential constitutive aspect of experience. Such a denial would be tantamount to a denial of the first-person perspective. It would entail the view that my own mind is either not given to me at all — I would be mind- or self-blind — or is presented to me in exactly the same way as the minds of others. There are also lines of argumentation in contemporary analytical philosophy of mind that is close to and consistent with the phenomenological conception of pre-reflective self-awareness. Alvin Goldman provides an example: [Consider] the case of thinking about x or attending to x. In the process of thinking about x there is already an implicit awareness that one is thinking about x. There is no need for reflection here, for taking a step back from thinking about x in order to examine it... When we are thinking about x, the mind is focused on x, not on our thinking of x. Nevertheless, the process of thinking about x carries with it a non-reflective self-awareness (Goldman 1970, 96). A similar view has been defended by Owen Flanagan, who not only argues that consciousness involves self-consciousness in the weak sense that there is something it is like for the subject to have the experience, but also speaks of the low-level self-consciousness involved in experiencing my experiences as mine (Flanagan 1992, 194). As Flanagan quite correctly points out, this primary type of self-consciousness should not be confused with the much stronger notion of self-consciousness that is in play when we are thinking about our own narrative self. The latter form of reflective self-consciousness presupposes both conceptual knowledge and narrative competence. It requires maturation and socialization, and the ability to access and issue reports about the states, traits, dispositions that make one the person one is. To claim that every kind of self-consciousness is conceptual is overly cognitive. Bermúdez (1998), to mention one further philosopher in the analytic tradition, argues that there are a variety of **nonconceptual forms of self-consciousness** that are "logically and ontogenetically more primitive than the higher forms of self-consciousness that are usually the focus of philosophical debate" (1998, 274; also see Poellner 2003). This growing consensus across philosophical studies supports the phenomenological view of pre-reflective self-consciousness. (References: Stanford Encyclopedia, Post Modernist Theories, Internet Encyclopedia of Philosophy) Korzybski pointed out that there are two ways to slice easily through life; to believe everything or to doubt everything. Both ways save us from thinking. So, from time to time you have to give yourself a Virtual Reality Check. The hallmark of postmodern philosophy has been disbelief or skepticism of all "metanarratives," or translations of reality. Postmodernism has even turned its profound skepticism on such important humanist concepts as "objective truth" and reason. Yet, for a deconstructionist postmodern society, individually we are still riddled with superstition and gullibility, and open to

manipulation through our belief systems as any politician, philosopher, clergy, or salesperson will attest. Further, most people are painfully naive when it comes to even the simplest scientific understanding. Most of us don't have a clue about the fundamental nature of physical reality or our own psychological nature, and our ability to be fooled by our senses and mind, or methods of social and technological persuasion. We may think we know, but the penetration is deeper than we dare imagine. Nature rejects the naïveté that seeks absolute truth. We are beginning to realize, individually and culturally, that "realities" are all human constructions. The task becomes one of "catching ourselves in the act" of creating our own "reality" from the flow of events. Human truth is always an engagement of mind with experience.. We don't need to fear the collapse of our personalistic belief system (the "box" we live in), nor our belief in absolute truth. **A strong desire to engage in the "quest for uncertainty" complements our anxiety that perhaps there is no absolute, objective ground to reality.** The warrant of Truth is ever elusive when we deconstruct the foundational justifications of our convenient notions about the way the world works. It is easy to confuse what is actually the creation of beliefs with the "discovery of Truth," a common goal of science and theology. Science offers no ontological Scientific Picture of the World but substitutes a number of applied theories as meta-theory. It's close, powerful and useful: it 'works' and allows use to do work. But the still mysterious "true" metatheory may reveal the basis of consciousness. But who knows when we will lift that veil? Permeating the living reality of our culture are certain contagious notions, fads and trends that have the ability to influence the way we think about the nature of Reality and ourselves. Some of them are toxic and can consume you; others are just whims of pop culture infectious and benign as the common cold. **Jung** described the concept of **psychic contagion** by certain archetypal forces inherent in the human psyche, which manifest in our spiritual lives and belief systems. He spoke of both conscious and unconscious contamination. Notions like this range from simple superstitions to scientific concepts, to urban myths. Notions sweep through our culture and **insinuate themselves** within its fabric, as fads, whether they are "real" or not, they can be influential. They mood alter us, make us feel we belong. An analysis of these notions is useful in distinguishing a common human phenomenon from any potential "alien influence" which may or may not be exerted on us from an unknown source. **Strange Attractors: Transference, Holography, and an Archetype** **Burke, J. (2003). Strange Attractors: Transference, Holography, and an Archetype (Doctoral dissertation, Pacifica Graduate Institute, 2003).** In consideration to the Nobel Prize for Chemistry this year(2014), a stability analysis of the problem in single molecules would be a contemporaneous question that has to be taken cognizance of and given importance to. Determination of molecular structure by geometry optimization became routine only after efficient methods for calculating the first derivatives of the energy with respect to all atomic coordinates became available. Evaluation of the related second derivatives allows the prediction of vibrational frequencies if harmonic motion is estimated. More importantly, it allows for the characterization of stationary points. The frequencies are related to the eigenvalues of the Hessian matrix, which contains second derivatives. If the eigenvalues are all positive, then the frequencies are all real and the stationary point is a local minimum. If one eigenvalue is negative (i.e., an imaginary frequency), then the stationary point is a transition structure. If more than one eigenvalue is negative, then the stationary point is a more complex one, and is usually of little interest. When one of these is found, it is necessary to move the search away from it if the experimenter is looking solely for local minima and transition structures. The total energy is determined by approximate solutions of the time-dependent Schrödinger equation, usually with no relativistic terms included, and by making use of the Born–Oppenheimer approximation, which allows for the separation of electronic and nuclear motions, thereby simplifying the Schrödinger equation. This leads to the evaluation of the total energy as a sum of the electronic energy at fixed nuclei positions and the repulsion energy of the nuclei. Notable exceptions are certain approaches called direct quantum chemistry, which treat electrons and nuclei on a common footing. Density functional methods and semi-empirical methods are variants on the major theme. **Orlando Alvarez** discusses Polyakov's quantization of the string in the presence of a boundary allowing for an arbitrary topology for the world sheet. In addition to the dynamical conformal factor discovered by Polyakov, there are a finite number of new degrees of freedom if the surface is more complicated than a sphere or a disc. The quantization of the Liouville theory in an arbitrary topology is also discussed. A one-loop calculation shows that the model is renormalizable if one performs a mass renormalization and an additive field renormalization. The renormalization group equations have a perturbative infrared unstable fixed point in all topologies. Copyright © 1983 Published by Elsevier B.V.

Nuclear Physics B Volume 216, Issue 1, 25 April 1983, Pages 125–184 Theory of strings with boundaries: Fluctuations, topology and quantum geometry Orlando Alvarez DOI: 10.1016/0550-3213(83)90490-X

Quantum geometry (the modern loop quantum gravity involving graphs and spin-networks instead of the loops) provides microscopic degrees of freedom that account for black-hole entropy. However, the procedure for state counting used in the literature contains an error and the number of the relevant horizon states is underestimated. In our paper a correct method of counting is presented by **Marcin Domagala and Jerzy Lewandowski**. Results lead to a revision of the literature of the subject. It turns out that the contribution of spins greater than $1/2$ to the entropy is not negligible. Hence, the value of the Barbero–Immirzi parameter involved in the spectra of all the geometric and physical operators in this theory is different than previously derived. Also, the conjectured relation between quantum geometry and the black-hole quasi-normal modes should be understood again. **Marcin Domagala and Jerzy Lewandowski 2004 Class. Quantum Grav 21 5233 doi:10.1088/0264-9381/21/22/014**

Black-hole entropy from quantum geometry Quantum geometry predicts that a universe evolves through an inflationary phase at small volume before exiting gracefully into a standard Friedmann phase. This does not require the introduction of additional matter fields with ad hoc potentials; rather, it occurs because of a quantum gravity modification of the kinetic part of ordinary matter Hamiltonians. Authors draw the cognizance of application of the same mechanism that can explain why the present day cosmological acceleration is so tiny. DOI: <http://dx.doi.org/10.1103/PhysRevLett.89.261301>

Inflation from Quantum Geometry Phys. Rev. Lett 89, 261301 – Published 12 December 2002 Martin Bojowald The loop quantum cosmology of the closed isotropic model is studied with special emphasis on a comparison with traditional results obtained in the Wheeler–DeWitt approach by **Martin Bojowald**. Investigation includes the relation of the dynamical initial conditions to boundary conditions such as the no-boundary or the tunneling proposal and a discussion of inflation from quantum cosmology. DOI: <http://dx.doi.org/10.1103/PhysRevD.67.124023>

Loop quantum cosmology, boundary proposals, and inflation Phys. Rev. D 67, 124023 – Published 23 June 2003 Martin Bojowald and Kevin Vandersloot In loop quantum cosmology, the universe avoids a big bang singularity and undergoes an early and short super-inflation phase. During super-inflation, non-perturbative quantum corrections to the dynamics drive an inflaton field up its potential hill, thus setting the initial conditions for standard inflation. **Shinji Tsujikawa et al** show that this effect can raise the inflaton high enough to achieve sufficient e-foldings in the standard inflation era. They also analyse the cosmological perturbations generated when slow-roll is **violated after super-inflation** and show that loop quantum effects can in principle leave an **indirect signature** on the largest scales in the CMB, with some loss of power and running of the spectral index. **Shinji Tsujikawa et al 2004 Class. Quantum Grav 21 5767 doi:10.1088/0264-9381/21/24/006**

Loop quantum gravity effects on inflation and the CMB Quantum geometry from quantum information is the study of **Florian Girelli and Etera R Livine**. Loop quantum gravity defines the quantum states of space geometry as spin networks and describes their evolution in time. We reformulate spin networks in terms of harmonic oscillators and show how the holographic degrees of freedom of the theory are described as matrix models. This allows us to make a link with non-commutative geometry and to look at the issue of the semi-classical limit of loop quantum gravity from a new perspective. This work is thought of as part of a bigger project of describing quantum geometry in quantum information terms. **Florian Girelli and Etera R Livine 2005 Class. Quantum Grav 22 3295 doi:10.1088/0264-9381/22/16/011**

Reconstructing quantum geometry from quantum information: spin networks as harmonic oscillators. Quantum geometric mechanics is the topic of study by **Dorje C. Brody Lane P. Hughston** establishing a clear relation between the quantum mechanics and quantum geometry and also the entangled states. The manifold of pure quantum states can be regarded as a complex projective space endowed with the unitary-invariant Fubini–Study metric. According to the principles of geometric quantum mechanics, the physical characteristics of a given quantum system can be represented by geometrical features that are preferentially identified in this complex manifold. Here we construct a number of examples of such features as they arise in the state spaces for spin, spin 1, spin and spin 2 systems, and for pairs of spin systems. A study is then undertaken on the geometry of entangled states. A locally invariant measure is assigned to the degree of entanglement of a given state for a general multi-particle system, and the properties of this measure are analysed for the entangled states of a pair of spin particles. With the specification of a quantum Hamiltonian, the resulting Schrödinger trajectories induce an isometry of the Fubini–Study manifold, and hence also an isometry of each of the

energy surfaces generated by level values of the expectation of the Hamiltonian. For a generic quantum evolution, the corresponding Killing trajectory is quasiergodic on a toroidal subspace of the energy surface through the initial state. When a dynamical trajectory is lifted orthogonally to Hilbert space, it induces a geometric phase shift on the wave function. The **uncertainty** of an observable in a given state is the length of the gradient vector of the level surface of the expectation of the observable in that state; a fact that allows us to calculate higher orders corrections to the Heisenberg relations. A general mixed state is determined by a probability density function on the state space, for which the associated first moment is the density matrix. The advantage of a general state is in its applicability in various attempts to go beyond the standard quantum theory, some of which admit a natural phase-space characterization. **Journal of Geometry and Physics Volume 38, Issue 1, April 2001, Pages 19–53 Geometric quantum mechanics Dorje C. Brody Lane P. Hughston DOI: 10.1016/S0393-0440(00)00052-8** Expansion (unmesa) of [consciousness] or the creative intuition {pratibha} is (=) [experienced] in the interval which divides two [moments] of differentiated perception (vikalpa) It is here that they arise and disappear. The Sastras and Agamas proclaim with reasoned argument that it is (=) free of thought-constructs {Nirvikalpa} and precedes (e) Path to Liberation all mental representations of any object. None can deny that a gap exists between perceptions insofar as two moments of thought are (=) invariably divided. This [gap] is (=) the undifferentiated unity of all the countless manifestations. Similarly, in the outer more objective sphere, where change consists (e) of the alterations in the configurations of manifest appearances (Abhasa\ the transition from one to (e&eb) another corresponds to (e&eb) a phase of pure luminosity that marks (eb) the beginning of one form and the end of another. The world of manifestation and differentiated perceptions (Yikalpa) thus extends from one Centre to (eb) the next. **What is meant again is the realisation or the evolution of individual consciousness or general ledger to be the cosmic general ledger is the sole criterion for the Satchidananda state where all dualities cease** Although it is never in fact divorced from (e) the subject who resides there, the ignorant fail (e) to grasp this fact and so, cut off from (e) the Centre, the world of objectivity becomes (=) for them the sphere of Maya. Bhagavatopala quotes the Light of Consciousness amvitprakdsa): This ever pure experience (suddhanubhava) is variegated by (e) each form [revealed within it] Even so it remains unstained (nirmala) when moving to another. Just as a cloth which is naturally white, once dyed, cannot (e) change colour without [first] becoming white again, similarly the pure power of awareness, (citi) once coloured by (e) form, is pure [again] at the Centre where that form is (=) abandoned and from whence it proceeds to (e&eb) another. In his Essence of Vibration (Spandasarpdoha), K\$Emaraja explains that the rise and fall of every individual perception in the field of awareness is (=) a specific pulsation of consciousness. **This is exactly the individual general ledger transaction lifted to the transaction in cosmic general ledger.** During The Initial Instant Of Perception, T Consciousness Is Manifestly Apparent And The Yogi, Participating In Its Plenitude, Observes The Outer World Without Being Attached To Any Particular Divine Body And Sacred Circle Of The Senses Singling It Out From Any Other, Like A Man Who Observes A City From A High Mountain Peak. He Sees The Outer World Reflected Within His Consciousness Free Of Thought-Constructs And So 'Stamps* The Outer On The Inner While Absorbing The Object And Means Of Knowledge In The Pure Subject Which Grasps Them As The Expansion Of His Own Nature. K\$Emaraja Says: By Penetrating Into Bhairavimudra, The Yogi Observes The Vast Totality Of Beings Rising From, And Dissolving Into, The Sky [Of Conscious- Ness], Like A Series Of Reflections Appearing And Disappearing Inside A Mirror. Through The Practice Of Bhairavimudra The Yogi Realises That He Is The Substratum Consciousness (Adhisfhatr) Which Both Underlies And Is The Essence Of All Things. He Discovers That Phenomena Have No Independent Existence Apart From Him And So Are, In This Sense, Void. At The Same Time, He Realises That Because All Things Are Consciousness, They Are Far From Unreal. He Views The Outer World Yet Sees It Not. Beyond Both Voidness And Non-Voidness He Penetrates Into The Supreme Abode (Param Padam) Of Siva's Consciousness. [The Powers Of The Senses] Endowed With The Attributes Of The Great Union [Between Subject And Object] Whose Form Is The Wakening Of Man's Spiritual Potential (Kuntfalinf), Fill [With Consciousness] The Outer Clatter Of Diversity (Bhedatfambara) Born Of Its Intense Power And Are Then Established In The Unobscured Abode Of The Void Of Consciousness To Shine [There] Eternally. Thus Residing Beyond Being And Non-Being, The Sole Protector Of The Unity Which Is Tranquil And Expanding [Consciousness], Whose Glory Is All-Embracing And Form Unobscured, Is Called Bhairavimudra. Through The Practice Of Bhairavimudra, The Yogi Unites The Universal Vibration Of T Consciousness With The Individual

Pulsation Of Objectivised 'This' Consciousness. The Two Aspects Of Consciousness Are Now In A State Of Equilibrium Like The Two Pans Of An Evenly Weighed Balance And The Yogi Experiences The Pure Knowledge (Suddhavidya) That: 'I Alone Am All Things'. Thus Becoming The Master Of The Wheel Of Energies He Is Free, Like Siva, To Create And Destroy. When [The Yogi] Is Well Established, Without Wavering, Solely In The Integral Egoity Of His Authentic Nature, The Spanda Principle, And Is Absorbed In Contemplation (Samavisfa), He Becomes One With It The Doctrine Of Vibration (Tanmaya). Then . . . Dissolving And Creating The Universe By Means Of His Introverted And Extroverted Absorption, He Destroys And Creates All Things Out Of Sarikara, His Innate Nature. [Thus] He Assumes The State Of The Universal Experienter And Having Absorbed All That Is To Be Experienced From [The Grossest Level] — Earth — To (The Subtlest) — Siva — He Reaches The State Of The Supreme Subject By Progressively Recognising [His Identity With Him]. Thus, Introverted And Extroverted Absorption Both Lead To The Recognition Of The Pulsation (Spanda) Of One's Own Consciousness. This Is Where Individual General Ledger Becomes Cosmic General Ledger. At The Level Of Consciousness Corresponding To Siva's Basic State (Saryibhavadvasta), The Alternation From Inner To Outer Is Instantaneously Resolved Into The Vibration Of His Nature. When The Yogi Finally Comes To Be Constantly Aware Of This Reality, His Enlightenment Is Full And Perfect. Freed Of All Means (Anupaya) And Delighting In The Power Of His Bliss (Anandasakti), He Knows And Does Whatever He Pleases. The Yogi Seeking Self-Realisation Must Acquire Mastery Over This Movement. Ksemaraja Stresses That The Doctrine Of Vibration Teaches That Liberation Can Only Be Achieved By First Withdrawing All Sense Activity In Introverted Contemplation (Nimilanasamadhi) To Then Experience The 'Great Expansion' (Mahdviksd) Of Consciousness While Recognising This To Be A Spontaneous Process Within It. This Is Done Through The Practice Of Kramamudra. A Passage From The Now Lost Kramasiitra Explains: Although The Adept's Attention [May Be] Outwardly Directed, He Enjoys Contemplative Absorption Through The Introverted Aspect Of Kramamudra. Initially He Turns Inward From The Outside World And [Then] From Within [Himself] He Exits Into The Outer World Under The Influence Of His Absorption. There Is No Necessity Of Any Glorification Or Mortification Here. It Is Just Like Joining A Job And Retiring And Realising All That Happened Is Just By His Own Effort And Nothing Else. Thus The Sequence (Krama) In This Attitude (Mudra) [Ranges Through] Both Inner And Outer. The Yogi Must Pervade The Surface Level Of Awareness (Yyutthana) With The Same Bliss He Experiences Plunging Into The Depths Of Contemplative Absorption (Samadhi). Note Here That Samadhi Is A Death Like State And Still Being Matter And Information Stored You Experience Consciousness. Submerging Himself And Emerging Repeatedly From Samadhi, He Eventually Recognises That The Unity Of Consciousness Pervades Both States: The Best Of Yogi's, Who Has Achieved A State Of Complete Absorption Even When Risen From Meditation, [Inwardly] Vibrating Like A Drunkard In Blissful Inebriation From The After-Effects Of The Nectar Of Contemplation, Sees All Things Dissolving In The Sky Of Consciousness Like A Cloud In The Autumn Sky. He Plunges Repeatedly Within Himself Divine Body And Sacred Circle Of The Senses And Becomes Aware Of His Identity With Consciousness By The Practice Of Introverted Contemplation. Thus Even When He Is Said To Have Risen From Absorption, He Is One With [His] Experience Of It. Ksemaraja Goes On To Explain That This Practice Is Called 'Mudra' Because It Both Fills The Adept With Bliss (Mud) And Is Itself The Bliss Of Consciousness. Moreover, It Dissolves Away (Dra) All Bondage And 'Stamps' The Universe Of Experience With The Seal (Mudra) Of The Fourth State (Turiya) Of Enlightened Consciousness Beyond, And Including, The Three States Of Waking, Dreaming And Deep Sleep. It Is Called 'Krama' Because It Is The Root Source Of All Emanation And All Other Conscious Processes Which Succeed One Another In Ordered Sequence (Krama) And Is, At The Same Time, Their Successive (Krama) Appearance As Well. By The Practice Of Kramamudra The Opposites Fuse And Siva And Sakti Unite. They Yogi Comes To Experience The Simultaneous Pervasion Of All The Lower, Grosser Categories Of Existence By The Higher And The Presence Of The Lower In The Higher. Commencing His Practice In A Low Form Of Bhairavimudra, The Yogi Conjoins The Outer With The Inner; Then, In Kramamudra, He Fills Both The Outer With The Inner And The Inner With The Outer. When He Achieves Perfection In This Two-Fold Movement, He Attains To The Highest Form Of Bhairavimudra In Which The Two Merge Completely In The Experience Of The Absolute (Anuttara), Free Of All Differentiation And Polarities. If He Fails To Maintain Awareness Of This State, He Again Falls Into Kramamudra Until He Has Finally Completely Merged All The Highest States In The Lower

And The Lower In The Higher. He Then No Longer Needs To Resort To Any Means (Anupaya) To Achieve Liberation. All He Says Or Does Anything He Perceives Or Thinks, Instantly Occasions In Him The Highest Level Of Consciousness. Thus The Fruit Of Bhairavimudra Is The Wonder (Camatkara) Or Amazement (Vismaya) That Overcomes The Yogi When He Reaches The Plane Of Union (Yogabhumika) Y Xsl Where All Opposites Merge In The Radiance Of The Great Light Of Consciousness. The Stanzas On Vibration Teach: How Can One Who, As If Astonished, Beholds His Own Nature As That Which Sustains [The Existence Of Everything] Be Subject To This Painful Round Of Transmigration (Kusrti)? The Yogi, Recognising His True Nature To Be The Supreme Subject, Is Astonished To Suddenly Discover That The Individual He Thought He Was, Caught Up In The Trammels Of Thought And Living In A World Enmeshed In The Web Of Time And Space, Does Not Really Exist At All. He Experiences A 'Turning About' (Paravrtti) In The Deepest Seat Of The Doctrine Of Vibration Consciousness As He Penetrates His True Nature. The Sudden Eruption Of This Intuition (Pratibha) Arouses In Him A Cry Of Amazement As He Transcends All Thought-Constructs And, Perfectly Absorbed In His Own Nature, Is Liberated. The Path To Liberation Essentially, Spanda Doctrine Is Concerned With Two Matters. The First Is To Impart To Those Who Are Fit To Receive The Teachings A Deeper Understanding Of The Ultimate Goal Of Life (Upeya). When We Have Understood What Truly Benefits Us And Is Worth Attaining And What, On The Contrary, Is Of No Real Value But Stands In The Way Of This Attainment, We Can Begin To Make Progress Towards Our Goal. This Is Spanda Doctrine's Second Concern, Namely, To Show The Way In Which We Can Develop Spiritually Through Siva's Grace And The Right Application Of The Means To Realisation That It Teaches. When Both These Aspects Of The Teaching Have Been Correctly Understood And Applied, The Spanda Yogi Achieves A Clear And Permanent Realisation Of His Goal And Is Liberated, Thus Fulfilling The Ultimate Aim Of The Teaching. The Doctrine Of Vibration Is Not Meant For The Spiritually Dull. It Is Not For The Worldly Whose Consciousness, Clouded By Ignorance, Is As If Dreaming, Even During The Waking State Of Daily Life, The Dream Of Its Own Thought-Constructs. The Teachings Are Meant For Those Who Are Awake (Prabuddha), Those Who, Full Of Faith And Reverence, Are Always Alert And Intent On Discerning The True Nature Of Ultimate Reality. This Reality Is Understood In Three Basic Ways. The First Is Purely Transcendental. The Stanzas Choose This Aspect As The One Which Formally Defines It Most Specifically. Ultimate Reality Transcends All The Opposites, Including Subject And Object. This Does Not Mean, However, That It Is An Unconscious Void, A Mere Absence Of All Existence. In Fact, This Negative Characterization Of Reality (Which Includes Also A Denial Of All That Is Unconscious) Implies A Positive Immanence In Which The Opposites Are United In The Oneness Of Pure Consciousness That Is Equally Siva And Spanda, His Universal Activity. These Two Seemingly Contrasting Aspects Are Reconciled In The Third, Namely, Reality Understood As The Essential Nature Of All Things. Although Universal And Everywhere The Same, It Is Understood To Be The Essential And Specific Nature Of Each Existent As Its 'Own Nature' (Svabhava). In The Case Of The Individual Soul It Is Even More Specific, More Personal As His Own 'Own Nature' (Svasvabhava). Note Here That Svabhava Is Not Svasbhava. In Fact We Made A Statement That Svabhava Is The Very Characteristics Of Space Time Like, Crime Zones, Silent Zones, School Zones We Classify Space In To. Belonging To None Other Than Oneself It Is The Pure Subjectivity That Perceives Experiences, Enjoys, Reflects, Thinks And Senses As Well As Being The Conscious Agent Who Creates Every Possible Form Of Experience In All The States Of Consciousness. The Liberating Knowledge Of Reality Thus Corresponds To Our Regaining Possession Of Ourselves (Svatmagraha). We Must Lay Hold Of Ourselves And Abide In Our Authentic Nature. Reality Coincides With Our Own Most Fundamental State Of Being (Svasthiti), Free Of All Contrasts And Contradictions. Once We Have Overcome The Negative Forces That Arise From Our Ignorance And Prevent Us From Abiding In Ourselves, We Are Liberated. To Do This, We Must Penetrate Through The Pulsing Fluctuations Of Objectively Experienced States And Perceptions At The Surface Level Of Consciousness And Gain Insight Into The Timeless Rhythm Of Our Own Nature Manifest In The Universal Arising And Falling Away Of All Things. We Are Not Freed Of The Trammels Of Perpetual Change By Setting It Aside; On The Contrary, We Must Gain Insight Into The Recurrent Cycles Of Creation, Persistence And Destruction, Or Else Be Bound By Our Ignorance. This Spiritual Ignorance Consists Essentially Of Our Contracted State Of Consciousness And So Can Only Be Effectively Countered By Expanding It To Reveal Our Own Authentic Nature As This Expanded State Itself, Which Is The Universal Vibration (Samanyaspanda) Of Consciousness. Vibration Of Strings Which Gives Shape To

Objects Is Itself A Information, A Sine Qua Non Of Individual Consciousness. The Spanda- Yogi Treads The Path Of Consciousness Expansion. The Movement From The Contracted To The Expanded State Marks The Transition From Ignorance To Understanding, From The Dispersion And Incompleteness Of A Form Of Consciousness Entirely Centred On An Objectively Perceived And Discursively Represented Reality To A Direct, Intuitive Awareness Of The Unity And Integral Wholeness Of Our Own Absolute Spanda Nature. Along The Way To This Supreme Realisation Consciousness Develops, As Veil After Veil Is Lifted, Until It Becomes Full And Perfect In The Absolute Which Encompasses Within Itself All Possible Formats Of Experience. As Abhinava Says: [This Realisation] Is The Supreme Limit Of Plenitude And As Such There Can Be No Higher Attainment. Any [Other] Attainment [We Can] Conceive Issues From A State That Falls Short Of [This] Perfection. Once* [This] Uncreated Fullness Has Been Attained, Pray Tell, What Other Fruit Can There Be [Beyond It]? The Fettered Soul's Contracted State Of Consciousness Binds Him Because He Is Deprived Thereby Of The Subtle, Intuitive Insight Into The Underlying Unity Of Existence And His Attention Is Focused Instead On Its Gross, Outer Diversity Easily Apparent To Everybody, However Restricted His Consciousness May Be. However, Although The Fettered Soul In This State Is Ignorant Of This Unity, This Does Not Mean That His Knowledge Of Diversity Is False. Ignorance Entails A Form Of Knowledge Which, Although Quite Correct, Is Binding. We Are Not Absolutely Ignorant Of Reality For If We Were We Would Be Totally Unconscious. Spiritual Ignorance Is Always Linked With Some Degree Of Consciousness. Those Subject To The Round Of Birth And Death Are Not Inert Clods Of Earth. This Also Leads To Conclusion That Shakti Responsible Is Impure Because Of Maya Imbued In It. Thus, Although Ignorance Obscures Consciousness, It Is Wrong To Think Of It, As Dualist Saivites Do, In Terms Of A Defiling Impurity That Shrouds It Like A Cloth Covering Ajar. Spiritual Ignorance Can Be Nothing But Consciousness Itself, Albeit In A Limited State. Siva, Who Is Universal Consciousness, Is The Innate Nature Of Both Its Contracted And Expanded States, Both Of Which Are Forms Of Knowledge, Namely: 1) Supreme Knowledge (Parajñāna) Defined As The Revelation Of One's Own Innate Nature As The One Reality Which Is The Being Of All Things. 2) Inferior Knowledge (Aparajñāna) Which Jayaratha Explains Results From The Mental Activity (Vyapāra) Of The Individual Subject Whose Consciousness Is Contracted. It Consists Of The Mental Representations (Vikalpa) He Forms Of Himself And His Object, Of The Type 'I Know This'. The Lower Knowledge Obscures The Higher And Binds The Soul By Breaking Up His Direct, Pervasive Awareness Of His Own Pure Consciousness Nature, Free Of Mental Representation. The Stanzas On Vibration Teach: Operating In The Field Of The Subtle Elements, The Arising Of Mental Representation Marks The Disappearance Of The Flavour Of The Supreme Nectar Of Immortality; Due To This [Man] Forfeits His Freedom. As We Have Already Seen, Three Factors Are Necessary For Perception And Thought To Be Possible, Namely, The Perceiving Subject, The Means Of Knowledge And The Object Perceived. Rajanaka Rama, In His Commentary On The Stanza Cited Above, Explains At Length That These Three Factors Correspond To Three Major Divisions In The Lower Thirty-One Categories Of Existence, Namely: 1) The Object. This Consists Essentially Of The Five Primary Sensations Which Are The Subtle Elements {Tanmatras} Of Smell, Taste, Sight, Touch And Sound Along With The Five Gross Elements — Earth, Water, Fire, Air And Ether — Of Which These Sensations Are The Perceivable Qualities. 2) The Means Of Knowledge. This Consists Of The Senses And The Inner Mental Organ. 3) The Subject. At This Level, The Subject Is The Individual Soul (Purusa) Whose Consciousness Is Contracted By The Five Obscuring Coverings (Kāncukas) Of Limited Knowledge And Action, Attachment, Natural Law And Time Along With Maya, Their Source. All These Categories Belong To The Impure Creation (Asuddhasrīti), Which Is The Sphere Of Maya Where The Lower Order Of Knowledge Operates And Subject And Object Are Divided. Above Them Are Five More Categories Which Belong To The Pure Creation (Suddhasrīti) Where Subject And Object Are Still United. The Highest Of Those Categories Are Siva And Śakti. Combined They Represent The State Of Pure T Consciousness And Its Sentient Subjectivity (Upalabdhrta), Respectively. The Next Category Is Called Sadasiva. Here Faint Traces Of Objectivity Appear In The Pervasive, Undivided Consciousness Of Siva And Śakti. Consciousness, Now Full Of The Power Of Knowledge (Jñānasakti), Views The All In A State Of Withdrawal (Nimesa), Shining Within, And At One With Its Own Nature. T Consciousness Predominates Over 'This' Consciousness Which It Encompasses In The Awareness That: 'I Am This [Universe]' (Aham-Idam). Next Comes The Category 'Isvara' Corresponding To The Awareness: 'This (Universe) Is Me' (Idam-Aham). 'This' Consciousness Takes The Upperhand Over T Consciousness And Unfolds Externally Full

Of The Creative Power Of Action (Kriya&Akti). The All Now Becomes More Clearly Manifest As An Independent Reality. It Is Still Experienced As One With Consciousness But Is No Longer Fully Merged Within It. Finally, When Both Subjective And Objective Aspects Share An Equal Status In The Two-Fold Awareness That: 'I Am This (Universe) And This (Universe) Is Me (Ahamidam-Idamaham)\ Pure Knowledge (Fuddhavidya), The Last Of These Categories, Emerges. The Pure Categories Are The Experience Of The Impure Categories When They Are Recognised To Be One With Consciousness. They Are Experienced Within The Domain Of The Pure Universal Subject The Enlightened Yogi Realises Himself To Be. Mental Representations (Yikalpa) Emerge From This Pure Awareness And Subside Into It In Consonance With The Rhythm Of The Emanation And Withdrawal Of The Lower Categories. Impelled By The Universal Will, This Movement Is Spontaneous And Free. Free Of All Hopes And Fears The Enlightened Yogi Sees All Things As Part Of This Eternal Cosmic Game, Played In Harmony With The Blissful Rhythm Of His Own Sportive Nature At One With All Things. The Stanzas On Vibration Teach: Everything Arises [Out Of] The Individual Soul And He Is All Things. Being Aware Of Them, He Perceives His Identity [With Them]. Therefore There Is No State In The Thoughts Of Words Or [Their] Meanings That Is Not Siva. It Is The Enjoyer Alone Who Always And Everywhere Abides As The Object Of Enjoyment. Or, Constantly Attentive, And Perceiving The Entire Universe As Play, He Who Has This Awareness (Sawvitti) Is Undoubtedly Liberated In This Very Life. According To The Doctrine Of Vibration, Only Liberation In This Life (Jivanmukti) Is Authentic Liberation. Liberation After Death (Yideha- Mukti) In Some Form Of Disembodied State Free Of All Perceptions And Notions Of The World Of Diversity Is Not The Ultimate Goal. Ksemaraja Stresses That Liberation Is Only Possible By Realising One's Own Identity With The Whole Universe, However Difficult This May Be. Similarly, He Maintains That The Suspension Of All Mental And Sensory Activity, Which Takes Place In The Introverted Absorption Of Contemplation With The Eyes Closed (Nimilanasamadhi) That Leads To Identification With Transcendent Consciousness Is Complemented And Fulfilled By The Cosmic Vision Had Through The Expansion Of Consciousness That Takes Place In Contemplative Absorption With The Eyes Open (Unmilanasamadhi). Consequently, Ksemaraja Explains That The First Of The Three Sections, Into Which He Divides The Stanzas, Deals With The Former Mode Of Contemplation And The Second Section With The Latter. Significantly, The Last Stanza Of The Second Section Ends With The Declaration That 'This Is The Initiation That Bestows Siva's True Nature'. In Other Words, This Realisation, Attained Through The Expanding Consciousness Of Contemplation With The Eyes Open, Initiates The Yogi Into The Liberated State, Which Is Identification With Siva Whose Body Is The Universe. In Order To Attain This Expanded State Of Liberated Consciousness, The Yogi Must Find A Spiritual Guide Because The Master (Guru) Is The Means To Realisation. The Master Is For His Disciple Siva Himself For It Is He Who Through His Initiation, Teaching And Grace, Reveals The Secret Power Of Spiritual Discipline. Instructing In The Purport Of Scripture He Does More Than Simply Explain Its Meaning: He Transmits The Realisation It Can Bestow. The Master Is At One With Siva's Divine Power Through Which He Enlightens His Disciple. It Is This Power That Matters And Makes The Master A True Spiritual Guide, Just As It Was This Same Power That Led The Disciple To Him In His Quest For The Path That Leads To The Tranquility That Can Only Be Found 'In The Abode Beyond Mind'. The Master Is The Ferry That Transports The Disciple Over The Ocean Of Thought — If, That Is, The Disciple Is Ready. The Disciple Must Be 'Awake' (Prabuddha) 2% Attending Carefully To The Pulse Of Consciousness. This Alert State Of Wakefulness Is At Once The Keen Sensitivity Of Insight As Well As The Receptivity Of One Who Has No Other Goal To Pursue Except Enlightenment. The Highest, Most Perfect Relationship The Disciple Can Have With His Master Is Such As It Is With Siva Himself: One Of Identity. The Exchange That Takes Place Between Them Is An Internal Dialogue Within Universal Consciousness, Their Common Identity (Svabhava). Limiting Itself To A Point Source (Anu) And Obscured By The Thought-Constructs Born Of Doubt And Ignorance, Consciousness Assumes The Guise Of The Disciple Who Seeks To Attain The Expanded Fullness Of His Master's Consciousness. The Master, On The Other Hand, Embodies The Aspect Of Consciousness Which Responds To The Inquiring Consciousness Of His Disciple. Free Of The Notions Of 'Self And 'Other', When The Disciple Is Liberated By His Grace, It Is The Master Who In Reality Liberates Himself. Although Ksemaraja Assures Us That The Master Can By Himself Enlighten His Disciple By The Initiation He Imparts To Him And The Other Means (Yukti) N He Adopts, Even So, He Is Not The Only Guide On The Path. Apart From The Master There Is Scripture And, Above All, One's Own Personal Experience, Because, As Abhinava

Says: The Knowledge [Acquired] By Gradually [Coming To Understand The Meaning Of] The Scriptures And Following The Master [Who Knows Them] Leads, [When] Confirmed For Oneself, To The Realisation Of One's Own Identity With Bhairava. It Is Important To Know The Scriptures. God Reveals Himself Through Them; They Are One Of The Forms In Which He Is Directly Apparent In This World. They Teach Man What Is Worth Attaining And What Should Be Avoided And So Like A Boat Convey Him Across The Ocean Of Profane Existence (Samsara) To The Other Shore Where God's True Nature Is Revealed To Him. However, The Study Of The Scriptures Is Of Value Only If Accompanied By The Spiritual Knowledge That Results From Personal Experience. Mahesvarananda Writes: Being Well Versed In The Nature Of Deity Is One Thing, But Being Well Versed In The Sacred Scripture Is Another, Just As The Peace Of That Abode Is One Thing And What Worldly People Experience Is Another. Vasugupta, Who Found The Sivasutra, Knew The Means To Realisation (Yukti) As Well As The Scriptures And Had Fully Experienced The One Ultimate Reality. Therefore, Ksemaraja Declares Him To Be Amongst The Best Of Teachers. The Stanzas On Vibration (That Ksemaraja Attributes To Vasugupta) Accordingly Transmit The Secrets Of The Sivasutra In Accord With Scripture, Sound Reasoning And Personal Experience. The Latter Is Particularly Important For The Spanda Yogi; He Is Not Interested In Wasting His Time In Useless Discussion About The Experience Of Consciousness Expansion And Its Fruits, For That Can Only Be Known For Oneself. The Yogi Can Achieve This Experience Either Through Faith In The Master Or Personal Insight (Svapatyayatah) Acquired By Unswerving Devotion To God. Ksemaraja Accordingly Quotes A Passage From The Bhagavadgita Where Krsna Says: Those I Deem To Be The Best Yogis Who Fix Their Thoughts On Me And Serve Me, Ever Integrated [In Themselves], Filled With The Highest Faith. But While The Yogi's Development Depends On Faith And Personal Experience Of The Higher States Of Consciousness, He Can, And Must, Strengthen His Conviction In The Light Of Reason. When Reason (Upapatti) And Direct Insight (Upalabधि) Work Together, They Serve As A Means To Liberation. Reason Alone Cannot Help Us, But When It Is Based On An Intuitive Insight Of Fundamental Principles Along With A Direct Experience Of Reality, Error Is Eradicated And The Yogi Is Freed. In This Way The Awakened Yogi Realizes His Inherent Spiritual Power (Svabala) With Which He Exerts Himself To Distinguish Between The **Motions Of Individualized Consciousness And The Universal Vibration (Samanyaspanḍa)** Of The Collective Consciousness That Is Their Ultimate Ground And Firm Foundation. Note The Information Of Individual, Collective And Cosmic General Ledger Is Alone Responsible For Creation And Also Destruction. Thus, Although The Doctrine Taught In The Stanzas On Vibration Accords With Scripture, It Is Supported By Reason And Above All By Personal Experience. Thus, For Example, The Seventeenth Stanza Describes The Difference In The Manner In Which The Well Awakened And The Unawakened Experience Their Own Nature (Atmopalambha While The Eighteenth Describes The Experience Of The Well Awakened In The Three States Of Waking, Dreaming And Deep Sleep. Indeed, Rajanaka Rama, One Of The Commentators, Explains That The First Sixteen Stanzas Establish On The Basis Of Personal Experience (Svanubhava) That One's Own True Nature Is Independent Of The Body. Similarly, The Remaining Stanzas Also Discuss The Direct Experience Of One's Own Nature, But This Time As The Unity Of All Things. This Direct Experience, In Its Diverse Aspects, Is Both The Means By Which The Yogi Develops His Consciousness As Well As His Ultimate Goal. **Doctrine Of Vibration: Mark S.G.Dyczkowski Models See Next Paper** From the point of view of the object, the expansion (unmesa) of this pulse is represented (eb) by the initial desire to perceive (didrkṣa) a particular object, while the contracted (nimesha) phase is (=) the withdrawal **of attention** from the object previously perceived. From the point of view of the perceiving subjectivity, the phases are (=) reversed, so that the initial desire to perceive marks (eb) the contraction (nimesha) of subjective consciousness while the falling away of the previous perception is (=) its expansion (unmesa). At the higher level, where **these two phases are experienced within (eb) consciousness**, they represent (eb) the state of the categories of Isvara ('this universe is me') and Sadasiva ('I am this universe'). Utpaladeva says: Expansion (unmesa\ which is in (eb) the external manifestation [of objectivity], is (=) Kvaratattva while contraction (Niemba), which is in the internal manifestation [of subjectivity], is Sadasiva. ***There is no glorification or mortification here. It just like joining the job and retiring the involution and evolution phases.*** (italics mine) The Doctrine of Vibration At this level all the powers of consciousness fuse (e&eb) and both phases are (=) manifest as part of one reality. This unity is in fact apparent (eb) to everybody at each moment. However, within the domain of Maya, which is the sphere of differentiated perceptions (Yikalpa), it is clearly

manifest (eb) only at the juncture (Madhya) **between two cognitions**. In this Centre resides (eb) the void (kha) of consciousness (free of (e) thought-constructs) which divested of (e) diversity, digests into (eb) itself all the psycho-physical processes that give (eb) life to the multiplicity of perceptions. **Here it is important to note that the when the cosmic general ledger zero reflects upon itself and it is projected on the individual consciousness you see a film an illusory animations wherein creation itself becomes subjective wherein the individual consciousness is both the perceiver the perceived being the image on the screen on the screen of individual consciousness**. The yogi moves from the particular vibrations of consciousness at its periphery to (e&eb) the universal throb of the Heart in the Centre. As Abhinava explains: The self-reflective awareness in (eb) the Heart of pure consciousness, present (eb) at the beginning and end of each perception, within (eb) which the entire universe is (=) dissolved away without residue, is (=) called in the scriptures, the universal vibration of consciousness (samanyaspana) and is (=) the outpouring (uccalana) [of awareness] within one's own nature. All the categories of existence (tattvas) are united in (eb) the Heart of the Centre where the life-giving elixir of Siva's consciousness floods (e&eb) one's own inner nature. To reside in the Centre is to abide by (e) the law of totality (gramadharmā) in a state which transcends (e) the workings of the mind (unmana). Consciousness (jnana) with Light as its support, residing in (eb) the Centre between being and non-being is (=) known as the act of abiding in (eb) one's own abode as (=) the perceiving subjectivity (dratftrva) free of (e) all obscuration. That which has been purified by (e) pure awareness (**suddhavi(jnana)**) is called the transcendent (viviktavastu), said to be (=) the mode of being (v/7//) of the law of totality (gramadharmā) through (e&eb) which everything is (=) easily attainable. The power in the Centre (madhyasakti) is (=) the eternal Present Beyond time it is the source of both past and future. To be established there is to abide without a break in Rama, the supreme enjoyer, in every action of one's life. Rama is Siva, the supreme cause Who pervades the fourteen aspects which embrace the entire universe of experience, namely, moving, standing, dreaming, waking, the opening and closing of the eyes, running, jumping, exertion, knowledge [born] of the power of the senses, the [three] aspects of the mind, living beings, names and all kinds of actions. This came as a natural development in Spanda doctrine not only for **this reason but also because the universal ego is experienced as the inner dynamics of absolute consciousness. To conclude our summarial exposition of the Divine Means, which is centred on the direct experience of this pure ego (and hence on Spanda in this form), we turn now to a brief description of its inner, cyclic activity. We shall do this** by examining Abhinava's esoteric exegesis of the symbolic significance of the word 'A HAM', which in Sanskrit means T, and symbolises by its form the ego's dynamic nature. "Dasein Does Not Fill Up A Track Or Stretch 'Of Life' — One Which Is Somehow Present-At-Hand — With The Phases Of Its Momentary Actualities. It Stretches Itself Along In Such A Way That Its Own Being Is Constituted In Advance As A Stretching-Along. The 'Between' Which Relates To Birth And Death Already Lies In The Being Of Dasein ... It Is By No Means The Case That Dasein 'Is' Actual In A Point Of Time, And That, Apart From This, It Is 'Surrounded' By The Non-Actuality Of Its Birth And Death. Understood Existentially, Birth Is Not ... Something Past In The Sense Of Something No Longer Present-At-Hand; And Death Is Just As Far From Having The Kind Of Being Of Something ... Not Yet Present-At-Hand But Coming Along ... Factual Dasein Exists As Born; And, As Born, It Is Already Dying, In The Sense Of Being-Towards-Death. As Long As Dasein Factually Exists, Both The 'Ends' And Their 'Between' *Are*, And They Are In The Only Way Possible On The Basis Of Dasein's Being As *Care* ... As Care, Dasein Is The 'Between'." — **Martin Heidegger (Thomas Piel - Traducteur), Being And Time** Path to Liberation On without knowledge that everything is manifest within consciousness is illusory or unreal in that sense alone. 78 Things are more real or more tangibly experienced according to their own essential nature (svabhava) to the degree in which we recognise that they are appearances (abhasa) within absolute consciousness As Jayaratha says: Just as images manifest in a mirror, for example, are essentially mere appearances, so too are [phenomena] manifest within conscious- ness. Thus, beause they are external, [phenomena] have no being (sattva) of their own. The Lord says this [not with the intention of saying anything about the nature of things] but in order to raise the level of consciousness of those people who are attached to outer things; thus everything in this sense is essentially a mere appearance. [Knowing this], in order to quell the delusion of duality, one should not be attached to anything external. 79 The ultimate experience is the realisation that everything is contained within consciousness. We can discover this in two ways. Either we merge the external world into the inner subject, or we look upon the outer as a gross form of the inner. In these two ways we come to recognise that all things reside within our own

consciousness just as consciousness resides within them. This all-embracing inwardness is only possible if there is an essential identity between the universe and consciousness. The events which constitute the universe are always internal events happening within consciousness because their essential nature is consciousness itself. 80 We can only account for the fact that things appear if there is an essential identity between consciousness and the object perceived. If a physical object were really totally material, that is, part of a reality independent of, and external to, consciousness, it could never be experienced. Abhinava says: The existence or non-existence of phenomena within the domain of the empirical (iha) cannot be established unless they rest within consciousness. In fact, phenomena which rest within consciousness are apparent (prakasamana). And the fact of their appearing is itself their oneness (abheda) with consciousness because consciousness is nothing but the fact of appearing (prakasa). If one were to say that they were separate from the light of [that consciousness] and that they appeared [it would be tantamount to saying that] 'blue' is separate from its own nature. However, [insofar as it appears and is known as such] one says: 'this is blue'. Thus, in this sense, [phenomena] rest in consciousness; they are not separate from consciousness.

The Doctrine of Vibration The universe and consciousness are two aspects of the whole, just as quality and substance constitute two aspects of a single entity. The universe is an attribute (dharma) of consciousness which bears (dharmiri) it as its substance. It is said that 'substance' is that resting in which this entire group of categories manifests and is made effective. Now, if you don't get angry [we insist that] this entire class of worlds, entities, elements and categories (tattva) rests in consciousness and [resting in it] is as it is. Thus consciousness contains everything in the sense that it is the ground or basis (adhard) of all things, their very being (satta) and substance from which they are made. But, unlike the Brahman of the Advaita Vedanta, it is not the real basis (adhithand) of an unreal projection or illusion. Consciousness and its contents are essentially identical and equally real. They are two forms of the same reality. Consciousness is both the substratum and what it supports: The perceiving awareness and its object. In this respect, the Kashmiri Saiva is frankly and without reserve an idealist. Although he does not deny the reality of the object, his position is at odds with most commonly accepted forms of realism. The realist maintains that the content perceived is independent of the act of perception. The content is only accidentally an object of perception and undergoes no change in the process of being perceived. His contention, however, is essentially unverifiable; to verify it, we would have to know an object without perceiving it. This, from the Kashmiri Saiva point of view, is not possible. Objects of which we have no knowledge may indeed exist, but they are knowable as objects only if they are related to subjects who perceive them. In this sense, if there were no subjects, there could be no objects. 86 The subject, however, as opposed to the object is, in terms of the phenomenology of perception, apparent to himself. He is self-luminous (svaprakasa). Thus, consciousness (the essence of subjectivity) is one's own awareness by virtue of which all things exist. The realist maintains that consciousness clearly differs from its object insofar as their properties are contrary to each other. The Saivite idealist. However, says that the object is a form of awareness (vijñanakara)TM The objective status of the object is cognition itself. Perception manifests its object and renders it immediately apparent (sphuta) to those who perceive it. It does not appear at any other time. If 'blue' were to exist apart from the cognition of blue\ two things would appear: 'blue' and its cognition, which is not the case. It is the perception of the object which constitutes its manifest nature. An entity becomes an Integral Monism of Kashmiri Saivism object of knowledge not by virtue of the entity itself but by our knowledge of it. If objects had the property of making other objects appear, it would be possible for one object to make another appear in its own likeness. 'Blue' is perceived to be 'blue' because it is manifest as such to the perceiver. As Abhinava points out: The [nature of an] object of knowledge could not be established through a means of knowledge totally unrelated to it — a crow does not become white because a swan [sitting next to it] is white. Perception, on the other hand, is immediately apparent to consciousness. It is self-luminous in the sense that it is directly known without need of being known by any ulterior acts of perception and makes its object known at the same time. Adopting the Buddhist Yogacara doctrine that things necessarily perceived together are the same (sahopalambhaniyamavada), the Saivite affirms that because the perceived is never found apart from perception, they are in fact identical. Reality (satya) is the point where the intelligible and the sensible meet in the common unity of being; it cannot be said to exist in itself outside, and apart from, knowledge or vision. Bhagavatopala in his commentary on the Stanzas on Vibration quotes: Once the object is reduced to its authentic nature, one knows [the true nature of] consciousness. What then [remains of] objectivity? What [indeed could be] higher than

consciousness? Consciousness is essentially active. Full of the vibration of its own energy engaged in the act of perception, it manifests itself externally as its own object. When the act of perception is over, consciousness reabsorbs the object and turns in on itself to resume its undifferentiated inner nature. Knowledge (jnana) manifests internally and externally as each individual entity.... Once knowledge has assumed that form it falls back [into itself]. The Yogacara Buddhist similarly maintains that consciousness creates its own forms. But, according to him, because the perceived and perception are identical; there is no perceived object at all. The so-called outer world is merely a flux of cognitions, it is not real. He is firmly committed to a doctrine of illusion. The reality of consciousness from The Doctrine of Vibration his point of view is established by proving the unreality of the universe. "All this consists of the act of consciousness alone", says Vasubandhu, "because unreal entities appear, just as a man with defective vision sees unreal hair or a moon, etc." He points to dreams as examples of purely subjective constructs which appear to be objective realities. The apparent reality dreams possess is not derived from any concrete, objective world, but merely from the idea of objectivity. While the Yogacara does not say that an idea has, for example, spatial attributes, it does have a form manifesting them. While he agrees with the Saiva idealist that appearances have no independent existence apart from their appearing to consciousness, he maintains that for this reason they are unreal. The creativity of consciousness consists in its diversification in many modes having apparent externality; it is not a creation of objects. While the Kashmiri Saivite agrees that the world is pure consciousness alone, he maintains that it is such because it is a real creation of consciousness. The effect is essentially identical with the cause and shares in its reality. Matter and the entire universe are absolutely real, as 'congealed' (sty ana) or 'contracted' (samkucita) forms of consciousness. "This God of consciousness", writes Ksemaraja, "generates the universe and its form is a condensation of His own essence (rasa)" m By boiling sugarcane juice it condenses to form treacle, brown sugar and candy which retains its sweetness. Similarly, consciousness abides unchanged even though it assumes the concrete material form of the five gross elements. The same reality thus abides equally in gross and subtle forms. Consequently no object is totally insentient. Even stones bear a trace (vasana) of consciousness, although it is not clearly apparent because it is not associated with the vital breath (prana) and other components of a psycho-physical organism. Somananda goes so far as to affirm that physical objects, far from being insentient, can only exist insofar as they are aware of themselves as existing. The jar performs its function because it knows itself to be its agent. Indeed, all things are pervaded by consciousness and at one with it and hence share in its omniscience. Thus, Siva, Who perceives Himself in the form of physical objects, is the one ultimate reality. "The jar knows because it is of my nature", writes Somananda, "and I know it because I am of the jar's nature. I know because I am of Sadasiva's nature and He knows because He is of my nature; Yajnadatta [knows] because he is of Siva's nature and Siva [knows] because He is of Yajnadatta's nature". Integral Monism of Kashmiri Saivism Everything in this sense is directly perceived by absolute consciousness, and this direct perception (pratyaksa) unifies the knowable into a single, undivided whole. This is the central concept behind a doctrine originally expounded by Narasimha called 'the non-dualism of direct perception' (pratyaksadvaita). This states that consciousness is essentially perceptive and that its perception of all things operates throughout the universe. Insofar as phenomena are clearly evident (sphufa) to us, everything is directly perceived by absolute consciousness, with which our individual consciousness is identical. This direct perception unfolds everywhere; the one true reality, it is alone and without companion or rival (nihsapatna). Even though it remains one, it can, by its very nature, perceive distinctions (bheda) between one entity and another, without this engendering any division within it. We distinguish between two entities in empirical terms on the basis of their mutual exclusion (anyonyabhava). The relative distinction {bheda} between them is essentially the perceived difference between their respective characteristics. Despite this difference they are united within the purview of a single cognition insofar as they are equally both manifest appearances. This cognition is the undivided essence (rasa) or 'own nature' (svabhava) of both. Encompassed by the 'fire of consciousness', there is no essential difference between them. Just as when an emerald and ruby reflect each other's light, the ruby is reddish-green and the emerald greenish-red, similarly everything is connected with everything else as part of the single variegated (vicitra) cognition of absolute consciousness. Mahesvarananda writes: The Supreme Lord's unique state of emotivity (asadharanabhava) is the outpouring of pure Being (mahasatta). It is manifest as the brilliance (sphuratta) of the universe which, if we ponder deeply, [is realized to be] the single flavour (ekarasa) of the essence of Beauty which is the vibration of the bliss of one's own nature. In

this way all things are in reality one although divided from the one another sharing as they do the 'single flavour' (ekarasa) of the pure vibration of consciousness. Kashmiri Saiva Realism Kashmiri Saivism as a whole has been variously called a form of 'realistic idealism', 'monistic idealism', 'idealistic monism' and 'concrete monism'. It is easy to understand why Kashmiri Saivism is The Doctrine of Vibration said to be 'idealistic' and 'monistic', but in what sense is it also 'realistic'? The answer to this question is of no small importance in trying to understand the central idea behind its metaphysics and the fundamental importance of the concept of Spanda, in this seemingly impossible marriage between monistic idealism and pluralistic realism. The Kashmiri Saiva approach understands the world to be a symbol of the absolute, that is, as the manner in which it presents itself to us. Again we can contrast this view with that of the Advaita Vedanta. The Advaita Vedanta understands the world to be an expression of the absolute insofar as it exists by virtue of the absolute's Being. Being is understood to be the real unity which underlies empirically manifest separateness and as such is never empirically manifest. It is only transcendently actual as 'being-in-itself. The Kashmiri Saiva position represents, in a sense, a reversal of this point of view. The nature of the absolute, and also that of Being, is conceived as an eternal becoming (satatodita), a dynamic flux or Spanda, 'the agency of the act of being'. It is identified with the concrete actuality of the fact of appearing, not passive unmanifest Being. Appearance (abhasa) alone is real. appearing (prakasamanatva) is equivalent to the fact of being (astitva). Ksemaraja writes in his commentary on the Stanzas on Vibration: Indeed, all things are manifest because they are nothing but manifestation. The point being that nothing is manifest apart from manifestation The absolutely unmanifest, from this point of view, can have as little existence as the space in a lattice window of a sky-palace. Nay, even less, because even that space can appear as an imagined mage manifest within consciousness. Everything is real according to the manner in which it appears. Even an illusion is in this sense real, insofar as it appears and is known in the manner in which it appears. The empirical and the real are identical categories of thought. As Abhinava says: Thus this is the supreme doctrine (upanisatf), namely that, when- ever and in whatever form [an entity] appears, that then is its particular nature. Perhaps at this stage a brief comparison with Heidegger's ideas might prove to be enlightening and not altogether out of place. According to Heidegger's phenomenology of Being, reality is intelligible in a two-fold Integral Monism of Kashmiri Saivism manner as 'phenomenon' and 'logos'. Heidegger defines what he means by 'phenomenon' as: "that-which-shows-itself. The manifest . . . phenomena are then the collection of that which lies open in broad daylight or can be brought to the light of day — what the Greeks at times implicitly identified as 'ta onta' (the things-which-are)". 127 In his later writings Heidegger drops the term 'phenomenon' in preference for the verbal form 'phainesthai' in order to emphasize even more the actuality or presentational property of Being. Explaining this new form of the term he writes: "Being disclosed itself to the ancient Greeks as 'physis'. The etymological roots 'phy-' and 'pha-' designate the same thing: 'phyein', the rising-up or upsurge which resides within itself as 'phainesthai', lighting-up, self-showing, coming-out, appearing-forth." Heidegger contrasted his notion of phenomenon with semblance (Schein) and with appearing (Erscheinung). In the case of semblance a thing can show itself as that which it is not, as when fool's gold shows itself to be gold. The ancients always allied semblance with non-being. Heidegger points out, however, that semblances are grounded in showings, and so does Abhinava. Both Heidegger and Abhinava consequently maintain that all semblances have a real basis and are to be treated as instances of phenomena along with the so-called real showing or manifestation of non-deceptive objects. So Heidegger states that: 'how- ever much seeming, just that much being'. Thus self-showing or appearing defines Being as phenomenon, but this definition of Being is as yet incomplete. Being is not only self-showing but 'logos' which Heidegger explains means 'discourse' (Rede) in the sense of 'apophansis': 'letting-be-seen'. Phenomenology, which according to Heidegger is the only correct study of Being, means 'letting-be-seen-that-which- shows-itself. This is true of Saiva Paramadvaita as well. The reality of the world demands recognition; we are forced to accept the direct presentation of the fact of our daily experience. As Abhinava says: "if practical life, which is useful to all persons at all times, places and conditions were not real, then there would be nothing left which could be said to be real." A thousand proofs could not make 'blue' other than the colour blue. The reality of whatever appears in consciousness cannot be denied. Objects appear; they do not cease to do so by a mere emphatic denial. The manifestation of an entity in its own specific form is a fact at one level of consciousness; it is real. The appearing of the same entity in the same form but recognised to be a direct representation of the absolute is also a fact, but at another level of consciousness. It is no more or less real than the first. 'As is the state of consciousness,

so is the experience,' says Abhinava. Although the nature of the absolute is discovered at a higher level of consciousness, The Doctrine of Vibration nonetheless it presents itself to us directly in the specific form in which we perceive things; otherwise there would be no way in which we could penetrate from the level of appearing to that of its source and basis. Abhinava writes: Real is the entity iyastu) that appears in the moment of direct perception (sak\$atkara), that is to say, within our experience of it. Once its own specific form has been clearly determined one should, with effort, induce it to penetrate into its pure conscious nature. All things are known to be just as they present themselves. The concrete actuality of being known (pramiti), irrespective of content, is itself the vibrant (spanda) actuality of the absolute. Liberating knowledge is gained not by going beyond appearances but by attending closely to them. "The secret," Mahesvarananda says, "is that liberation while alive (jivanmukti) is the profound contemplation of Maya's nature." No ontological distinction can be drawn between the absolute and its manifestations because both are an appearing (dbhasa), the latter of diversity and the former of 'the true light of consciousness which is beyond Maya and is the category Siva'. Those who have attained the category of Pure Knowledge above Maya and have thus gone beyond the category of Maya, see the entire universe as the light of consciousness . . . Just as the markings [on a feather] are nothing apart from the feather, the feather [is nothing apart from] them, similarly, when the light of consciousness is manifest, the whole group of phenomena is manifest as the light of consciousness itself. Within the sphere of Maya, every entity's 'own nature' (svabhdva) corresponds to its specific manifest form. Accordingly it is defined as that which distinguishes it from all else and from which it never deviates. Above the sphere of Maya, that is, above the level of objectivity, is the domain of the subject. At this level, everything is realised to be part of the fullness of the experience r and hence no longer bound by the conditions which impinge on the object. Here the part is discovered to be the whole, that is, consciousness in toto. In this sphere beyond relative distinctions, the yogi realises that (all) the categories of existence are present in every single category. m The yogi experiences every individual particular as the sum total of everything else. He recognises that all things have one nature and that every particular is all things. This is the 'essence' (sard) or co-extensive unity (samarasya) of all things. Integral Monism of Kashmiri SaivismWe have established that reality is manifest according to how [and the degree in which] the freedom of consciousness reveals it and that [this freedom] is the womb of all forms. Just as 'sweetness' is present in its entirety in every atom of the sugarcane, so each and every atom [of the universe] bears within itself the emanation of all things. This is the level of consciousness in which the absolute reflects on itself realising to its eternal delight and astonishment (camatkara) its own integral nature. The reality of the world of diversity is not denied, but experienced in a new mode of awareness free of time and space in the eternal omnipresence of the Here and Now. [Phenomenal forms of awareness] such as 'this [exists]', born of the colouring [imparted to the absolute] by the limitations engendered by the diversifying power of time {kalakalana) also emanate within the Supreme Principle. There [at that level], Fullness {purriata) is the one nature [of all things] and so everything is omnipresent; otherwise, associated with division (khancjlana), the Fullness [of the absolute] would not be full. The content of absolute consciousness consists of diverse appearances (abhdsa) which, because they are manifest through it in this way, do not compromise the wholeness of consciousness. Everything we perceive is a momentary collocation of a number of such manifestations which combine together like 4 a row of altar lamps' (dipavalT) to form the single radiant picture of the universe. The individual objects which constitute the universe are specific collocations of such 'atomic' appearances. Together they form a single unified particular which appears according to its own defining features (svalaksand). A jar, for example, consists of a number of appearances such as 'round', 4 fat\ 'earthen', 'red', etc., which together discharge a single function (arthakriya), in this case, that of carrying the appearance 'water'. They unite with each other much as the scattered rays of a lamp come together when focused, or as the various currents of the sea together give rise to waves. Atomic appearances can combine in any number of ways, provided that they are not contrary to one another as established by the dictates of natural law (niyati). An appearance of 'form', for example, cannot combine with that of 'air'. Insofar as they share a common basis (samanyadhikaranya), a given cluster of appearances appears as a single whole. This common basis is the most prominent member of the group; the appearance 'jar' is such in the example quoted above. Any one appearance in a cluster may assume a more important or subordinate role. The result is a specific The Doctrine of Vibration awareness of an object of the form: 'here this is such.' While individual appearances do not lose their separate identity {svarupabheda) when they rest on a common basis, even so the particular object which appears according to

its own characteristics (svalakṣaṇa) is an individual reality in its own right. It is a different kind of appearance characterised by its association with the appearance of the specific location and time in which it is made manifest. The form of our experience is thus 'I now see this here'. But when we perceive each particular constituent appearance separately, each assumes a separate fixed function. Abhinava cites the following colourful example to illustrate how the various combinations of appearances account for the variety of experience: Thus even though the appearance of the beloved may manifest externally, it is as if far away in the absence of another appearance, namely, that of 'embracing'. So when the [appearing of the beloved] is associated with another appearance [namely that of 'far away'] the power (arthakriya) it formerly had of giving pleasure appears as its contrary. The form our experience assumes depends, not only on the nature of the object perceived, but also on personal factors entirely peculiar to ourselves. This theory explains this in two ways. In one sense, the object remains the same, but one or other of its constituent appearances comes to the fore according to the inclinations of the perceiver. From another point of view, we can say that the perceived object is different for each perceiver according to the difference in the prominent appearance manifest to him. Abhinava, citing as an example a golden jar, illustrates how the same object appears differently to different perceivers according to the use they wish to make of it and to their state of mind: When a person who is depressed and feels that there is nothing [of value for him in the world] sees the jar, he merely perceives the appearance 'exists' [in the form of the awareness that] 'it is'. He is not conscious of any other [of its constituent appearances] at all. An individual who desires to fetch water [perceives] the appearance 'jar'. The man who simply wants something that can be taken somewhere and then brought back [perceives] the appearance 'thing'. The man who desires money [perceives] the appearance 'gold'. The man who desires a pleasing object [perceives] the appearance 'brightness' while he who wants something solid sees the appearance 'hardness'. These 'atomic events' or appearances emerge from the pure subject's consciousness and combine together to form a total event at each moment. Integral Monism of Kashmiri Saivism Daily life (vyavahara) goes on by virtue of this ever renewed flux of appearances. They are connected together and work towards a single unified experience because they appear within the field of consciousness of the universal subject. The aggregate of appearances arises in the [supreme] subject as do [sprouts in] a rice field. Even though each sprout germinates from its own seed, they are perceived as a collective whole. Appearances rest in this way within the universal subject. 'Externality' is itself another appearance; it arises from a distinction between appearances and the individual subject. So, although all manifestation always occurs within the subject, it appears to be external due to the power of Maya which separates the individual subject from his object. This split must occur for daily life to be possible. Only externally manifest appearances can perform their functions; when they are merged within the subject and at one with him, they cannot do so. Daily life proceeds on the basis of the operation and withdrawal of the conditions necessary for fruitful action to be possible. Appearance in this sense represents the actualisation of a potential hidden in consciousness made possible by virtue of its dynamic, Spanda nature which is both the flow from inner to outer and back as well as the power that impels it. The emergence from, and submergence into, pure consciousness of each individual appearance is a particular pulsation (visesaspaṇḍa) of differentiated awareness. Together these individual pulsations constitute the universal pulse (samanyaspaṇḍa) of cosmic creation and destruction. Thus, every single thing in this way forms a part of the radiant vibration (sphuratta, sphurana) of the light of absolute consciousness. Light and Awareness: The Two Aspects of Consciousness Absolute consciousness understood as the unchanging ontological ground of all appearing is termed 'Prakasa'. As the creative awareness of its own Being, the absolute is called 'Vimarsa'. Prakasa and Vimarsa — the Divine Light of consciousness and the reflective awareness this Light has of its own nature — together constitute the all-embracing fullness (purnata) of consciousness. The Recognition (pratyaabhijñā) school of Kashmiri Saivism develops this concept of the absolute which finds its fullest expression in Utpaladeva's Stanzas on the Recognition of God. Even though neither of these two key terms appear in the Stanzas on Vibration or the Aphorisms of Siva, they recur frequently in their commentaries. Thus, although the original formulation of the Doctrine of Vibration differs from the theology of Recognition in this respect, it was extended in the course of its development to accommodate this concept of the absolute as well. This was possible, and quite justified, insofar as the absolute understood in Pratyabhijñā terms does not, as we shall see, differ essentially from that of the Spanda school. We can, as Kashmiri Saivites themselves have done, explain one in terms of the other. The Doctrine of Vibration Prakasa: The Light of Consciousness Prakasa is the pure luminosity' (bhdna) or 'self-showing' that constitutes the

essence and ultimate identity (atman) of phenomena. That things appear at all is due to the light of consciousness, and their appearing (avabhasana) is itself this Light which bestows on all things their evident, manifest nature. Established in the light of consciousness everything appears there according to its own specific nature (svabhava). Anything that supposedly does not rest in this Light is as unreal as a sky-flower. 3 Thus, according to Rajanaka Rama, unlike the light of the sun, or any other light, this Light not only makes all things apparent, it is also their ultimate source. 4 Full of its divine vibration the Light makes all things manifest and withdraws them into itself. This supra- temporal activity characterises it most specifically; devoid of it, it would be no better than an inert physical phenomenon. At the same time, this light is the conjunction (slesa) or oneness (aikdmya) of its countless manifest forms, 6 and the collective whole (sarpina'ana) of all the categories of existence. The universe is nothing but the shining of the Light within itself. It is the radiant vibration (sphuratta) of this Light, the state (avast hand) in which consciousness becomes manifest. Although the Light shines as all things at all times and hence also makes their diversity manifest, 9 penetrating each object individually as well as collectively, it is not totally 'merged' (magna) or identified with the object so as to suffer any division within itself. Our experience of any object is of the form: 'I see this': it is not itself an object, but the manifest form the object assumes as a luminous principle of experience. The Light is ever revealed and can never be obscured; objectivity can never cast a shadow on the light of consciousness. The Stanzas on Vibration declare: That in which all this creation is established and from whence it arises is nowhere obstructed because it is unconditioned by [its very] nature. This Light is the highest reality (paramartha). It is the 'Ancient Light' (puranaprakasa) that makes all things new and fresh every moment. It is 'always new and secret, ancient and known to all'. It is the form of the Present (vartamanarupa), the Eternal Now. Time and space are relations between the contents of consciousness; they cannot impinge on the integrity of the absolute itself. 16 Neither space nor time can divide it, for they are one with the Light that illumines them Light and Awareness: Two Aspects of Consciousness and makes them known as elements of experience. But this Light is the shining of the absolute; it is not an impersonal principle. It is the living Light of God, indeed it is God Himself, the Master Who instructs the entire universe. 18 Siva is this 'auspicious lamp', Who illumines all things. He is the Light of consciousness that reveals the presence of both the real and the unreal, of light' and 'darkness'. 20 Abhinavagupta writes: Thus Bhairava, the Light, is self-evident (svatahsiddha); without beginning, He is the first and last of all things, the Eternal Present. And so what else can be said of Him? The unfolding of the categories of existence (tattva) and creation, which are the expansion of His own Self, He illumines, luminous with His own Light, in identity with Himself, and because He illumines Himself, so too He reflects on His own nature, without His wonder (camatara) being in any way diminished. Since Zwicky's early observations, similar data from other galaxy clusters has yielded the same result – we consistently see that clusters of galaxies appear to By developing an awareness of the Centre, the yogi experiences the bliss of consciousness. 117 Through this gap he plunges into introverted absorption (nimilanasamadhi) and then emerges again to pervade the field of awareness between Centres and so experience the Cosmic Bliss (jagadananda) of the universal vibration of consciousness. 118 He then recognises that this state pervades every aspect of experience. In this way the yogi's consciousness is no longer afflicted by the power which obscures it, hemming the Centre in on both sides with thought-constructs that seemingly deprive it of its fullness. As he realises directly his pure conscious nature as the universal ego free of all mental representations, it expands out to embrace all things within itself. Thus the realisation the Divine Means leads to, and is directly based upon, is that this pure ego is in all things just as all things are within it. In the Spanda tradition, as recorded in the Stanzas on Vibration, no such ego is recognised. '19 Man's authentic nature is, however, understood in personal terms as every individual's own 'own nature' (svasvabhava) which is Siva, the universal vibration of pure subjectivity (upalabdhta). It is not surprising, therefore, that later commentators found these two conceptions to be essentially the same and accordingly identified one's own inner nature with the pure ego. This came as a natural development in Spanda doctrine not only for **this reason but also because the universal ego is experienced as the inner dynamics of absolute consciousness. To conclude our summarial exposition of the Divine Means, which is centred on the direct experience of this pure ego (and hence on Spanda in this form), we turn now to a brief description of its inner, cyclic activity. We shall do this** by examining Abhinava's esoteric exegesis of the symbolic significance of the word 'A HAM', which in Sanskrit means T, and symbolises by its form the ego's dynamic nature. The objective world of perceptions is, as we have seen, essentially a chain of thought-constructs

(prapanca) closely linked to one another and woven into the fabric of diversity (vicitrata). This thought (vikalpa) is a form of speech (vac) uttered internally by the mind (citta), which is itself an outpouring of consciousness. Consciousness also, in its turn, resounds with the silent, supreme form of speech {para vac} which is the reflective awareness through which it expresses itself to itself. Consequently, the fifty letters of the Sanskrit alphabet, which are the smallest phonemic units into which speech can be analysed, are symbolic of the principal elements of the activity of consciousness. Letters come together to generate words and words go on to form sentences. In the same way the fifty phases in the cycle of consciousness represent, in the realms of denoted meaning (vacya) the sum total of its universal activity (kriya) corresponding to the principal forces (kala) which come together to form the metaphysical categories of 186 The Doctrine of Vibration experience, which in their turn appear in the grossest, most explicitly 'articulate' form as the one hundred and eighteen world-systems (bhuvana). 'A', the first letter of both AHAM and the Sanskrit alphabet, is the point of departure or initial emergence of all the other letters and hence denotes Anuttara — the absolute. 'Ha', is the final letter of the alphabet and represents the point of completion when all the letters have emerged. It represents the state in which all the elements of experience, in the domains of both inner consciousness and outer unconsciousness, are fully displayed. It is also the generative, emission (visarga) which, like the breath, casts the inner into the outer, and draws what is outside inward. The two letters 'A' and 'Ha' thus represent Siva, the transcendental source and Sakti, His cosmic outpouring that flows back into Him. The combined 'A-Ha' contains within itself all the letters of the alphabet — every phase of consciousness, both transcendental and universal. (For a graphic representation of this analysis, see figure 1.) M, the final letter of AHAM, is written as a dot placed above the letter which precedes it. It comes at the end of the vowel series and before the consonants and so is called 'anus vara' (lit. 'that which follows the vowels') and also 'bindu' (lit. 'dot,' 'drop,' 'point' or 'zero'). While the consonant 'M' symbolises the individual soul (purusa), 'bindu' represents the subtle vibration of T, which is the life force (jivakala) and essence of the soul's subjectivity manifest at the transcendental, supra-mental level (unmana). 12 ° It is the zero-point in the centre between the series of **negative numbers, in this case the vowels which represent the processes happening internally within Siva**, and the series of positive numbers — the consonants which symbolise the processes happening externally within Sakti. Bindu, as a point without area, symbolises the non-finite nature of the pure awareness (pramitibhava) of AHAM. It is the pivot around which the cycle of energies from 'A' to 'Ha' rotates, the Void in the centre from which all the powers emanate and into which they collapse. As such, it is the supreme power of action which holds subject, object and means of knowledge together in a potential state in the one Light that shines as all three 121 containing them in its repose 122 (visrdnti). Bindu is the 'knower' (jndtr), who is essentially consciousness that, though omniscient does not manifest its intelligence, like a man who knows the scriptures but having no occasion to explain them to others silently bears this knowledge within himself. As such, it symbolises the union of Siva and Sakti (sivasakti- mithunapina'a) 123 in a state of heightened potency in which they have not yet divided to generate the world of diversity. It stands, in other words, at the threshold of differentiation in the stream of emanation still contained within Siva. Expansion Commences Bindu — The Individual Soul Withdrawal Commences Figure 1 188 The Doctrine of Vibration Then, to the degree in which that which is to be accomplished by the power of action residing within it [as a potential] penetrates into the absolute, it appears initially as bindu, which is the light of pure consciousness. 124 When outer objectivity is reabsorbed into its transcendent source, bindu is the point into which all the manifest powers of consciousness are gathered and fused together. The universal potency of all the letters is thus contained in bindu which, as the reflective awareness of supreme T consciousness, 125 gives them all life. Thus bindu also marks the beginning of Siva's internal movement back to the undifferentiated absolute and so stands at the threshold of both emission and absorption without being involved in either. The three aspects of AH AM together constitute a movement from the undifferentiated source of transcendental consciousness — 'A' — through the expansion or emission of its power — 4 Ha' — to the subject — 'M' — which contains and makes manifest the entire universe of experience. The reverse of this movement, which of withdrawal (samhara), is represented by M-Ha-A. AHAM and M-Ha-A alternate in the rotation (ghurnana) of the reflective awareness of T consciousness as immanent Sakti emerges from transcendental Siva to then merge back into Him. As Abhinava says: The universe rests within Sakti and She on the plane of the absolute (anutara) and this again within Sakti ... for the universe shines within consciousness and [consciousness shines] there [within the universe by the power of] consciousness. These three

poles, forming a couple and merging, make up the one supreme nature of Bhairava Whose essence is AHAM- 126 At the microcosmic level, 'A' represents the initial moment when the subject begins to rise out of himself to view the object. The movement from 'A' to 'Ha' marks the emergence of sensation within the field of awareness, which is represented by the fifty letters of the alphabet symbolic of the fifty aspects of the flux of consciousness leading to objectified perception. 'N' is the subject who, resting content within himself when he has perceived his object, merges through the inner flow of awareness into 'K' the absolute. Then from the absolute (A) its emission (Ha) flows back into the pure subject (M) set to perceive his object. Thus all the cycles of creation and destruction are contained within AHAM through which they are experienced simultaneously as the spontaneous play of the absolute. The yogi who recognises this recurrent pulse of awareness to be the movement of his own consciousness merges his limited ego with the Path to Liberation universal ego. **This is nature's general ledger** Thus he realises that its power to create, sustain and destroy all things is his own inner strength (svabhava) that he exerts effortlessly in the same state of mystical absorption (samadhi) in universal consciousness that the absolute itself enjoys. In this way he shares in the three-fold awareness Siva Himself has of His own nature which Abhinava describes as follows: 4 I make the universe manifest within myself in the Sky of Consciousness. I, who am the universe, am its creator! ' — this awareness is the way in which one becomes Bhairava. 'AH of manifest creation (sadhya) is reflected within me, I cause it to persist 1 — this awareness is the way in which one becomes the universe. The universe dissolves within me. I who am the flame of the [one] great and eternal fire of consciousness' — seeing thus one achieves peace. 127 The experience of the liberated thus coincides with the realisation of their own divine nature which, through its power, rules and guides the cosmic order. Thus this attainment (siddhi), which is liberation itself, is in the Doctrine of Vibration technically called 'Mastery over the Wheel of Energies' (cakrasvaratasiddhi) because the liberated soul, identified with Siva, now governs, as does Siva, the cycle of the powers that bring about the creation and destruction of all things. **Here we are talking of individual consciousness becoming cosmic general ledger and the liberationist realises all destruction and creation is by him. This clearly means that he is in his own world of illusion. Creation is subjective quintessentially** The Empowered Means (Saktipada) All the practices taught in the Stanzas on Vibration are internal. Whenever ritual is mentioned, it is invariably interpreted in terms of the dynamics of the inner processes the yogi experiences and implements in the course of his yogic practice. The Doctrine of Vibration, Ksemaraja affirms, 129 is concerned entirely with these inner disciplines centred, as it is, in one way or another, on consciousness or, at least, on the inner activity of the mind. Thus the Empowered Means which, like the other categories we have discussed, is entirely internal, includes an important part of Spanda practice. Spanda practice belonging to the Divine Means centres on one's own inherent nature (svabhava) as Siva, the universal perceiver and agent, that belonging to the Empowered Means on His power. Instead of arriving directly at the all-embracing emptiness of subjective consciousness, the yogi practising the Empowered Means realises his true nature through the fullness of its energy. Practising the Divine Means, the yogi plunges, as it were, straight into the fire of consciousness; practising the 190 The Doctrine of Vibration Empowered Means he merges with its rays. Either way the yogi is centred equally on ultimate reality. The power of consciousness is no less absolute than its possessor. To make this point Abhinava quotes the Matanga- tantra: This reality consists of the rays of [Siva's] power and is variously said to be the abode of the Lord's manifestation . . . That same [power] illumined [by Siva] is itself also luminous, unshaken and unmoving. That very [power] is the supreme state, subtle, omnipresent, the nectar of immortality, free of obscuration, peaceful, yearning for pure Being alone (vastumatra) and devoid of beginning and end. Perfectly pure, it is said to be the body [of ultimate reality]. 110 The yogi concentrates on the powers operating in all of life's activities as particular pulsations (visvaspada) in the universal rhythm (samanyaspada) of the power of consciousness. In this way he rises progressively from the particular to the universal until he reaches pure Being (satta), the greatest of all universals (mahasamanyd) and the highest form of Siva's power. Thus the creative power of Maya, manifest through countless lesser powers, no longer causes the yogi to stray from Siva's consciousness but becomes the means through which it can be realised 131 in the illuminating brilliance (sphuratta) which is Siva's pure Being. Thus by discovering the true nature of Sakti, the yogi realises himself to be Siva, its possessor Who consists of all its countless powers. Thus practice belonging to this Means leads to the same pure consciousness free of thought-constructs realised through the Divine Means. Although the ultimate realisation is instantaneous, the yogi rises to it gradually by freeing his consciousness of the limitations imposed upon it by

thought. Abhinava explains: The same occurs in the Empowered Means [as does in the Divine]. At the discursive level of consciousness {yaikalpikibhumi} [where the Empowered Means functions] knowledge and action, although evident, are, for the reasons explained previously, contracted. A blazing energy [is revealed within] the one who dedicates himself to removing the burden of this contraction. [This energy eventually] brings about the inner manifestation {antarabhasa} of pure consciousness he seeks. 132 Consciousness is individualized and its power of knowledge and action contracted by the thought-constructs born of ignorance. The arising of these mental representations, as the Stanzas on Vibration say, deprives the soul of its freedom and immortal life. 133 The practise of the Path to Liberation 191 Empowered Means is meant to free the fettered soul of this constriction on his consciousness. It operates within the mental sphere (cetasyaTM) and is designed to purify thought (vikalpasamskara) in order to reveal the pure consciousness which is its ground and ultimate source. Thus, the Empowered Means is concerned with the second instant of perception, during which the subject forms mental representations of his object. Thought functions on the basis of an awareness of relative distinctions between specific particulars, distinguishing them from one another and thus seemingly fragmenting the essential unity of reality. 135 The vibrant vitality of consciousness, universally manifest, is clouded like a mirror by a child's breath 136 and the soul is deprived of the liberating intuition of the one reality free of thought-constructs (nirvikalpa). Abhinava writes: The [fettered soul] is like a dancing girl who although wishing to leave the dancehall is collared by the doorkeeper of thought and thrown back onto the stage of Maya. 137 All thought is centred on objectivity and hence dislodges awareness from the plenitude of pure subjective consciousness. Thus, to regain the original state of rest (visranti) consciousness enjoys, the yogi must rid himself of thought. As thought-forms decrease, pure, thought-free awareness is strengthened 138 until the yogi is fully established in a state in which the relative distinctions (bheda) conceived between entities dissolve away. Everything appears to him as pure Being (sattdmdtra) 139 and the entire universe shines before him pervaded by Siva's radiance. 140 His intuitive faculty (mati) thus purified, the yogi gains both the perfections (siddhi) of yogic practice and liberation (mukti). His consciousness is now like a well-polished mirror which reflects everything he desires and grants it to him. 141 Abhinava writes: Just as a man who has been ill for a long time forgets his past pain completely when he regains his health, absorbed as he is in the ease of his present condition, so too those who are grounded in pure awareness free of thought-constructs are no longer conscious of their previous [fettered] state. Consciousness, the sole truly existent reality, free of thought-constructs is made fully and evidently manifest by eliminating these differentiated perceptions. The wise man should therefore exert himself to attend closely to this [state of awareness]. 142 The thought-constructs generated within consciousness do not in reality affect it at all. They can neither break up nor add anything to the Light which shines as all things. 143 They are in fact nothing but 192 The Doctrine of Vibration consciousness itself 144 which perceives, through its power of reflective awareness (yimarsa), the multitude of objects in diverse ways, and so assumes this form. 145 Although thought-constructs are mental representations of objects once seen or present, they are products of the power of consciousness and not of the objects they represent. 146 Thought is both analytic and synthetic; 147 it serves the useful purpose of separating individual elements of experience from others and linking together those that appear to be distinct from one another so that they can be better understood. 148 It does not consist merely of false mental constructs projected onto reality that need to be wholly rejected. Thought obscures consciousness and distracts it only when it appears in the form of doubt, vacillating between alternatives. 149 Once this conflicting duality (dvaitddhivasa) 150 is eliminated, thought is purified and rests in itself as the 'thought-less thought' of pure consciousness. 151 By gradually eliminating the multitude of conflicting notions that agitate him, the yogi ultimately achieves the certainty (niscaya) corresponding to a direct awareness of his own divine nature. 152 Abhinava explains: Thought is in reality none other than pure consciousness. Even so, it serves as a means to liberation for the individual soul (anu) only when it takes the form of certainty (niscaya). 153 The yogi must eliminate every doubt and misguided notion that leads him to believe himself to be other than Siva. By developing the thought: 'I am Siva', it ultimately affirms itself directly as a pure awareness beyond thought without any intervening mental representations. Abhinava says: Just as the man who thinks intensely that he is a sinner becomes such, just so one who thinks himself to be Siva, and none other than He, becomes Siva. This certainty (ddrtfhyā), which penetrates and affirms itself in our thoughts, coincides with an awareness free of thought-constructs engendered by a series of differentiated mental representations, the object of which is our identity with Siva. 154

As thought is gradually purified, it becomes progressively clearer until its object becomes maximally apparent (sphufatama). 155 The stream of perceptive consciousness (pramana) progressively reveals each aspect of its object which, thus affirming itself with increasing clarity, reveals its ultimate nature. The yogi reflects repeatedly upon it as the object of his realisation and loving devotion, for all that is perceptible and need be known (jñeyam) is Siva alone. As Abhinava says: Path to Liberation 193 What should we say of those who before they are satisfied have to see their beloved again and again, caress her and think about her for a long time? 156 The yogi practising the Empowered Means is initiated into the Great Sacrificial Rite (mahayaga), eternally enacted at the interface between the inner and outer aspects of consciousness, by a direct infusion of awareness from his master who is the embodiment and outer symbol of the yogi's enlightened identity. 157 The rite begins with ritual bathing (snana) which is in this case the immersion of the body of thought in the white ashes of the cosmic fuel of duality, burnt in the fire of consciousness. 158 The yogi then goes on to worship (puja) by uniting all that is pleasing to the senses in the oneness of consciousness. 159 The ritual formula (mantra) he recites is the eternal resonance of the awareness which is the pulsation of the Heart of his own consciousness. 160 Repetition (Japa) of the formula is every activity, perception, breath or thought which arises within him while plunged in the universal awareness of his true nature. 161 The mental image he visualises meditating (dhyana) on the Deity in the course of the rite, is whatever the yogi spontaneously imagines and contemplates as the outpouring of the universal creativity of consciousness. 162 Ritual gesture (mudra) is whatever bodily posture the yogi may assume when, fully absorbed in consciousness, he moves, staggering about (ghumita) as it were, drunk with the wine of self-realisation. 163 Oblation is performed by offering with devotion and awareness all the sensations which flow in through the channels of the senses to the fire of his subjectivity, which is thus inflamed (uddipita) and makes all things one with itself. 164 The outer ritual which commences in the sphere of the Individual Means thus leads naturally to the inner rite of the Empowered Means. When the yogi's practice (abhyasa) reaches fruition, the rite merges with the spontaneous activity of consciousness. This is fullness (purnata), the completion and reunification of the forces within consciousness which, through the power of ignorance, were formerly dispersed and divided. "Just as a horse driven here and there", writes Abhinava, "over plains, hills and dales follows the will of its rider, so also consciousness, driven by various expedients (bhāgi), quiescent or terrific, abandoning duality, becomes Bhairava. Just as by looking repeatedly at one's own face in a mirror one comes to know that it is the same [as the image reflected], so also, [one sees] in the mirror of mental representations of meditation {dhyana), ritual (puja) and worship (area) one's own Self as Bhairava and so quickly identifies with Him. This identification is the realisation that takes place in the absolute (anuttara)." 165 194 The Doctrine of Vibration By ridding himself of the relative distinctions engendered by thought, the yogi practising the Empowered Means, illumined by the power of self-awareness of Pure Knowledge {suddhavidya), transcends the distinction between right and wrong, purity and impurity. He is led to the conviction that the pure consciousness, which is his true nature, is unaffected by whatever action he may do, whether conventionally accepted as good or bad. Abhinava quotes the Malinivijaya as saying: All here is enjoined and all prohibited. This alone, O Lord of the gods, is here prescribed as obligatory, namely that the mind be firmly applied to the true reality. It matters little how this is achieved. He whose mind is firmly established in [this] reality, even if he eats poison, is as little affected by it as are lotus petals by water. 166 Impurity is a state of seeming separation from consciousness. 167 The yogi who has freed himself of all false notions comes to realise that the true nature of consciousness can never be sullied or limited by any object appearing within it. 168 This is the realisation the ancient sages achieved through a direct intuition of reality free of intruding thought-constructs {avikalpabhava), but kept secret in order not to confuse the worldly. 169 Similarly, in reality nobody is ever bound. It is ignorance to believe bondage exists and to contrast it with a conceived state of liberation. 170 If the Self is one with Siva, how can it be either bound or released? 171 Nothing essentially distinguishes those who are bound from those who are free. 172 The difference between their states is merely conceptual. 173 Pure consciousness abides free of all such distinctions. Thus Bhagavatopala, in his commentary on the Stanzas, repeatedly stresses that thought-constructs obscure consciousness and misguide the individual soul. 174 Those who are bound are convinced that they are dull witted, conditioned by Karma, sullied by their sin and helplessly impelled to action by some power beyond their control. He who manages to counter this conviction with its opposite achieves freedom. 175 He who considers himself to be free is free indeed, while he who thinks himself bound remains so. Thus at the highest level of realisation, as Abhinava says: Nothing new is achieved

nor is that which in reality is unmanifest, revealed- [only] the idea is eradicated that the luminous being shines not. 176 Nothing is impure, all is perfect, including Maya and the diversity it engenders. To say that illusion exists and that ignorance must be Path to Liberation 195 eradicated implies that it has a separate existence apart from conscious- ness. If this is so, it has as little reality as the shadow of a shadow, but if not, then it must be consciousness itself. Thus, as Kallata says, bondage, the binder and the bound are in fact one. 177 It is Siva Himself Who freely obscures His own nature. Siva binds Himself by Himself. 178 Concealing and revealing Himself, Siva plays His timeless game. At the Divine (sambhava) level of pure Siva-consciousness, the Spanda yogi directly lays hold of the power inherent in his own conscious nature (svabala) which gives life to the psycho-physical organism and impels the senses and mind to action. 179 In this way every thought- construct, and with it the ego, is instantly annulled in the immediacy of the pure subjectivity that remains unaltered throughout every perception and state of consciousness. The same takes place at the Empowered level b> attending to the recurrent activity — Spanda — of the subject, that is, the flux of awareness through the cyclic movement of the powers of consciousness. 180 By attending (avadhana) to this movement the thought- constructs that emerge and subside in the course of perception are seen to be part of this universal process, and, in this way purified, are no longer binding. Thus, Ksemaraja says that the Spanda teachings are concerned most directly with the Empowered Means. m The yogi who is always alert to discern the pulse of Spanda quickly realises his own authentic state of being (niharfi bhavam). 1 * 2 He is then truly awake, not only literally, but also in the deeper sense that he is awake to his authentic nature, its power and activity. When attention (avadhana) slackens, this movement takes place unconsciously and so the thought- constructs and perceptions generated through it appear to take on an autonomous existence of their own, just as happens when we dream. 183 The spontaneity of the movement that travels between subject and object and holds them together in the pure awareness of the universal subject's identity with his cosmic object devolves into the creative activity of waking and dreaming. Man, in other words, becomes a victim of his states of consciousness and the contents that they, by their very nature, generate within themselves. 184 The Spanda teachings are not only concerned with the structure of thought and its functions, but also with the powers and properties of its vehicle, namely, speech. Speech issues out of consciousness, develops into thought to then become articulated sound. A focal point of Spanda doctrine is thus the role speech plays in the formation of thought- constructs and their purification. Although this takes place at all levels of practise below the Divine (sambhava), the Spanda teachings, meant as they are for advanced yogis, ignore the outer forms of spiritual discipline to concentrate on practise in the Empowered (sakta) psychic sphere (cetas) and what lies beyond it where speech is the pure inner awareness (yimarsa) **Although Abhinava does not differentiate between individual consciousness and cosmic general ledger, one tends to comprehend that right interpretation is the transfor mentioned the individual consciousness to cosmic consciousness responsible for the creation and destruction which the sadhaka sees as his own power. Every one is rama and every one is Shiva. only thing that is essential is striving and concerted efforts and sustained struggle to reach.** The Doctrine of Vibration of the light of consciousness. The Doctrine of Vibration identifies this, the highest level of speech (para vac), with the universal pulse of consciousness that resounds spontaneously within it as the inner flow of its own undifferentiated awareness. 185 Beyond the realms of language, it is the transcendental consciousness in which all language is rooted and pervades all that language denotes as its essential being. Utpaladeva writes: The Supreme Voice is consciousness. It is self-awareness spontaneously arisen, the highest freedom and sovereignty of the Supreme Lord. That pulsing radiance (sphuratta) is pure Being, unqualified by time and space. As the essence [of all things] it is said to be the Heart of the Supreme Lord. 186 When the intention arises within consciousness to discern its own brilliance manifest in the world of denotations and denoted meanings, speech turns from the supreme transcendental level to that of immanence and assumes the form of a pure intuitive awareness (pratibha) which perceives and comprehends its universal manifestation. This is the voice of intuition (pasyanti), which grasps the meaning inherent inwardly in all words and externally in all that they denote. Analogous to the non- discursive, instinctual knowledge animals possess, it is a pure generic perception not yet formed into language in which the act of denotation, its object and that which denotes it are indistinguishable. Illumined by the voice of intuition birds migrate in their due seasons, the cock crows at dawn and young mammals suck at the breast. 187 Infants similarly reflect and respond instinctively to their environment by virtue of this intuitive sense 188 and through it come to grasp the link between words and the

objects they denote. As they learn to speak, they begin to form concepts and so the next two levels of speech develop. One is the outer corporeal speech (yaikhari) and the other the subtler, inner discourse (antah- saryijalpa) of thought that forms at the intermediate level (madhyama) where the ratiocinating mind stands between the higher levels of intuition and its outer verbal expression. In this way the development of speech in infancy reflects its progressive actualisation in every spoken word. A hymn to the Goddess quoted by Bhagavatopala describes this process well: Therefore, O Supreme Goddess, the highest form of speech should be worshipped as the [universal] cause that establishes the existence of all things by insight [niscaya] into their nature (artha) brought about by their manifestation through the superimposition [of verbal designations]. O Mother, insight into the true nature [of things] is nothing but the Path to Liberation 197 act of intent of that [same speech], apart from which [speech itself and all that it expresses] could not attain to its own nature. Again, in that state [speech] is said to be the light of one's own nature. Free of division and succession it is attainable [only] by the yogi. Then from the state of intent, O Siva, speech [assumes] the nature of thought as the radiant pulse (sphuraria) of desire to speak of that which is in the domain of words. Then consisting of words, it bears a clearly expressed meaning, for if [speech] were not such, meaning could not be understood. 189 personal experience clearly proves that thought is invariably associated with speech. 190 Thought is a function of language. Through it we communicate to ourselves a mental image of the world about us and can construct complex ideas about ourselves. Language is the fabric from which our world of ideas is woven. Mental representation which orders the influx of sensation and presents us with a meaningful, picture of the world, memory, the elaboration of ideas and the shifting tides of emotion are all intimately connected with language and through it to the consciousness which underlies them. To think of language as nothing more than a system of denotation based on a commonly accepted convention (sanketd) fails to fully account for the inherent power to convey meaning (vacakasakti). In order to learn the convention we must be born with an innate ability to grasp meaning, and this ability is not itself learned nor found anywhere within the domain of convention. Lacking this ability we would be caught in an infinitely expanding system of denotation in which each element pointed to some other within it, without ever coming to rest anywhere. Unless we can couple the word 'jar' with the object it denotes, explaining that the word 'pot' is a synonym of the word 'jar' would leave us none the wiser. 191 The connection between word and meaning is only explicable if we postulate that it is an inherent property of the power of awareness to link one with the other. Language must be grounded in the pure cognitive awareness (prama) of consciousness which stands beyond, and yet illumines, the sphere of experience we define and understand through the medium of language. As Abhinava says: Someone may hear another person speak, but if his awareness (prama) is obscured, he is unable to rise, unconscious as he is, to the level of the experiencing subject [who understands] what has been said. He only grasps the outer successive (sound) of what the other person says and thus can only repeat it as would a parrot. An understanding of its meaning presupposes that he has caught hold of his own power of awareness (prama) by attaining the autonomy [of the conscious, universal subject]. **Note that we have defined sachitananada as one which understand Truth and Untruth, Bliss and Unbliss, and Knowledge and ignorance where all these are equal. Truth=Untruth, Bliss=Unbliss and Ignorance=Knowledge** The Doctrine of Vibration Outer, articulate speech consists of a series of ordered phonemic elements produced and combined by the vocal organs to form meaningful words. In order for this to be possible, these elements must also be grounded in consciousness (prama). The articulated phonemes are merely outer, gross manifestations of the phonemic energies (yarnagrama) held in a potential state within consciousness. This 'mass of sounds' (sabdarsi) is the light of consciousness (prakdsd) which makes the universe manifest and contains all things within itself. In other words, it is the totality of consciousness expressed as the collective awareness symbolised by all the letters corresponding to the introverted subjectivity of Siva Himself. The power through which this potential actualises itself into speech and the world of denotation is technically called * Mdrkd*. It is the reflective awareness (vimarsa) and radiance (sphuratta) of the supreme subject — the 'mass of sounds' (Jabdarasi) — and the undivided wonder Siva experiences when He contemplates the universe He gathers up into Himself in the form of countless words (vdcaka) and their meanings (vdcya). 193 Mdrkdsakti is manifest in the second movement of consciousness after the primal vibration of the pure luminosity of the 'mass of sounds', as the state of pure potency which arises when its unsullied subjectivity begins to turn away from itself and is associated with faint traces of objectivity (dmrsya- cchdyd). 194 Mdrkd contains within itself the various aspects of objectivity that, although not yet manifest, are ready to issue forth. Thus this power, at

one with Siva, is called 'Mdrkd' because she is the mother (mdtrkd) of the universe that she contains within herself as does a pregnant woman her child. 195 The circle of the powers of Mdrkd (mdtrkdcakra) consists of the phonemic energies contained in AH AM, the universal ego. 196 When grasped in its entirety at its source, these energies elevate the consciousness of the enlightened, but when split up and dispersed give rise to the obscuring forces (kald) which lead the ignorant away from realisation. The fettered soul is ignorant of the pure egoity that is the source of speech and so it generates, through its powers, the many thought-constructs that deprive him of the awareness of unity and obscure Siva's universal activity. 197 The Stanzas on Vibration declare: He who is deprived of his power by the forces of obscurity {kala} and a victim of the powers arising from the mass of sounds (sabdarasi) is called the fettered soul. 198 The powers [of speech] are always ready to obscure his true nature as no mental representation can arise that is not penetrated by speech. 199 The rays of phonemic energies emanate from the light of Siva, the Path to Liberation 199 'mass of sounds' (sabdarasi) in eight groups. They constitute the powers of the inner mental organ and the five senses, figuratively arranged in a circle around the sacred shrine {pi(ha) of Matrka&akti who manifests externally as the body. 200 The eight classes and the names of the deities presiding over them are as follows: 201 Gutturals Brahman! Intellect (buddhi) Palatals Mahegvari Ego (ahankara) Cerebrals Kaumarl Mind (manas) Dentals Narayani Hearing Labials Varahi Touch Semivowels Aindri Sight Sibilants Camuntfa Taste Vowels Mahalak\$ml Smell The yogi who grasps the true nature of the power of Matfka and its phonemic forces is liberated 202 by recognising that the activity of the senses and the discursive representations of the mind are in fact emanations of universal consciousness. Conversely, when ignorant, he is affected by its power in its multiple negative aspects known as 'Mahdghora' ('greatly terrible') and, unable to rest within himself free of the sense of diversity, he is constantly disturbed by the flux of extroverted perceptions. Abhinava explains: When the [phonemic energies] are not known to be [emanations of the Lord] they conceal the wonder (camatkara) of consciousness which is the one essential non-discursive awareness [present throughout perception] and even in discursive thought. They obscure it with thought-constructs constituted by the diverse configurations of phonemes and syllables which [although also] a form of the deity [are no longer benevolent but] most terrible. Inducing doubt and fear, they engender the fettered soul's state, bound by the shackles of transmigrati on. But once their true nature is understood correctly in this way, they bestow freedom in this very life This knowledge of their intimate being [at one with the absolute] consists of this, namely, that even in the midst of all these fluctuations, free at their inception of discursive representations, thought-constructs do not conjoin [individualised consciousness] with the wheel of energies consisting of the totality of phonemes, even though [these constructs] are coloured by the many diverse words generated by the aggregate of phonemes. Language has a powerful effect on us. A few words we may hear or read can inspire us with joy, fear or sadness, and the constant inner 200 The Doctrine of Vibration dialogue of thought arouses intense feelings within us. This power hidden in language, which binds us through the thought-constructs it generates, can also be used to free us of them by channeling it through Mantra. Mantric practice begins at the Individual (driava) level where Mantras are recited in consonance with the rising and falling away of the breath. In this way they are charged with the vibration (spandd) of consciousness and, in their turn, make consciousness vibration (Doctrine of Vibration S.G.Dyczkowski) On without knowledge that everything is manifest within consciousness is illusory or unreal in that sense alone. 78 Things are more real or more tangibly experienced according to their own essential nature (svabhava) to the degree in which we recognise that they are appearances (abhasa) within absolute consciousness As Jayaratha says: Just as images manifest in a mirror, for example, are essentially mere appearances, so too are [phenomena] manifest within conscious- ness. Thus, because they are external, [phenomena] have no being (sattva) of their own. The Lord says this [not with the intention of saying anything about the nature of things] but in order to raise the level of consciousness of those people who are attached to outer things; thus everything in this sense is essentially a mere appearance. [Knowing this], in order to quell the delusion of duality, one should not be attached to anything external. 79 The ultimate experience is the realisation that everything is contained within consciousness. We can discover this in two ways. Either we merge the external world into the inner subject, or we look upon the outer as a gross form of the inner. In these two ways we come to recognise that all things reside within our own consciousness just as consciousness resides within them. This all-embracing inwardness is only possible if there is an essential identity between the universe and consciousness. The events which constitute the universe are always internal events happening within consciousness because their

essential nature is consciousness itself. 80 We can only account for the fact that things appear if there is an essential identity between consciousness and the object perceived. If a physical object were really totally material, that is, part of a reality independent of, and external to, consciousness, it could never be experienced. Abhinava says: The existence or non-existence of phenomena within the domain of the empirical (iha) cannot be established unless they rest within consciousness. In fact, phenomena which rest within consciousness are apparent (prakasamana). And the fact of their appearing is itself their oneness {abheda) with consciousness because consciousness is nothing but the fact of appearing {prakasa). If one were to say that they were separate from the light of [that consciousness] and that they appeared [it would be tantamount to saying that] 'blue' is separate from its own nature. However, [insofar as it appears and is known as such] one says: 'this is blue 1 . Thus, in this sense, [phenomena] rest in conscious- ness; they are not separate from consciousness. The Doctrine of Vibration The universe and consciousness are two aspects of the whole, just as quality and substance constitute two aspects of a single entity. The universe is an attribute (dharma) of consciousness which bears (dharmiri) it as its substance. It is said that 'substance' is that resting in which this entire group of categories manifests and is made effective. Now, if you don't get angry [we insist that] this entire class of worlds, entities, elements and categories (tattva) rests in consciousness and [resting in it] is as it is. Thus consciousness contains everything in the sense that it is the ground or basis (adhard) of all things, their very being (satta) and substance from which they are made. But, unlike the Brahman of the Advaita Vedanta, it is not the real basis (adhithand) of an unreal projection or illusion. Consciousness and its contents are essentially identical and equally real. They are two forms of the same reality. Consciousness is both the substratum and what it supports: The perceiving awareness and its object. In this respect, the Kashmiri Saiva is frankly and without reserve an idealist. Although he does not deny the reality of the object, his position is at odds with most commonly accepted forms of realism. The realist maintains that the content perceived is independent of the act of perception. The content is only accidentally an object of perception and undergoes no change in the process of being perceived. His contention, however, is essentially unverifiable; to verify it, we would have to know an object without perceiving it. This, from the Kashmiri Saiva point of view, is not possible. Objects of which we have no knowledge may indeed exist, but they are knowable as objects only if they are related to subjects who perceive them. In this sense, if there were no subjects, there could be no objects. 86 The subject, however, as opposed to the object is, in terms of the phenomenology of perception, apparent to himself. He is self-luminous (svaprakdsa). Thus, conscious- ness (the essence of subjectivity) is one s own awareness by virtue of which all things exist. The realist maintains that consciousness clearly differs from its object insofar as their properties are contrary to each other. The Saivite idealist. However, says that the object is a form of awareness (vijnanakara)TM The objective status of the object is cognition itself. Perception manifests its object and renders it immediately apparent (sphuta) to those who perceive it. It does not appear at any other time. If 'blue' were to exist apart from the cognition of k blue\ two things would appear: 'blue' and its cognition, which is not the case. It is the perception of the object which constitutes its manifest nature. An entity becomes an Integral Monism of Kashmiri Saivism object of knowledge not by virtue of the entity itself but by our knowledge of it. If objects had the property of making other objects appear, it would be possible for one object to make another appear in its own likeness. 'Blue' is perceived to be 'blue' because it is manifest as such to the perceiver. As Abhinava points out: The [nature of an] object of knowledge could not be established through a means of knowledge totally unrelated to it — a crow does not become white because a swan [sitting next to it] is white. Perception, on the other hand, is immediately apparent to conscious- ness. It is self-luminous in the sense that it is directly known without need of being known by any ulterior acts of perception and makes its object known at the same time. Adopting the Buddhist Yogacara doctrine that things necessarily perceived together are the same (sahopa- lambhaniyamavada), the Saivite affirms that because the perceived is never found apart from perception, they are in fact identical. Reality (satya) is the point where the intelligible and the sensible meet in the common unity of being; it cannot be said to exist in itself outside, and apart from, knowledge or vision. **Bhagavatotpala in his commentary on the Stanzas on Vibration quotes: Once the object is reduced to its authentic nature, one knows [the true nature of] consciousness. What then [remains of] objectivity? What [indeed could be] higher than consciousness? Consciousness is essentially active.** Full of the vibration of its own energy engaged in the act of perception, it manifests itself externally as its own object. When the act of perception is over, consciousness reabsorbs the object and turns in on itself to resume its undifferentiated inner nature. Knowledge (jnana) manifests

internally and externally as each individual entity.... Once knowledge has assumed that form it falls back [into itself]. The Yogacara Buddhist similarly maintains that consciousness creates its own forms. But, according to him, because the perceived and perception are identical; there is no perceived object at all. The so-called outer world is merely a flux of cognitions, it is not real. He is firmly committed to a doctrine of illusion. The reality of consciousness from The Doctrine of Vibration his point of view is established by proving the unreality of the universe. "All this consists of the act of consciousness alone", says Vasubandhu, "because unreal entities appear, just as a man with defective vision sees unreal hair or a moon, etc." He points to dreams as examples of purely subjective constructs which appear to be objective realities. The apparent reality dreams possess is not derived from any concrete, objective world, but merely from the idea of objectivity. While the Yogacara does not say that an idea has, for example, spatial attributes, it does have a form manifesting them. While he agrees with the Saiva idealist that appearances have no independent existence apart from their appearing to consciousness, he maintains that for this reason they are unreal. The creativity of consciousness consists in its diversification in many modes having apparent externality; it is not a creation of objects. While the Kashmiri Saivite agrees that the world is pure consciousness alone, he maintains that it is such because it is a real creation of consciousness. The effect is essentially identical with the cause and shares in its reality. Matter and the entire universe are absolutely real, as 'congealed' (sty ana) or 'contracted' (samkucita) forms of consciousness. "This God of consciousness", writes Ksemaraja, "generates the universe and its form is a condensation of His own essence (rasa)" m By boiling sugarcane juice it condenses to form treacle, brown sugar and candy which retains its sweetness. Similarly, **consciousness abides unchanged even though it assumes the concrete material form of the five gross elements.** The same reality thus abides equally in gross and subtle forms. Consequently no object is totally insentient. Even stones bear a trace (vasana) of consciousness, although it is not clearly apparent because it is not associated with the vital breath (prana) and other components of a psycho-physical organism. Somananda goes so far as to affirm that physical objects, far from being insentient, can only exist insofar as they are aware of themselves as existing. The jar performs its function because it knows itself to be its agent. Indeed, all things are pervaded by consciousness and at one with it and hence share in its omniscience. Thus, Siva, Who perceives Himself in the form of physical objects, is the one ultimate reality. "The jar knows because it is of my nature", writes Somananda, "and I know it because I am of the jar's nature. I know because I am of Sadasiva's nature and He knows because He is of my nature; Yajnadatta [knows] because he is of Siva's nature and Siva [knows] because He is of Yajnadatta's nature". Integral Monism of Kashmiri Saivism Everything in this sense is directly perceived by absolute consciousness, and this direct perception (pratyaksa) unifies the knowable into a single, undivided whole. This is the central concept behind a doctrine originally expounded by Narasimha called 'the non-dualism of direct perception' (pratyaksadvaita). This states that consciousness is essentially perceptive and that its perception of all things operates throughout the universe. Insofar as phenomena are clearly evident (sphufa) to us, everything is directly perceived by absolute consciousness, with which our individual consciousness is identical. This direct perception unfolds everywhere; the one true reality, it is alone and without companion or rival (nihsapatna). Even though it remains one, it can, by its very nature, perceive distinctions (bheda) between one entity and another, without this engendering any division within it. We distinguish between two entities in empirical terms on the basis of their mutual exclusion (anyonyabhava). The relative distinction {bheda) between them is essentially the perceived difference between their respective characteristics. Despite this difference they are united within the purview of a single cognition insofar as they are equally both manifest appearances. This cognition is the undivided essence (rasa) or 'own nature' (svabhava) of both. Encompassed by the 'fire of consciousness', there is no essential difference between them. Just as when an emerald and ruby reflect each other's light, the ruby is reddish-green and the emerald greenish-red, similarly everything is connected with everything else as part of the single variegated (vicitra) cognition of absolute consciousness. Mahesvarananda writes: The Supreme Lord's unique state of emotivity (asadharanabhava) is the outpouring of pure Being (mahasatta). It is manifest as the brilliance (sphuratta) of the universe which, if we ponder deeply, [is realized to be] the single flavour (ekarasa) of the essence of Beauty which is the vibration of the bliss of one's own nature. In this way all things are in reality one although divided from the one another sharing as they do the 'single flavour' (ekarasa) of the pure vibration of consciousness. Kashmiri Saiva Realism Kashmiri Saivism as a whole has been variously called a form of 'realistic idealism', 'monistic idealism', 'idealistic monism' and 'concrete monism'. It is

easy to understand why Kashmiri Saivism is The Doctrine of **Vibration said to be 'idealistic' and 'monistic', but in what sense is it also 'realistic'?** The answer to this question is of no small importance in trying to understand the central idea behind its metaphysics and the fundamental importance of the concept of Spanda, in this seemingly impossible marriage between monistic idealism and pluralistic realism. The Kashmiri Saiva approach understands the world to be a symbol of the absolute, that is, as the manner in which it presents itself to us. Again we can contrast this view with that of the Advaita Vedanta. The Advaita Vedanta understands the world to be an expression of the absolute insofar as it exists by virtue of the absolute's Being. Being is understood to be the real unity which underlies empirically manifest separateness and as such is never empirically manifest. It is only transcendently actual as 'being-in-itself. The Kashmiri Saiva position represents, in a sense, a reversal of this point of view. The nature of the absolute, and also that of Being, is conceived as an eternal becoming (satatodita), a dynamic flux or Spanda, 'the agency of the act of being'. It is identified with the concrete actuality of the fact of appearing, not passive unmanifest Being. Appearance (abhāsa) alone is real. appearing (prakasamanatva) is equivalent to the fact of being (astitva). Ksemaraja writes in his commentary on the Stanzas on Vibration: Indeed, all things are manifest because they are nothing but manifestation. The point being that nothing is manifest apart from manifestation. The absolutely unmanifest, from this point of view, can have as little existence as the space in a lattice window of a sky-palace. Nay, even less, because even **that space can appear as an imagined image manifest within consciousness. Everything is real according to the manner in which it appears. Even an illusion is in this sense real, insofar as it appears and is known in the manner in which it appears. The empirical and the real are identical categories of thought. As Abhinava says: Thus this is the supreme doctrine (upaniṣat), namely that, whenever and in whatever form [an entity] appears, that then is its particular nature. Perhaps at this stage a brief comparison with Heidegger's ideas might prove to be enlightening and not altogether out of place. According to Heidegger's phenomenology of Being, reality is intelligible in a two-fold Integral Monism of Kashmiri Saivism manner** as 'phenomenon' and 'logos'. **Heidegger defines** what he means by 'phenomenon' as: "that-which-shows-itself. The manifest . . . phenomena are then the collection of that which lies open in broad daylight or can be brought to the light of day — what the Greeks at times implicitly identified as 'ta onta' (the things-which-are)". 127 In his later writings Heidegger drops the term 'phenomenon' in preference for the verbal form 'phainesthai' in order to emphasize even more the actuality or presentational property of Being. Explaining this new form of the term he writes: "Being disclosed itself to the ancient Greeks as 'physis'. The etymological roots 'phy-' and 'pha-' designate the same thing: 'phyein', the rising-up or upsurge which resides within itself as 'phainesthai', lighting-up, self-showing, coming-out, appearing-forth." Heidegger contrasted his notion of phenomenon with semblance (Schein) and with appearing (Erscheinung). In the case of semblance a thing can show itself as that which it is not, as when fool's gold shows itself to be gold. The ancients always allied semblance with non-being. Heidegger points out, however, that semblances are grounded in showings, and so does Abhinava. Both Heidegger and Abhinava consequently maintain that all semblances have a real basis and are to be treated as instances of phenomena along with the so-called real showing or manifestation of non-deceptive objects. So Heidegger states that: 'how- ever much seeming, just that much being'. Thus self-showing or appearing defines Being as phenomenon, but this definition of Being is as yet incomplete. Being is not only self-showing but 'logos' which Heidegger explains means 'discourse' (Rede) in the sense of 'apophansis': 'letting-be-seen'. Phenomenology, which according to Heidegger is the only correct study of Being, means 'letting-be-seen-that-which- shows-itself. This is true of Saiva Paramadvaita as well. The reality of the world demands recognition; we are forced to accept the direct presentation of the fact of our daily experience. As Abhinava says: "if practical life, which is useful to all persons at all times, places and conditions were not real, then there would be nothing left which could be said to be real" **A thousand proofs could not make 'blue' other than the colour blue. The reality of whatever appears in consciousness cannot be denied. Objects appear; they do not cease to do so by a mere emphatic denial.** The manifestation of an entity in its own specific form is a fact at one level of consciousness; it is real. The appearing of the same entity in the same form but recognised to be a direct representation of the absolute is also a fact, but at another level of consciousness. It is no more or less real than the first. 'As is the state of consciousness, so is the experience,' says Abhinava. Although the nature of the absolute is discovered at a higher level of consciousness, The Doctrine of Vibration nonetheless it presents itself to us directly in the specific form in which we perceive things; otherwise there would be no way in which we could penetrate from the level of appearing to that of

its source and basis. Abhinava writes: Real is the entity (yastu) that appears in the moment of direct perception (sak\$atkara), that is to say, within our experience of it. Once its own specific form has been clearly determined one should, with effort, induce it to penetrate into its pure conscious nature. All things are known to be just as they present themselves. The concrete actuality of being known (pramiti), irrespective of content, is itself the vibrant (spanda) actuality of the absolute. Liberating knowledge is gained not by going beyond appearances but by attending closely to them. "The secret," Mahesvarananda says, "is that liberation while alive (jivanmukti) is the profound contemplation of Maya's nature." No ontological distinction can be drawn between the absolute and its manifestations because both are an appearing (dbhasa), the latter of diversity and the former of 'the true light of consciousness which is beyond Maya and is the category Siva'. Those who have attained the category of **Pure Knowledge above Maya and have thus gone beyond the category of Maya, see the entire universe as the light of consciousness . . . Just as the markings [on a feather] are nothing apart from the feather, the feather [is nothing apart from] them, similarly, when the light of consciousness is manifest, the whole group of phenomena is manifest as the light of consciousness itself.** Within the sphere of Maya, every entity's 'own nature' (svabhdva) corresponds to its specific manifest form. Accordingly it is defined as that which distinguishes it from all else and from which it never deviates. Above the sphere of Maya, that is, above the level of objectivity, is the domain of the subject. At this level, everything is realised to be part of the fullness of the experience and hence no longer bound by the conditions which impinge on the object. Here the part is discovered to be the whole, that is, consciousness in toto. In this sphere beyond relative distinctions, the yogi realises that (all) the categories of existence are present in every single category. The yogi experiences every individual particular as the sum total of everything else. He recognises that all things have one nature and that every particular is all things. This is the 'essence' (sard) or co-extensive unity (samarasya) of all things. Integral Monism of Kashmiri Saivism We have established that reality is manifest according to how [and the degree in which] the freedom of consciousness reveals it and that [this freedom] is the womb of all forms. Just as 'sweetness' is present in its entirety in every atom of the sugarcane, so each and every atom [of the universe] bears within itself the emanation of all things. This is the level of consciousness in which the absolute reflects on itself realising to its eternal delight and astonishment (camatkara) its own integral nature. The reality of the world of diversity is not denied, but experienced in a new mode of awareness free of time and space in the eternal omnipresence of the Here and Now. [Phenomenal forms of awareness] such as 'this [exists]', born of the colouring [imparted to the absolute] by the limitations engendered by the diversifying power of time {kalakalana} also emanate within the Supreme Principle. There [at that level], Fullness {purriata} is the one nature [of all things] and so everything is omnipresent; otherwise, associated with division (khancjlana), the Fullness [of the absolute] would not be full. The content of absolute consciousness consists of diverse appearances (abhd\$sa) which, because they are manifest through it in this way, do not compromise the wholeness of consciousness. Everything we perceive is a momentary collocation of a number of such manifestations which combine together like a row of altar lamps' (dipavalT) to form the single radiant picture of the universe. The individual objects which constitute the universe are specific collocations of such 'atomic' appearances. Together they form a single unified particular which appears according to its own defining features (svalaksand). A jar, for example, consists of a number of appearances such as 'round', 'earthen', 'red', etc., which together discharge a single function (arthakriya), in this case, that of carrying the appearance 'water'. They unite with each other much as the scattered rays of a lamp come together when focused, or as the various currents of the sea together give rise to waves. Atomic appearances can combine in any number of ways, provided that they are not contrary to one another as established by the dictates of natural law (niyati). An appearance of 'form', for example, cannot combine with that of 'air'. Insofar as they share a common basis (samanyadhikaranya), a given cluster of appearances appears as a single whole. This common basis is the most prominent member of the group; the appearance 'jar' is such in the example quoted above. Any one appearance in a cluster may assume a more important or subordinate role. The result is a specific The Doctrine of Vibration awareness of an object of the form: 'here this is such.' While individual appearances do not lose their separate identity {svarupabhedha} when they rest on a common basis, even so the particular object which appears according to its own characteristics (svalak\$ana) is an individual reality in its own right. It is a different kind of appearance characterised by its association with the appearance of the specific location and time in which it is made manifest. The form of our experience is thus 'I now see this here'. But when we perceive each particular constituent

appearance separately, each assumes a separate fixed function. Abhinava cites the following colourful example to illustrate how the various combinations of appearances account for the variety of experience: Thus even though the appearance of the beloved may manifest externally, it is as if far away in the absence of another appearance, namely, that of 'embracing'. So when the [appearing of the beloved] is associated with another appearance [namely that of 'far away'] the power (arthakriya) it formerly had of giving pleasure appears as its contrary. The form our experience assumes depends, not only on the nature of the object perceived, but also on personal factors entirely peculiar to ourselves. This theory explains this in two ways. In one sense, the object remains the same, but one or other of its constituent appearances comes to the fore according to the inclinations of the perceiver. From another point of view, we can say that the perceived object is different for each perceiver according to the difference in the prominent appearance manifest to him. Abhinava, citing as an example a golden jar, illustrates how the same object appears differently to different perceivers according to the use they wish to make of it and to their state of mind: When a person who is depressed and feels that there is nothing [of value for him in the world] sees the jar, he merely perceives the appearance 'exists' [in the form of the awareness that] 'it is'. He is not conscious of any other [of its constituent appearances] at all. An individual who desires to fetch water [perceives] the appearance 'jar'. The man who simply wants something that can be taken somewhere and then brought back [perceives] the appearance 'thing'. The man who desires money [perceives] the appearance 'gold'. The man who desires a pleasing object [perceives] the appearance 'brightness' while he who wants something solid sees the appearance 'hardness'. These 'atomic events' or appearances emerge from the pure subject's consciousness and combine together to form a total event at each moment. Integral Monism of Kashmiri Saivism Daily life (vyavahara) goes on by virtue of this ever renewed flux of appearances. They are connected together and work towards a single unified experience because they appear within the field of consciousness of the universal subject. The aggregate of appearances arises in the [supreme] subject as do [sprouts in] a rice field. Even though each sprout germinates from its own seed, they are perceived as a collective whole. Appearances rest in this way within the universal subject. 'External-ity' is itself another appearance; it arises from a distinction between appearances and the individual subject. So, although all manifestation always occurs within the subject, it appears to be external due to the power of Maya which separates the individual subject from his object. This split must occur for daily life to be possible. Only externally manifest appearances can perform their functions; when they are merged within the subject and at one with him, they cannot do so. Daily life proceeds on the basis of the operation and withdrawal of the conditions necessary for fruitful action to be possible. Appearance in this sense represents the actualisation of a potential hidden in consciousness made possible by virtue of its dynamic, Spanda nature which is both the flow from inner to outer and back as well as the power that impels it. The emergence from, and submergence into, pure consciousness of each individual appearance is a particular pulsation (visesaspana) of differentiated awareness. Together these individual pulsations constitute the universal pulse (samanyaspana) of cosmic creation and destruction. Thus, every single thing in this way forms a part of the radiant vibration {sphuratta, sphurana) of the light of absolute consciousness. Light and Awareness: The Two Aspects of Consciousness Absolute consciousness understood as the unchanging ontological ground of all appearing is termed 'Prakasa'. As the creative awareness of its own Being, the absolute is called 'Vimarsa'. **Prakasa and Vimarsa — the Divine Light of consciousness and the reflective awareness this Light has of its own nature — together constitute the all-embracing fullness (purnata) of consciousness. The Recognition (pratyabhijñā) school of Kashmiri Saivism develops this concept of the absolute which finds its fullest expression in Utpaladeva's Stanzas on the Recognition of God.** Even though neither of these two key terms appear in the Stanzas on Vibration or the Aphorisms of Siva, they recur frequently in their commentaries. Thus, although the original formulation of the Doctrine of Vibration differs from the theology of Recognition in this respect, it was extended in the course of its development to accommodate this concept of the absolute as well. This was possible, and quite justified, insofar as the absolute understood in Pratyabhijñā terms does not, as we shall see, differ essentially from that of the Spanda school. We can, as Kashmiri Saivites themselves have done, explain one in terms of the other. The Doctrine of Vibration Prakasa: The Light of Consciousness Prakasa is the pure luminosity' (bhdna) or 'self-showing' that constitutes the essence and ultimate identity (atman) of phenomena. That things appear at all is due to the light of consciousness, and their appearing (avabhasana) is itself this Light which bestows on all things their evident, manifest nature. Established in the light of consciousness everything appears there according to its own specific nature (svabhava).

Anything that supposedly does not rest in this Light is as unreal as a sky-flower. 3 Thus, according to Rajanaka Rama, unlike the light of the sun, or any other light, this Light not only makes all things apparent, it is also their ultimate source. 4 Full of its divine vibration the Light makes all things manifest and withdraws them into itself. This supra- temporal activity characterises it most specifically; devoid of it, it would be no better than an inert physical phenomenon. At the same time, this light is the conjunction (slesa) or oneness (aikdmya) of its countless manifest forms, 6 and the collective whole (sarpina'ana) of all the categories of existence. The universe is nothing but the shining of the Light within itself. It is the radiant vibration (sphuratta) of this Light, the state (avast hand) in which consciousness becomes manifest. Although the Light shines as all things at all times and hence also makes their diversity manifest, 9 penetrating each object individually as well as collectively, it is not totally 'merged' (magna) or identified with the object so as to suffer any division within itself. Our experience of any object is of the form: 'I see this': it is not itself an object, but the manifest form the object assumes as a luminous principle of experience. The Light is ever revealed and can never be obscured; objectivity can never cast a shadow on the light of consciousness. The Stanzas on Vibration declare: That in which all this creation is established and from whence it arises is nowhere obstructed because it is unconditioned by [its very] nature. This Light is the highest reality (paramartha). It is the 'Ancient Light' (puranaprakasa) that makes all things new and fresh every moment. It is 'always new and secret, ancient and known to all'. It is the form of the Present (vartamanarupa), the Eternal Now. Time and space are relations between the contents of consciousness; they cannot impinge on the integrity of the absolute itself. 16 Neither space nor time can divide it, for they are one with the Light that illumines them Light and Awareness: Two Aspects of Consciousness and makes them known as elements of experience. But this Light is the shining of the absolute; it is not an impersonal principle. It is the living Light of God, indeed it is God Himself, the Master Who instructs the entire universe. 18 Siva is this 'auspicious lamp', Who illumines all things. He is the Light of consciousness that reveals the presence of both the real and the unreal, of light' and 'darkness'. 20 Abhinavagupta writes: Thus Bhairava, the Light, is self-evident (svatahsiddha); without beginning, He is the first and last of all things, the Eternal Present. And so what else can be said of Him? The unfolding of the categories of existence (tattva) and creation, which are the expansion of His own Self, He illumines, luminous with His own Light, in identity with Himself, and because He illumines Himself, so too He reflects on His own nature, without His wonder (camatkara) being in any way diminished. Although It Is Possible To Catch Glimpses Of The Highest Reality In Advanced States Of Contemplation Before Attaining Perfect Enlightenment, These States, However Long They Last, Are Transitory (Kadacitka) And When They End The Vision Of The Absolute Ceases With Them. The Highest Realisation, However, Persists In All States Of Consciousness. It Happens Once And Need Never Occur Again. A Passage From A Lost Tantra Declares: "The Self Shines Forth But Once, It Is Full [Of All Things] And Can Nowhere Be Unmanifest." All Spiritual Discipline Culminates In This Moment Of Realisation. Accordingly, Abhinava Stresses That The Goal Of All The Means To Realisation, Even The Individual Means, Is This Absolute Consciousness. Finally, It Is Worth Noting That Although Abhinava Affirms That The Teachings Concerning Anupaya Are Found In The Siddhayogesvarimata And The Malinxvijaya, Both Of Which, According To Abhinava, Are Major Tantras Of The Trika School, It Is In The Theology Of The School Of Recognition That It Is Best Exemplified. Abhinava Himself Refers To Somananda, The Founder Of This School, As Teaching It And Alludes To The Following Passage In The Vision Of Siva To Support His Own Exposition: When Siva, Who Is Everywhere Present, Is Known Just Once Through The Firm Insight Born Of Right Knowledge (Pramana), The Scripture And The Master's Words, No Means [To Realisation] Serves Any Purpose And Even Contemplation {Bhavana) [Is Of No Further Use]. Anupaya Is Therefore, According To Abhinavagupta, The Recognition Of One's Own Authentic Siva-Nature, Which All The Higher Tantric Traditions Teach Is The Ultimate Realisation. This Is Also True Of The Doctrine Of Vibration Whose Precedents Are Clearly Traceable To These Same Traditions. Thus, Although The Stanzas Themselves Never Refer Directly To Enlighten- Ment As An Experience Of Recognition, There Can Be Little Doubt That Spanda Practice Leads To This Same Realisation. Accordingly, Commentators Stress That We Realise The Vibration Of Consciousness By Recognising Its Activity And That Liberation Depends On The Recognition Of This As One's Own Nature. Ksemaraja Describes What Happens In This Moment Of Recognition According To The Doctrine Of Vibration Thus: At The End Of Countless Rebirths, The Yogi's [Psycho-Physical] Activity [Which Issues From Ignorance] Is Suddenly Interrupted By The Recognition Of His Own Transcendent

Nature, Full Of A Novel And Supreme Bliss. He Is Like One Struck With Awe. And In This Attitude Of Astonishment (Vismaya- Mudra) Achieves The Great Expansion [Of Consciousness] {Mahavikasa}. Thus He, The Best Of Yogis, Whose True Nature Has Been Revealed [To Him] Is Well Established [At The Highest Level Of Consciousness], Which He Grasps Firmly And His Hold Upon It Never Slackens. Thus He Is No Longer Subject To Profane Existence (Pravrtti), The Abhorrent And Continuing Round Of Birth And Death, Which Inspires Fear In All Living Beings, Because Its Cause, His Own Impurity, No Longer Exists. The Divine Means (Sambhavopaya) In Anupaya The Yogi Does Not Need To Deal With The World Of Diversity At All; Only Paramasiva Exists There. Beyond Both Immanence And Transcendence, He Has Nothing To Do With The World Of Practice And Realisation. Anupaya Is The Experience Of The Undefinable (Qnakhya) Light Of Consciousness, Which Is The Pure Bliss Beyond Even The Supreme State (Parattha) Of Sivatatva. At A Slightly Lower Level, Corresponding To The Divine Means, A Subtle Distinction Emerges Between The Goal And The Path. The Yogi Now Practises Within The Domain Of The Outpouring Of The Power Of Consciousness. From This Level He Penetrates Directly Into The Universal Egoity Of Pure Consciousness By The Subtle Exertion (Udyama) Of Its Freedom (Svatantrya) And Reflective Awareness. The Yogi Who Practises The Divine Means Is Not Concerned With Any Partial Aspect Of Reality But Centers His Attention Directly On Its Abounding Plenitude. Hence This Means Is Based On Siva's Own State (Sambhavavastha) In Which Only The Power Of Freedom Operates As The Pure Being (Satta) Or Essence Of All The Other Powers. This State Is The Light Of Consciousness Which, Free Of All Thought- Forms, Is The Basis Of All Practice." The Yogi Who Recognises That Pure Consciousness, Free Of Thought-Constructs (Nirvikalpa), Is His Basic State, Can Practice In Any Way He Chooses; Even The Most Common Mantra Will Lead Him Directly To The Highest State. Thus The Forms Of Contemplative Absorption, Empowered (Sakta) And Individual (Anava), Which Are The Fruits Of The Other Means To Realisation Both, Attain Maturity In This Same Undifferentiated Awareness. This Awareness Is The Pure Ego Manifest At The Initial Moment Of Perception (Prathamikalocana), When The Power Of The Will To Perceive Is Activated. It Is The Subtle State Of Consciousness That Reveals The Presence And Nature Of Its Object Directly: That Which Shines And Is Directly Grasped In The First Moment Of Perception While It Is Still Free Of Differentiated Representations And Reflects Upon Itself Is [The Basis Of The Divine Means] Said To Be The Will. Just As An Object Appears Directly To One Whose Eyes Are Open Without The Intervention Of Any Mental Cogitation (Anusaidhana), So, For Some, Does Siva's Nature. The Movement Of Awareness At This Level Of Practice Attains Its Goal Quickly. While Consciousness Is Heightened Progressively In The Other Means, Here It Expands Freely To The Higher Levels, Unconfined By Any Intruding Thought-Constructs. The Divine Means Is A 'Thoughtless Thought', A 'Processless Process', That Occurs At The Juncture Between Being And Becoming. Abhinava Explains: When The Heart [Of Consciousness] Is Pure And [Free Of Thought- Constructs], It Harbours The Light Which Illumines The Radiant, Primordial Plane (pragbhumi) Together With All The Categories Of Existence. [The Yogi] Then Realises Through It His Identity With Siva Who Is Pure Consciousness The Yogi Must Catch The Initial Moment Of Awareness (Adiparamarsa) Just When Perception Begins^ He Must Not Move On From The First Pure Sensation Of The Object But Return To Its Original Source In His Own T Consciousness. Observing In This Way The Objective Field Of Consciousness Without Laboring To Distinguish Particulars, The Yogi Penetrates Into His Own Subjectivity Which, Vacuous And Divested Of All Outer Supports (Niralamba), Is Not Directed Anywhere Outside Itself (Ananyamukha- Preksin). Here He Can Hold Of The Power Inherent In His Own Consciousness Through Which He Discerns The True Nature Of Whatever Appears Before Him. Thus The Stanzas On Vibration Teach: Just As An Object, Which Is Not Seen Clearly At First Even When The Mind Attends To It Carefully, Becomes Later Fully Evident When Observed With The Effort Exerted Through One's Own [Inherent] Strength (Svabala), In The Same Way, When [The Yogi] Lays Hold Of That Same Power, Then Whatever [He Perceives Manifests To Him] Quickly According To Its True Nature, Whatever Be Its Form, Locus, Time Or State. Thus, Although The Practice Of This Divine Means Starts By Catching Hold Of The Will In The First Moment Of Awareness, It Also Concerns The Second And Third Moments In Which The Means Of Knowledge And The Object Are Made Manifest. When Practice At This Level Proceeds Smoothly And Without Interruption, The Three Powers Of Will, Knowledge And Action Fuse Into The Trident (Trisula) Of Power, Which Is The Subject Free Of All Obscuration (Niranjana), M At One With The Power Of Action In Its Most Powerful And Evident Form. The Kaula Schools Call This State The Stainless (Niranjanatattva).

Equated In The Spanda Tradition With The Dawning Of The Vibration Of Consciousness {Spandodaya), It Is The Enlightenment The Spanda Yogi Seeks. Spanda Practice Is Based On The Experience Of Spanda Which, As We Have Seen, Is Defined As The Intent (Aunmukhya) Of Consciousness, Unrestricted To Any Specific Object And Hence Free Of Thought-Constructs. Spanda Can Therefore Be Experienced Directly When A Powerful Intention Develops Within Consciousness, Whatever Is Its Ultimate Goal Or Cause. We Have Already Noted That Intense Anger, Joy, Grief Or Confusion Is Such Occasions. Similarly, The Yogi Can Make Contact With The Omnipotent Will, Which He As Siva Possesses, Through Intense Prayer. Directing His Entire Attention To Siva, The Benefactor Of The World, Entreating Him Fervently And Without Break, His Will Merges With Siva's Universal Will, Which Is The Source Of Every Impulse And Perception. As He Looks About Him, The Yogi Realises That It Is Siva Himself, The Universal Consciousness And The Yogi's Authentic Identity, Who Ordains His Every Action, Thought And Perception. Thus The Yogi's Cognitive Intent On His Object Coincides With The Universal Will To Make That Object Known To Him, Whether The Yogi Is Awake Or Dreaming. He Is Thus No Longer Like The Worldly Man Who Cannot Dream As He Wishes And Is Forced To Experience Whatever Spontaneously Happens In These States Of Consciousness. Ultimately The Yogi Manages, By Siva's Grace, To Maintain A Constant Awareness Of His Own Pure Perceptive Consciousness (Upalabdhrta) Divested Of All Obscuring Thought-Constructs In Deep Sleep As Well As In The Contemplative State (Turiya) Beyond It. When He Rises To The Higher Levels Of Contemplation In Which The Breath Is Suspended And All Sensory And Mental Activity Ceases, The Yogi Who Manages To Sustain This Pure, Undifferentiated Awareness Does Not Succumb To Sleep As Do Less Developed Yogis. Perfection In The Practice Of The Divine Means Thus Coincides With The Goal Of Spanda Practice, Namely, A Constant, Alert Attention To The Perceiving Subjectivity Which Persists Unchanged In Every State Of Consciousness Both As The Perceiver And Agent Of All That It Experiences. Another Important Spanda Practice Belonging To This Means Is Centering. The Spanda Yogi Seeks To Find The Centre (Madhya) Between One Cognition And The Next; For It Is There That He Discovers The Expansion (Unmesd) Of Consciousness Free Of Thought-Constructs From Whence All Differentiated Perceptions (Yikalpa) Emerge. 107 Abhinava Explains That This Pure Awareness Is Called: . . . The Expansion (Unme\$A) Of [Consciousness] Or The Creative Intuition {Pratibha) [Experienced] In The Interval Which Divides Two [Moments] Of Differentiated Perception (Vikalpa). It Is Here That They Arise And Disappear. The Sastras And Agamas Proclaim With Reasoned Argument That It Is Free Of Thought-Constructs {Nirvikalpa) And Precedes All Mental Representations Of Any Object. None Can Deny That A Gap Exists Between Perceptions Insofar As Two Moments Of Thought Are Invariably Divided. This [Gap] Is The Undifferentiated Unity Of All The Countless Manifestations. Similarly, In The Outer More Objective Sphere, Where Change Consists Of The Alterations In The Configurations Of Manifest Appearances (Abhasa) The Transition From One To Another Corresponds To A Phase Of Pure Luminosity That Marks The Beginning Of One Form And The End Of Another. The World Of Manifestation And Differentiated Perceptions (Yikalpa) Thus Extends From One Centre To The Next. Although It Is Never In Fact Divorced From The Subject Who Resides There, The Ignorant Fail To Grasp This Fact And So, Cut Off From The Centre, The World Of Objectivity Becomes For Them The Sphere Of Maya. Bhagavatotpala Quotes The Light Of Consciousness (Samvitprakdsa): This Ever Pure Experience (Suddhanubhava) Is Variegated By Each Form [Revealed Within It]; Even So It Remains Unstained (Nirmala) When Moving To Another. Just As A Cloth Which Is Naturally White, Once Dyed, Cannot Change Colour Without [First] Becoming White Again, Similarly The Pure Power Of Awareness, (Citi) Once Coloured By Form, Is Pure [Again] At The Centre Where That Form Is Abandoned And From Whence It Proceeds To Another. In His Essence Of Vibration (Spandasarpdoha), K\$Emaraja Explains That The Rise And Fall Of Every Individual Perception In The Field Of Awareness Is A Specific Pulsation Of Consciousness. From The Point Of View Of The Object, The Expansion (Unmesa) Of This Pulse Is Represented By The Initial Desire To Perceive (Didrksa) A Particular Object, While The Contracted (Nimesa) Phase Is The Withdrawal Of Attention From The Object Previously Perceived. From The Point Of View Of The Perceiving Subjectivity, The Phases Are Reversed, So That The Initial Desire To Perceive Marks The Contraction (Nimesa) Of Subjective Consciousness While The Falling Away Of The Previous Perception Is Its Expansion (Unmesa). At The Higher Level, Where These Two Phases Are Experienced Within Consciousness, They Represent The State Of The Categories Of Isvara ('This Universe Is Me') And Sadasiva ('I Am This Universe'). Utpaladeva Says: Expansion (Unme\$A) Which Is In The

External Manifestation [Of Objectivity], Is Kvaratattva While Contraction (Nime\$A), Which Is In The Internal Manifestation [Of Subjectivity], Is Sadasiva. At This Level All The Powers Of Consciousness Fuse And Both Phases Are Manifest As Part Of One Reality. This Unity Is In Fact Apparent To Everybody At Each Moment. However, Within The Domain Of Maya, Which Is The Sphere Of Differentiated Perceptions (Yikalpa), It Is Clearly Manifest Only At The Juncture (Madhya) Between Two Cognitions. In This Centre Resides The Void (Kha) Of Consciousness (Free Of Thought-Constructs) Which, Divested Of Diversity, Digests Into Itself All The Psycho-Physical Processes That Give Life To The Multiplicity Of Perceptions. The Yogi Moves From The Particular Vibrations Of Consciousness At Its Periphery To The Universal Throb Of The Heart In The Centre. As Abhinava Explains: The Self-Reflective Awareness In The Heart Of Pure Consciousness, Present At The Beginning And End Of Each Perception, Within Which The Entire Universe Is Dissolved Away Without Residue, Is Called In The Scriptures, The Universal Vibration Of Consciousness (Samanyaspana) And Is The Outpouring (Uccalana) [Of Awareness] Within One's Own Nature. All The Categories Of Existence (Tattvas) Are United In The Heart Of The Centre Where The Life-Giving Elixir Of Siva's Consciousness Floods One's Own Inner Nature. To Reside In The Centre Is To Abide By The Law Of Totality (Gramadharma) In A State Which Transcends The Workings Of The Mind (Unmana). Consciousness (Jnana) With Light As Its Support, Residing In The Centre Between Being And Non-Being Is Known As The Act Of Abiding In One's Own Abode As The Perceiving Subjectivity (Drahtftva) Free Of All Obscuration. That Which Has Been Purified By Pure Awareness (Suddhavijnana) Is Called The Transcendent (Viviktavastu), Said To Be The Mode Of Being (V/7//) Of The Law Of Totality (Gramadharma) Through Which Everything Is Easily Attainable. The Power In The Centre (Madhyasakti) Is The Eternal Present. Beyond Time It Is The Source Of Both Past And Future. To Be Established There Is To Abide Without A Break In Rama, The Supreme Enjoyer, In Every Action Of One's Life. Rama Is Siva, The Supreme Cause Who Pervades The Fourteen Aspects Which Embrace The Entire Universe Of Experience, Namely, Moving, Standing, Dreaming, Waking, The Opening And Closing Of The Eyes, Running, Jumping, Exertion, Knowledge [Born] Of The Power Of The Senses, The [Three] Aspects Of The Mind, Living Beings, Names And All Kinds Of Actions. By Developing An Awareness Of The Centre, The Yogi Experiences The Bliss Of Consciousness. Through This Gap He Plunges Into Introverted Absorption (Nimilanasamadhi) And Then Emerges Again To Pervade The Field Of Awareness Between Centres And So Experience The Cosmic Bliss (Jagadananda) Of The Universal Vibration Of Consciousness. 1 18 He Then Recognises That This State Pervades Every Aspect Of Experience. In This Way The Yogi's Consciousness Is No Longer Afflicted By The Power Which Obscures It, Hemming The Centre In On Both Sides With Thought-Constructs That Seemingly Deprive It Of Its Fullness. As He Realises Directly His Pure Conscious Nature As The Universal Ego Free Of All Mental Representations, It Expands Out To Embrace All Things Within Itself. Thus The Realisation The Divine Means Leads To, And Is Directly Based Upon, Is That This Pure Ego Is In All Things Just As All Things Are Within It. In The Spanda Tradition, As Recorded In The Stanzas On Vibration, No Such Ego Is Recognised. ' 19 Man's Authentic Nature Is, However, Understood In Personal Terms As Every Individual's Own 'Own Nature* (Svasvabhava) Which Is Siva, The Universal Vibration Of Pure Subjectivity (Upalabdhfta). It Is Not Surprising, Therefore, That Later Commentators Found These Two Conceptions To Be Essentially The Same And Accordingly Identified One's Own Inner Nature With The Pure Ego. This Came As A Natural Development In Spanda Doctrine Not Only For This Reason But Also Because The Universal Ego Is Experienced As The Inner Dynamics Of Absolute Consciousness. To Conclude Our Summarial Exposition Of The Divine Means, Which Is Centred On The Direct Experience Of This Pure Ego (And Hence On Spanda In This Form), We Turn Now To A Brief Description Of Its Inner, Cyclic Activity. We Shall Do This By Examining Abhinava' S Esoteric Exegesis Of The Symbolic Significance Of The Word 'A HAM', Which In Sanskrit Means T, And Symbolises By Its Form The Ego's Dynamic Nature. The Objective World Of Perceptions Is, As We Have Seen, Essentially A Chain Of Thought-Constructs (Prapanca) Closely Linked To One Another And Woven Into The Fabric Of Diversity (Vicitrata). **This Thought (Vikalpa) Is A Form Of Speech (Vac) Uttered Internally By The Mind (Citta), Which Is Itself An Outpouring Of Consciousness. Consciousness Also, In Its Turn, Resounds With The Silent, Supreme Form Of Speech {Para Vac) Which Is The Reflective Awareness Through Which It Expresses Itself To Itself. Consequently, The** Fifty Letters Of The Sanskrit Alphabet, Which Are The Smallest Phonemic Units Into Which Speech Can Be Analysed, Are Symbolic Of The Principal Elements Of The Activity Of

Consciousness. Letters Come Together To Generate Words And Words Go On To Form Sentences. In The Same Way The Fifty Phases In The Cycle Of Consciousness Represent, In The Realms Of Denoted Meaning (Vacya), The Sum Total Of Its Universal Activity (Kriya) Corresponding To The Principal Forces (Kala) Which Come Together To Form The Metaphysical Categories Of Experience, Which In Their Turn Appear In The Grossest, Most Explicitly 'Articulate' Form As The One Hundred And Eighteen World-Systems (Bhuvana). 'A', The First Letter Of Both AHAM And The Sanskrit Alphabet, Is The Point Of Departure Or Initial Emergence Of All The Other Letters And Hence Denotes Anuttara — The Absolute. 'Ha', Is The Final Letter Of The Alphabet And Represents The Point Of Completion When All The Letters Have Emerged. It Represents The State In Which All The Elements Of Experience, In The Domains Of Both Inner Consciousness And Outer Unconsciousness, Are Fully Displayed. It Is Also The Generative, Emission (Visarga) Which, Like The Breath, Casts The Inner Into The Outer, And Draws What Is Outside Inward. The Two Letters 'A' And 'Ha' Thus Represent Siva, The Transcendental Source And Sakti, His Cosmic Outpouring That Flows Back Into Him. The Combined 'A-Ha' Contains Within Itself All The Letters Of The Alphabet — Every Phase Of Consciousness, Both Transcendental And Universal. (For A Graphic Representation Of This Analysis, See Figure 1.) M, The Final Letter Of AHAM, Is Written As A Dot Placed Above The Letter Which Precedes It. It Comes At The End Of The Vowel Series And Before The Consonants And So Is Called 'Anus Vara' (Lit. 'That Which Follows The Vowels') And Also 'Bindu' (Lit. 'Dot,' 'Drop,' 'Point' Or 'Zero'). While The Consonant 'M' Symbolises The Individual Soul (Purusa), 'Bindu' Represents The Subtle Vibration Of T, Which Is The Life Force (Jivakala) And Essence Of The Soul's Subjectivity Manifest At The Transcendental, Supra-Mental Level (Unmana). 12 ° It Is The Zero-Point In The Centre Between The Series Of Negative Numbers, In This Case The Vowels Which Represent The Processes Happening Internally Within Siva, And The Series Of Positive Numbers — The Consonants Which Symbolise The Processes Happening Externally Within Sakti. Bindu, As A Point Without Area, Symbolises The Non-Finite Nature Of The Pure Awareness (Pramitibhava) Of AHAM. It Is The Pivot Around Which The Cycle Of Energies From 'A' To 'Ha' Rotates, The Void In The Centre From Which All The Powers Emanate And Into Which They Collapse. As Such, It Is The Supreme Power Of Action Which Holds Subject, Object And Means Of Knowledge Together In A Potential State In The One Light That Shines As All Three Containing Them In Its Repose (Visrnti). Bindu Is The 'Knower' (Jndtr), Who Is Essentially Consciousness That, Though Omniscient, Does Not Manifest Its Intelligence, Like A Man Who Knows The Scriptures But Having No Occasion To Explain Them To Others Silently Bears This Knowledge Within Himself. As Such, It Symbolises The Union Of Siva And Sakti (Sivasakti- Mithunapina'a) In A State Of Heightened Potency In Which They Have Not Yet Divided To Generate The World Of Diversity. It Stands, In Other Words, At The Threshold Of Differentiation In The Stream Of Emanation Still Contained Within Siva. The Absolute. Expansion Commences Bindu — The Individual Soul Withdrawal Commences Then, To The Degree In Which That Which Is To Be Accomplished By The Power Of Action Residing Within It [As A Potential] Penetrates Into The Absolute, It Appears Initially **As Bindu, Which Is The Light Of Pure Consciousness. When Outer Objectivity Is Reabsorbed Into Its Transcendent Source, Bindu Is The Point Into Which All The Manifest Powers Of Consciousness Are Gathered And Fused Together. The Universal Potency Of All The Letters Is Thus Contained In Bindu Which, As The Reflective Awareness Of Supreme T Consciousness, Gives Them All Life. Thus Bindu Also Marks The Beginning Of Siva's Internal Movement Back To The Undifferentiated Absolute And So Stands At The Threshold Of Both Emission And Absorption Without Being Involved In Either.** The Three Aspects Of AH AM Together Constitute A Movement From The Undifferentiated Source Of Transcendental Consciousness — 'A' — Through The Expansion Or Emission Of Its Power — 'Ha' — To The Subject — 'M' — Which Contains And Makes Manifest The Entire Universe Of Experience. The Reverse Of This Movement, That Of Withdrawal (Samhara), Is Represented By M-Ha-A. AHAM And M-Ha-A Alternate In The Rotation (Ghurnana) Of The Reflective Awareness Of T Consciousness As Immanent Sakti Emerges From Transcendental Siva To Then Merge Back Into Him. As Abhinava Says: The Universe Rests Within Sakti And She On The Plane Of The Absolute (Anutiara) And This Again Within Sakti ... For The Universe Shines Within Consciousness And [Consciousness Shines] There [Within The Universe By The Power Of] Consciousness. These Three Poles, Forming A Couple And Merging, Make Up The One Supreme Nature Of Bhairava Whose Essence Is AHAM- At The Microcosmic Level, 'A' Represents The Initial Moment When The Subject Begins To Rise Out Of Himself To View The Object. The Movement From 'A' To

'Ha' Marks The Emergence Of Sensation Within The Field Of Awareness, Which Is Represented By The Fifty Letters Of The Alphabet Symbolic Of The Fifty Aspects Of The Flux Of Consciousness Leading To Objectified Perception. 'NI' Is The Subject Who, Resting Content Within Himself When He Has Perceived His Object Merges Through The Inner Flow Of Awareness Into K A\ The Absolute. Then From The Absolute (A) Its Emission (Ha) Flows Back Into The Pure Subject (M) Set To Perceive His Object. Thus All The Cycles Of Creation And Destruction Are Contained Within AHAM Through Which They Are Experienced Simultaneously As The Spontaneous Play Of The Absolute. The Yogi Who Recognises This Recurrent Pulse Of Awareness To Be The Movement Of His Own Consciousness Merges His Limited Ego With The Universal Ego. Thus He Realises That Its Power To Create, Sustain And Destroy All Things Is His Own Inner Strength (Svabaia) That He Exerts Effortlessly In The Same State Of Mystical Absorption (Turlyya) In Universal Consciousness That The Absolute Itself Enjoys. In This Way He Shares In The Three-Fold Awareness Siva Himself Has Of His Own Nature Which Abhinava Describes As Follows: I Make The Universe Manifest Within Myself In The Sky Of Consciousness. I, Who Am The Universe, Am Its Creator! ' — This Awareness Is The Way In Which One Becomes Bhairava. 'AH Of Manifest Creation(Sadadhvari) Is Reflected Within Me, I Cause It To Persist I — This Awareness Is The Way In Which One Becomes The Universe. The Universe Dissolves Within Me. I Who Am The Flame Of The [One] Great And Eternal Fire Of Consciousness' — Seeing Thus One Achieves Peace. The Experience Of The Liberated Thus Coincides With The Realisation Of Their Own Divine Nature Which, Through Its Power, Rules And Guides The Cosmic Order. Thus This Attainment (Siddhi), Which Is Liberation Itself, Is In The Doctrle Of Vibration Technically Called 'Mastery Over The Wheel Of Energies' (Cakresvaratvasiddhi) Because The Liberated Soul, Identified With Siva, Now Governs, As Does Siva, The Cycle Of The Powers That Bring About The Creation And Destruction Of All Things, The Empowered Means (Saktopaya) All The Practices Taught In The Stanzas On Vibration Are Internal. Whenever Ritual Is Mentioned, It Is Invariably Interpreted In Terms Of The Dynamics Of The Inner Processes The Yogi Experiences And Implements In The Course Of His Yogic Practice. The Doctrine Of Vibration, Ksemaraja Affirms, Is Concerned Entirely With These Inner Disciplines Centred, As It Is, In One Way Or Another, On Consciousness Or, At Least, On The Inner Activity Of The Mind. Thus The Empowered Means Which, Like The Other Categories We Have Discussed, Is Entirely Internal Includes An Important Part Of Spanda Practice. Spanda Practice Belonging To The Divine Means Centres On One's Own Inherent Nature (Svasvabhava) As Siva, The Universal Perceiver And Agent, That Belonging To The Empowered Means On His Power Instead Of Arriving Directly At The All-Embracing Emptiness Of Subjective Conscious- Ness, The Yogi Practising The Empowered Means Realises His True Nature Through The Fullness Of Its Energy. Practising The Divine Means, The Yogi Plunges, As It Were, Straight Into The Fire Of Consciousness; Practising The Empowered Means He Merges With Its Rays. Either Way The Yogi Is Centred Equally On Ultimate Reality. The Power Of Consciousness Is No Less Absolute Than Its Possessor. To Make This Point Abhinava Quotes The Matanga- Tantra: This Reality Consists Of The Rays Of [Siva's] Power And Is Variously Said To Be The Abode Of The Lord's Manifestation . . . That Same [Power] Illumined [By Siva] Is Itself Also Luminous, Unshaken And Unmoving. That Very [Power] Is The Supreme State, Subtle, Omnipresent, The Nectar Of Immortality, Free Of Obscuration, Peaceful, Yearning For Pure Being Alone {Vastumatra) And Devoid Of Beginning And End. Perfectly Pure, It Is Said To Be The Body [Of Ultimate Reality]. The Yogi Concentrates On The Powers Operating In All Of Life's Activities As Particular Pulsations (Visesaspana) In The Universal Rhythm (Samanyaspana) Of The Power Of Consciousness. In This Way He Rises Progressively From The Particular To The Universal Until He Reaches Pure Being (Satta), The Greatest Of All Universals (Mahasamanyd) And The Highest Form Of Siva's Power. Thus The Creative Power Of Maya, Manifest Through Countless Lesser Powers, No Longer Causes The Yogi To Stray From Siva's Consciousness But Becomes The Means Through Which It Can Be Realised In The Illuminating Brilliance (Sphuratta) Which Is Siva's Pure Being. Thus By Discovering The True Nature Of Sakti, The Yogi Realises Himself To Be Siva, Its Possessor Who Consists Of All Its Countless Powers. Thus Practise Belonging To This Means Leads To The Same Pure Consciousness Free Of Thought-Constructs Realised Through The Divine Means. Although The Ultimate Realisation Is Instantaneous, The Yogi Rises To It Gradually By Freeing His Consciousness Of The Limitations Imposed Upon It By Thought. Abhinava Explains: The Same Occurs In The Empowered Means [As Does In The Divine]. At The Discursive Level Of Consciousness {Yaikalpikibhumi) [Where The Empowered

Means Functions] Knowledge And Action, Although Evident, Are, For The Reasons Explained Previously, Contracted. A Blazing Energy [Is Revealed Within] The One Who Dedicates Himself To Removing The Burden Of This Contraction. [This Energy Eventually] Brings About The Inner Manifestation {Antarabhasa) Of Pure Consciousness He Seeks. Consciousness Is Individualized And Its Power Of Knowledge And Action Contracted By **The Thought-Constructs Born Of Ignorance**. The Arising Of These Mental Representations, As The Stanzas On Vibration Say, Deprives The Soul Of Its Freedom And Immortal Life. The Practise Of The Empowered Means Is Meant To Free The Fettered Soul Of This Constriction On His Consciousness. It Operates Within The Mental Sphere (Cetasy™ And Is Designed To Purify Thought (Vikalpasamskara) In Order To. Reveal The Pure Consciousness Which Is Its Ground And Ultimate Source. Thus, The Empowered Means Is Concerned With The Second Instant Of Perception, During Which The Subject Forms Mental Representations Of His Object. Thought Functions On The Basis Of An Awareness Of Relative Distinctions Between Specific Particulars, Distinguishing Them From One Another And Thus Seemingly Fragmenting The Essential Unity Of Reality. The Vibrant Vitality Of Consciousness, Universally Manifest, Is Clouded Like A Mirror By A Child's Breath And The Soul Is Deprived Of The Liberating Intuition Of The One Reality Free Of Thought-Constructs (Nirvikalpa). Abhinava Writes: The [Fettered Soul] Is Like A Dancing Girl Who Although Wishing To Leave The Dancehall Is Collared By The Doorkeeper Of Thought And Thrown Back Onto The Stage Of Maya All Thought Is Centred On Objectivity And Hence Dislodges Awareness From The Plenitude Of Pure Subjective Consciousness. Thus, To Regain The Original State Of Rest (Visranti) Consciousness Enjoys, The Yogi Must Rid Himself Of Thought. As Thought-Forms Decrease, Pure, Thought-Free Awareness Is Strengthened Until The Yogi Is Fully Established In A State In Which The Relative Distinctions (Bheda) Conceived Between Entities Dissolve Away. Everything Appears To Him As Pure Being (Sattdmdtra) L} And The Entire Universe Shines Before Him Pervaded By Siva's Radiance. His Intuitive Faculty (Mati) Thus Purified, The Yogi Gains Both The Perfections (Siddhi) Of Yogic Practice And Liberation (Mukti). His Consciousness Is Now Like A Well-Polished Mirror Which Reflects Everything He Desires And Grants It To Him. Abhinava Writes: Just As A Man Who Has Been Ill For A Long Time Forgets His Past Pain Completely When He Regains His Health, Absorbed As He Is In The Ease Of His Present Condition, So Too Those Who Are Grounded In Pure Awareness Free Of Thought-Constructs Are No Longer Conscious Of Their Previous [Fettered] State. Consciousness, The Sole Truly Existent Reality, Free Of Thought- Constructs Is Made Fully And Evidently Manifest By Eliminating These Differentiated Perceptions. The Wise Man Should Therefore Exert Himself To Attend Closely To This [State Of Awareness]. The Thought-Constructs Generated Within Consciousness Do Not In Reality Affect It At All. They Can Neither Break Up Nor Add Anything To The Light Which Shines As All Things. They Are In Fact Nothing But Consciousness Itself Which Perceives, Through Its Power Of Reflective Awareness (Yimarsa), The Multitude Of Objects In Diverse Ways, And So Assumes This Form. Although Thought-Constructs Are Mental Representations Of Objects Once Seen Or Present, They Are Products Of The Power Of Consciousness And Not Of The Objects They Represent. Thought Is Both Analytic And Synthetic; It Serves The Useful Purpose Of Separating Individual Elements Of Experience From Others And Linking Together Those That Appear To Be Distinct From One Another So That They Can Be Better Understood. It Does Not Consist Merely Of False Mental Constructs Projected Onto Reality That Need To Be Wholly Rejected. Thought Obscures Consciousness And Distracts It Only When It Appears In The Form Of Doubt, Vacillating Between Alternatives. Once This Conflicting Duality (Dvaitddhivasa) Is Eliminated, Thought Is Purified And Rests In Itself As The 'Thought-Less Thought' Of Pure Consciousness. By Gradually Eliminating The Multitude Of Conflicting Notions That Agitate Him, The Yogi Ultimately Achieves The Certainty (Niscaya) Corresponding To A Direct Awareness Of His Own Divine Nature. Abhinava Explains: Thought Is In Reality None Other Than Pure Consciousness. Even So, It Serves As A Means To Liberation For The Individual Soul (Anu) Only When It Takes The Form Of Certainty (Niscaya). **This surprising result--that information capacity depends on surface area--has (e) a natural explanation if the holographic principle (proposed in 1993 by Nobelist Gerard't Hooft of the University of Utrecht in the Netherlands and elaborated by Susskind) is true. In the everyday world, a hologram is a special kind of photograph that generates a full three-dimensional image when it is illuminated in the right manner**. All the information describing the 3-D scene is encoded into the pattern of light and dark areas on the two-dimensional piece of film, ready to be regenerated. The holographic principle contends that an analogue of this visual magic

applies to the full physical description of any system occupying a 3-D region. It proposes that another physical theory defined only on the 2-D boundary of the region completely describes the 3-D physics. If a 3-D system can be fully described by a physical theory operating solely on its 2-D boundary, one would expect the information content of the system not to exceed that of the description on the boundary. Can we apply the holographic principle to the universe at large? The real universe is a 4-D system. It has volume and extends in time. If the physics of our universe is holographic, there would be an alternative set of physical laws, operating on a 3-D boundary of spacetime somewhere, which would be equivalent to our known 4-D physics. We do not yet know of any such 3-D theory that works in that way. Indeed, what surface should we use as the boundary of the universe? One step toward realizing these ideas is to study models that are simpler than our real universe. A class of concrete examples of the holographic principle at work involves so-called anti-de Sitter spacetimes. The original de Sitter spacetime is a model universe first obtained by Dutch astronomer Willem de Sitter in 1917 as a solution of Einstein's equations, including the repulsive force known as the cosmological constant. De Sitter's spacetime is empty, expands at an accelerating rate and is very highly symmetrical. In 1997 astronomers studying distant supernova explosions concluded that our universe now expands in an accelerated fashion and will probably become increasingly like a de Sitter spacetime in the future. Now, if the repulsion in Einstein's equations is changed to attraction, de Sitter's solution turns into the anti-de Sitter spacetime, which has equally as much symmetry. More important for the holographic concept, it possesses a boundary, which is located "at infinity" and is a lot like our everyday spacetime. Using anti-de Sitter spacetime, theorists have devised a concrete example of the holographic principle at work: a universe described by superstring theory functioning in an anti-de Sitter spacetime is completely equivalent to a quantum field theory operating on the boundary of that spacetime. Thus, the full majesty of superstring theory in an anti-de Sitter universe is painted on the boundary of the universe. Juan Maldacena, then at Harvard University, first conjectured such a relation in 1997 for the 5-D anti-de Sitter case, and it was later confirmed for many situations by Edward Witten of the Institute for Advanced Study in Princeton, N.J., and Steven S. Gubser, Igor R. Klebanov and Alexander M. Polyakov of Princeton University. Examples of this holographic correspondence are now known for spacetimes with a variety of dimensions. This result means that two ostensibly very different theories--not even acting in spaces of the same dimension--are equivalent. Creatures living in one of these universes would be incapable of determining if they inhabited a 5-D universe described by string theory or a 4-D one described by a quantum field theory of point particles. (Of course, the structures of their brains might give them an overwhelming "commonsense" prejudice in favor of one description or another.)

Holographic Space-Time (Crystal Links, Tom Hanks and Wikipedia) Francisco Di Biase propose a quantum-informational holographic model of brain-consciousness-universe interactions based in the holonomic neural networks of Karl Pribram, in the holographic quantum theory developed by David Bohm, and in the non-locality property of the quantum field described by Hiroshi Umezawa. He considers this model an extension of the interactive dualism of Sir John Eccles, of an interconnection between brain and spirit by means of quantum microsites named dendrons and psychons. Francisco Di Biase proposes a dynamic concept of consciousness seen as a holoinformational flux interconnecting the holonomic informational quantum brain dynamics, with the quantum informational holographic nature of the universe. This self-organizing flux is generated by the holographic mode of treatment of neuronal information and can be optimized through practices of deep meditation, prayer, and others states of higher consciousness that underlie the coherence of cerebral waves. In brain mapping studies performed during the occurrence of these harmonic states we can see the spectral array of brain waves highly synchronized and perfectly ordered like a unique harmonic wave, as if all frequencies of all neurons from all cerebral centers played the same symphony. This highly coherent brain state generates the non-local holographic informational cortical field of consciousness that interconnect the human brain and the holographic cosmos. The comprehension of this holonomic quantum informational nature of brain-consciousness-universe interconnectedness allows us to solve the old mind-matter Cartesian hard problem, unifying science, philosophy, and spiritual traditions in a more transdisciplinary, holistic, integrated paradigm. In this new arrangement cosmovision, consciousness and transpersonal phenomena becomes part of Science and of the very holoinformational nature of the Holographic Conscious Multiverse. DOI: 10.14704/nq.2009.7.4.259

Quantum-Holographic Informational Consciousness Francisco Di Biase NeuroQuantology Holography suggests a considerable reduction of degrees of freedom in theories with gravity.

However it seems to be difficult to understand how holography could be realized in a closed re-contracting universe. In this Letter we claim that a scenario which achieves that goal will eliminate all spatial degrees of freedom. This would require a different concept of quantum mechanics and would imply an intriguing increase of power for the natural laws. **Physics Letters B Volume 451, Issues 1–2, 1 April 1999, Pages 19–26 Richard Dawid doi:10.1016/S0370-2693(99)00205-1** **Vijay Balasubramanian, Per Kraus, Albion Lawrence, and Sandip P. Trivedi** describe probes of anti-de Sitter spacetimes in terms of conformal field theories on the AdS boundary. Basic tool is a formula that relates bulk and boundary states—classical bulk field configurations are dual to expectation values of operators on the boundary. At the quantum level they relate the operator expansions of bulk and boundary fields. Using our methods, we discuss the CFT description of local bulk probes including normalizable wave packets, fundamental and D-strings, and D-instantons. Radial motions of probes in the bulk spacetime are related to motions in scale on the boundary, demonstrating a scale-radius duality. They also discuss the implications of these results for the holographic description of black hole horizons in the boundary field theory. DOI: <http://dx.doi.org/10.1103/PhysRevD.59.104021> **Holographic probes of anti-de Sitter spacetimes Phys. Rev. D 59, 104021 – Published 26 April 1999 Vijay Balasubramanian, Per Kraus, Albion Lawrence, and Sandip P. Trivedi** The existence of a fundamental scale, a lower bound to any output of a position measurement, seems to be a model-independent feature of quantum gravity. In fact, different approaches to this theory lead to this result. The key ingredients for the appearance of this minimum length are quantum mechanics, special relativity and general relativity. As a consequence, classical notions such as causality or distance between events cannot be expected to be applicable at this scale. They must be replaced by some other, yet unknown, structure. **LUIS J. GARAY, Int. J. Mod. Phys A, 10, 145 (1995) DOI: 10.1142/S0217751X95000085 QUANTUM GRAVITY AND MINIMUM LENGTH** A digital computer is generally believed to be an efficient universal computing device; that is, it is believed able to simulate any physical computing device with an increase in computation time by at most a polynomial factor. This may not be true when quantum mechanics is taken into consideration. This paper considers factoring integers and finding discrete logarithms, two problems which are generally thought to be hard on a classical computer and which have been used as the basis of several proposed cryptosystems. Efficient randomized algorithms are given for these two problems on a hypothetical quantum computer. These algorithms take a number of steps polynomial in the input size, e.g., the number of digits of the integer to be factored. Copyright © 1997 Society for Industrial and Applied Mathematics **SIAM J. Comput., 26(5), 1484–1509. (26 pages) Polynomial-Time Algorithms for Prime Factorization and Discrete Logarithms on a Quantum Computer ISSN (print): 0097-5397 ISSN (online): 1095-7111 Publisher: Society for Industrial and Applied Mathematics Peter W. Shor DOI: 10.1137/S0097539795293172** The classical dynamics of M-dimensional extended objects arising from stationary points of the world volume swept out in space time is discussed from various points of view. A introduction to the Hamiltonian mechanics of bosonic compact M (em) branes is given, emphasizing the diversity of the different formulations and gauge choices. For moving hypersurfaces, a graph description—including its nonlinear realization of Lorentz invariance—and hydrodynamic formulations (in light-cone coordinates as well as when choosing the time coordinate of a Lorentz observer as the dependent variable) are presented. A matrix regularization for $M = 2$ (existing for all topologies) is explained in detail for the 2-sphere, as well as multilinear formulations for $M > 2$. The recently found dynamical symmetry that exists for all M and related reconstruction algebras are covered, just as some explicit solutions of the level-set equations. **Jens Hoppe 2013 J. Phys A: Math. Theor. 46 023001 doi:10.1088/1751-8113/46/2/023001 Relativistic membranes** **N Bodendorfer et al** rederive the results of our companion paper, for matching space–time and internal signature, by applying in detail the Dirac algorithm to the Palatini action. While the constraint set of the Palatini action contains second class constraints, by an appeal to the method of gauge unfixing, we map the second class system to an equivalent first class system which turns out to be identical to the first class constraint system obtained via the extension of the ADM phase space performed in our companion paper. Central to analysis is again the appropriate treatment of the simplicity constraint. Remarkably, the simplicity constraint invariant extension of the Hamiltonian constraint, that is a necessary step in the gauge unfixing procedure, involves a correction term which is precisely the one found in the companion paper and which makes sure that the **Hamiltonian constraint derived from the Palatini Lagrangian** coincides with the ADM Hamiltonian constraint when Gauß and simplicity constraints are satisfied. **N Bodendorfer et al** therefore have

rederived new connection formulation of general relativity from an independent starting point, thus confirming the consistency of this framework. **N Bodendorfer et al 2013 Class. Quantum Grav 30 045002 doi:10.1088/0264-9381/30/4/045002 new variables for classical and quantum gravity in all dimensions: II. Lagrangian analysis**

As Kṣemaraja points out, none of the practices taught in the Stanzas on Vibration belong to the Individual Means and so it does not, strictly speaking, concern Spanda doctrine, if that is, we consider the Stanzas to be the basic text of the Spanda school. From Kṣemaraja's point of view, however, the third section of the Aphorisms of Siva (Sivasutra), which is both the last and most extensive, is largely an exposition of this category of practice. The Stanzas and Aphorisms have been traditionally linked together and so, even though we feel that they should be distinguished so far as the Stanzas rather than the Aphorisms teach the Doctrine of vibration as such, we are nonetheless justified in referring to the Aphorisms as its major source. Our exposition of the Individual Means will therefore be largely based on Kṣemaraja's interpretation of the third section of Aphorisms and we will present it, as he does, as an exposition of a possible mystical journey of individualised (anava) consciousness to realisation. We follow Kṣemaraja because he understood the practise taught in the Aphorisms in these terms, thereby not only illustrating for us how it fits into this scheme but also how he understood the basic categories of practice and their relationship to one another. According to Kṣemaraja, the first Aphorism of each section of the Sivasutra characterises the condition and nature of the Self at the corresponding three levels of practice. In other words, they indicate the yogi's basic state at each level in terms of his self-identification. **This identification corresponds to his existential condition as a degree of self- realisation in the process leading to the authentic self-awareness of the liberated. The very first Aphorism starts directly with this, the highest state, by declaring that the Self is pure, dynamic and universal consciousness (caitanya).** This is true for the yogi who has awakened to his authentic nature at the Divine (sambhava) level of being. At the Individual (anava) level, however, the situation has changed. In this sphere of consciousness the intermediate processes of discernment, analysis and classification of perceptions, which bridge the gap in the flow of awareness from the universal subject to a specific object of knowledge, appear to take over the status of the perceiving subjectivity which underlies them. **The universal Self recedes into the background as a pure, undefinable awareness, and the individual ego, consisting of the perceptions, thoughts and emotions generated by the contact between the universal perceiver and the perceived, emerges in the juncture between them.** Thus at this level, as the Aphorisms say, the Self is the mind. This is the Self which moves (atati) from one state of being to another, from one body to the next carrying with it subtle traces left behind by its sensory and mental activity. Together these are said to constitute, and be caused by, the subtle body technically called the 'City of Eight' (puryasfaka) with which consciousness is identified and due to which it is subject to the constant alterations of pleasure, pain and inertia. The Stanzas teach: [The soul] is bound by the City of Eight (puryaffaka) that resides in the **mind, intellect and ego** and consists of the arising of the [five] subtle elements [of sensory perception]. He helplessly suffers worldly pleasure and pain (bhoga) which consists of the arising of mental representations born of that [City of Eight] and so its existence subjects him to transmigration. **Whereas consciousness itself is the subject who practises the Divine Means (Sambhavopayd), the subject who practises the Individual Means is the mind. Unlike the Empowered Means, however, the mind is not directed inwards onto itself. At the Empowered level, enlivened by the direct intuition (pratibha) consciousness has of its own nature, mind ceases to function merely in the paradigmatic, formative manner which gives rise to mental representations, but operates instead as the subtle introverted activity of reflective awareness (vimarsa), the power of consciousness (sakti).** This activity, as we have seen, is the essence of Mantra which, independent of the senses, is no longer restricted in any way. At the Individual level, however, the creative powers of consciousness reflected through the eAtroverted mind are greatly attenuated. All that remains is the power to form thought-constructs and make determined resolutions (sankalpa) which go on to issue through the body into outer action to make the private creations of the mind apparent to others. The Individual Means, therefore, deals with the objectively perceived contents of consciousness and hence with the individual subject as a composite aggregate of objective elements, ranging from the subtle life force (prdrta) to the physical body and its outer environment. The practices belonging to this Means are thus of two types. One is concerned with the individual subject who resides in, and as, the psycho-physical organism; the other with external reality. What this implies essentially is that practice at this level is not concerned as much with the will or **cognitive consciousness** **(refer hiedigger who proposes the same philosophy:**

italics mine) as are the other two Means, but with the power of action applied, in the context of the practice taught in the Aphorisms, to the spiritual activity of Yoga. According to Ksemaraja, the Individual Means culminates in the Empowered state and hence leads to the levels of practice beyond it. 251 This is possible because, despite their differences, there is an essential similarity between them. **The aim of both the Individual and Empowered Means is to purify the discursive representations of differentiated perceptions (vikalpasamskara) 252 and so lead the yogi to the expanded (vikasita) consciousness of the Divine (sambhava) state.** The other levels of practice therefore both sustain and complement it. The activity of individual consciousness can be fully perfected only when it operates through the flow of the conative and cognitive powers which together constitute the pure activity of universal consciousness beyond all means (anupaya). In fact, according to Ksemaraja, all three soteriological types function together in various ways, their corresponding states representing dimensions of the same experience. For example, the upsurge of consciousness (udiyama) which is the supreme, **illuminating intuition (parapralibha)** of the Divine state (sambhava) is concomitant with the gathering together of all the powers of consciousness in the Empowered state. The Divine Means, in other words, leads to the experience of Power (sakti) which in its turn, when fully affirmed, marks the attainment of a permanent contemplative consciousness (turiyatita) at the Divine level which persists unaltered in every state of consciousness. Consequently, Ksemaraja concludes his exposition of the first section of the Aphorisms which exemplifies, according to him, the Divine Means, by saying: Thus we have explained the **first expansion** which starts with [the Aphorism] 'the **Self is pure dynamic consciousness**' (*note this happens in evolutionary process and in normal human being it remains as witness consciousness italics mine*) and expounds the nature of the realisation (prathana) attained through the Divine Means. It is the intuitive insight (samapatti) of Bhairava's nature which is, as we have said, the upsurge of consciousness that quells all bondage, namely, the ignorance of that freedom which makes it manifest. Transforming all things into the nectar of one's own innate bliss, it bestows every yogic accomplishment (siddhi) including mystic absorption in the vitality of Mantra, the highest of them all. Accordingly, we have, in the course of this exposition, explained the nature of Sakti in order to show that the Divine nature (Sambhavarupa) possesses [every] power. Another way in which **the Means are related to one another is illustrated by the recurrence of the same Aphorism in different sections of the Sivasutra which indicates, according to Ksemaraja, that the same practice belongs to more than one Means.** Both times this happens, the Aphorism appears first in the section dealing with the Divine Means and then recurs in that concerned with the Individual Means. In one case, Ksemaraja tells us this is because practice at the Divine level requires no effort whereas at the Individual level, the yogi must exert himself to achieve the same state that at the Divine level dawns spontaneously. At the Empowered level also, as the Sivasutra says, 'effort achieves the goal'. Here, however, because as the Empowered Means is, according to Ksemaraja, predominantly concerned with the contemplation (anusamdhhi) of the vitality of Mantra, the effort exerted is that required to bring the practice of Mantra to fulfillment. It is, as Ksemaraja says, 'the spontaneous effort exerted to grasp the initial expansion of intention to apply oneself to the contemplation [of Mantra]. It is this exertion which wins the favour of the gods of Mantra and identifies the adept with them.' The second case of the same practice being taught in different sections of the Aphorisms concerns the realisation of the **Fourth State of contemplative consciousness (turiya)** in the other three states of waking, dreaming and deep sleep. At the Divine level this takes place by **'violently digesting' (hafhapaka)** the three states in the Fourth. At the Individual level the Fourth state is first experienced at the junctures between the other three states and then induced gradually to spread out from these Centres to pervade the other states like oil extending slowly through a piece of cloth. The difference in this case between the levels of practice is not only that at the Divine level it reaches fulfillment spontaneously, but it is also sudden and complete, leading directly to the liberated state of consciousness Beyond the Fourth (turiyatita) At the Individual level, however, practice is gradual and even when the yogi manages to rise to states of contemplation, he must take care not to fall to lower levels of consciousness. Indeed, until the yogi attains the sudden and direct realisation of perfect enlightenment, whatever be his state of consciousness or level of practice, he is bound to rise and fall because **his contemplative state is necessarily transitory (kadacitka)** however long it may last. The yogi is more prone to these ups and downs the lower his basic state of consciousness. Consequently, the last section of the Sivasutra repeatedly instructs the yogi not only how to rise to higher levels of consciousness and maintain them, but also in what way he is liable to fall from them and how to regain them. Ksemaraja stresses that the rise from one level of consciousness to another is marked by the

transition from a lower Means to a higher. Conversely, a fall from the higher level to the lower entails practice of a lower Means. The measure of the yogi's level of consciousness and that which sustains him in it allowing him to progress further, is his attentiveness (avadhana) to the higher realities he experiences in the more elevated states. Thus the last Aphorism of the second section of the Sivasutra warns the yogi that if his pure awareness (suddhavidya) of his oneness with all things slackens, he will fall from his awakened state to dream the dream of thought-constructs. From Ksemaraja's point of view this means that the negligent yogi must now resort to the Individual Means described in the next section to return to his former, higher Empowered practice in which he experiences this oneness. Ksemaraja expounds practice at the Individual level, as he sees it in the Aphorisms, as extending from one Means to the next. For example, practice at the Individual level diverts the flow of the vital breath (prana) from its more usual course and induces it to enter the Central Channel (susumna) along which it rises as a pure conscious energy (technically called 'kuritfalini'). This leads the yogi to the Empowered state in which he enjoys the pure awareness of unity. If he manages to make it truly his own and it becomes his basic state of being, he enters the Divine plane (sambhavadh) of identity with Siva. The Individual Means is both a point of departure to higher levels of practice and the level to which the yogi returns if he falls. Thus although the practices taught in the last section of the Aphorisms may belong to any one of the three Means, they are collectively treated as part of the Individual Means because they start from it and because it is the yogi's abiding standby if he falls. Let us turn now to the basic practice at the Individual level, as Ksemaraja understands it. This is essentially Yoga. According to the Classical Yoga system taught by Patanjali in the Aphorisms of Yoga (Yogasutra), Yoga is defined as 'the quelling of the fluctuations of the mind' (cittavrttinirodha). The aim is to sever the spiritual essence of the Person (purusa) from the defiling materiality of Nature (prakriti), even though the word 'Yoga' means to 'unite' or 'yoke together.' Here, however, Yoga combines both union and cessation. It is the act (kriya) of removing the latent traces (yasana) of differentiated perceptions (yikalpa) born of the impurities (mala) which contract consciousness (like pravrutti and Nivrutti; italeis mine). This is achieved by uniting all the elements of experience (tattva) together in the wholeness of the activity of consciousness. As Jayaratha explains: The [wise] consider Yoga to be the union of one thing with another/ thus, in accord with this dictum, Yoga is the [act] of uniting [all] the metaphysical principles together within consciousness . . . , Ksemaraja seeks initially to establish the best form of Yoga for the yogi to practice at the Individual level. His sources are two Tantras he knew well and considered to be amongst the most important, namely, The Tantra of (Siva's Third) Eye (Netratanttra) and The Tantra of the Liberated Bhairava (Svacchandabhairavatantra). The basic model is that of the Eight-limbed Yoga (asfariga) taught by Patanjali which consists of: 1) The five restraints (yama\ namely, abstention from violence (ahirrisa), falsehood (satya), dishonesty (asteya), sexual intercourse (brahmacarya) and desire for more than the essential (aparigraha) 2) The five disciplines (niyama), namely, cleanliness (sauca), contentment (santo\$), austerity (tapas\ study (svadhyaya) and reverence for God (Uvarapranidhand) 3) Posturing of the body (asana) in a manner conducive to the practice of meditation and physical health. 4) Regulation of the breath (pranayama) 5) Withdrawal of the senses from their objects (pratyahara) 6) Focusing of attention (dharana) 7) Meditation (dhyana), that is, steady, uninterrupted concentration. 8) Contemplation (samadhi). Ksemaraja rejects Patanjali's system because he believes it to be a form of Yoga that can, at best, lead only to limited yogic attainments (mitasiddhi). In the Netratanttra, however, Siva teaches a different, higher form of the Eight Limbs of Yoga which lead to perfect penetration into the supreme, transcendental principle 280 of which the Netratanttra says: Speech cannot express, nor the eye see, the ears hear, or the nose smell, the tongue taste, the skin touch or the mind conceive that which is eternal. Free of all colour and flavour, endowed with all colours and flavours, it is beyond the senses and cannot be objectively perceived. O goddess, those yogis who attain it become immortal gods! By great practice and supreme dispassion . . . one attains Siva, the supreme imperishable, eternal and unchanging reality. A necessary preliminary of all Tantric Yoga is a process technically called the 'purification of the elements' (bhutasuddhi), through which the body is homologized with the macrocosm and so made a fit vessel for the pure, conscious presence of the Deity within it. Ksemaraja equates this with the meditation (dhyana) which, according to the Mdlinivijayatantra, characterises the Individual Means. In order to practice this meditation the yogi must visualise the dissolving away of all the forces in the body. There are two ways in which this can be done. The first is called 'the contemplation of dissolution' (layabhdvana). Through it the progressive differentiation of consciousness from its causal, pre-cosmic form to its phenomenal

manifestation is reversed As the Vijñanabhairava teaches: "One should meditate on the All in the form of the Paths of the world- orders etc. considered in their gross, subtle and supreme forms until, at the end, the mind dissolves away." **Mediated by consciousness, the macrocosm rests in the microcosm which is emitted along with it successively in the emptiness of the individual subject, vital breaths, mind, psychic nerves (ndtṛi), senses and external body.** The yogi reproduces this process by visualising the totality of reality including the world-systems, metaphysical principles and cosmic forces along with the Mantras, letters and syllables which represent them, as arising successively throughout the psycho-physical body so as to constitute it. Deployed in this way they form the Cosmic Path along which the yogi ascends, absorbing as he does so, the lower elements into the higher, thus strengthening and extending his unifying awareness (anusamdhnd) of the configuration of the Path. Thus, moving from the gross elements constituting the outer physical body, to pure sensations (tanmdtra), then to the senses and mind back to their primordial source, the yogi rises from the **embodied subjectivity** of the waking state to the Fourth State (turiya) of contemplation where he is one with the pervasive intent which initiates the creative vision of consciousness. Abhinava writes: Once [the yogi] has known [this] Path in its completeness, he must then dissolve it into the deities who sustain it and these successively into the body, breath, **mind [and emptiness]** as before, and all these into his own consciousness. Once this is full and an object of constant worship, it destroys, like the fire at the end of time, the ocean of transmigration. Thus, the second method Kṣemaraja teaches to dissolve away the diversity of sensory, mental and physical energies into the unity of consciousness is a meditation on the Fire of Consciousness (dahacintḍ) which the yogi visualises as burning away all division. At the Divine level (sambhavopaya) the yogi witnesses the sudden and violent withdrawal of all objectivity into the pure ego (aharri), like the pouring of fuel into a raging fire. He does not need to visualise this process but merely attend to it with a passive, receptive attitude. At the Individual level the yogi must exert his imagination to induce this process and so rise to the Divine level through the Empowered. The Vijñanabhairava teaches: Visualise the fortress [of your body] burning with the Fire of Time (kalagni) risen from the Abode of Time; then at the end peace manifests. The Fire of Time {kalagni} resides underneath the hell worlds at the bottom of the Cosmic Egg (Brahman^a). It issues from Ananta — a form of Siva who presides over the lower regions. He floats on a boat in the causal waters supporting the Egg, his mind all the while fixed on Bhairava. The flames of the Fire of Time rise up to the hell-worlds heating them intensely and radiate its energy throughout the universe. At the end of each period of creation the flames rise higher and destroy the old cosmic order to make room for a new one. At the microcosmic level the yogi reproduces this process by mentally placing the letters of the alphabet, in the prescribed order, on the limbs of his body starting from the left toe to the top of the head. As his attention progresses upwards, he visualises the Fire of Time moving with it in such a way that his bodily consciousness, together with the **universe of differentiated perceptions**, is gradually burnt away leaving in its place the white ashes of the undivided light of consciousness. Kṣemaraja considers this meditation (dhyana) to be a limb of a programme of yogic practice at the Individual level 291 of which the remaining limbs are as follows: Posture (A sana). The yogi fixes his attention on the centre between the inhaled and exhaled breath, absorbing in this way the flux of his awareness into the unfolding power of knowledge which rises initially as the upward flowing breath (udanaprana) in the Central Channel (suṣumna) between the other two breaths. The Prank aspect of this flow disappears as it moves upward and the yogi experiences the **spontaneous rise of the omniscience of consciousness within himself.** The mind reverts back to its original, pervasive conscious nature and understands the infinite fact of Siva's omnipresence. This is the firm seat (asana) upon which the yogi sits to practice. Regulation of the Breath (Pranayama) .To regulate the movement of the breath, the yogi must first cleanse the right and left channels of the ascending and descending breath by blocking the left nostril while exhaling and the right while inhaling a few times. This ensures that the movement of the breath is firm and evenly distributed. Next, without attempting to control it in any way, he attends to the flow of his breathing. As the mind becomes steadier and in closer harmony with the rhythm of its movement, the duration of each inhalation and exhalation gradually alters **until they become equal.** At this stage they unite and merge in the upward flowing current of vitality in the Central Channel (Susumna). This is when true Pranayama begins. The yogi's mind pure and tranquil, he returns, as it were, to a prenatal state and the external breathing cycle is internalised, so that it no longer moves through the lungs but passes directly to the universal source of vitality. The yogi, now at the Empowered level of practice, experiences this movement as travelling from the Heart centre upwards to a point distant twelve fingers above the head where it

merges in the void of consciousness. Free of its outer gross form, the breath moves freely through the Central Channel and soon transcends even this subtle movement to become one with the supreme vibration of consciousness. In this way, the yogi's breathing becomes one with the spontaneous rise and fall of energy from the bosom of the absolute. Abhinava quotes the Tantra of the Line of Heroes (Viravalitantra) as saying: When, by constantly merging the mind in Siva, Who is the pure conscious nature, the Sun and Moon [of the two breaths] have dissolved away and the Sun of Life, which is one's own consciousness, has reached the twelve-finger space, this is termed liberation. Breath control [at this stage] serves no useful purpose. Breath control which merely inflicts pain on the body is not to be practised. He who knows this secret is both themselves liberated and liberates others.

Focusing of Attention (Dharana): Attention is fixed on the psychic centres in the body corresponding to the five gross elements. In this way the vital breath is successively directed to these centres from the Heart of consciousness to refresh and stimulate their activity. First it moves to the Earth centre in the throat which regulates the firmness of the bones and flesh of the body; then to the Water centre in the glottis responsible for the balance of the bodily fluids. After this it travels to the navel which is the Fire centre dealing with digestion and anabolism and catabolism in general. It then moves to the Wind centre in the toe of the left foot which governs the movement of gases to and from the cells via the circulatory system. When the yogi has thus achieved control over these forces, the breath rises from the Heart to the top of the head and he becomes master of the Ether element and so attains every yogic power.

Meditation (Dhyana) The highest form of meditation stills the flux of the qualities (guna) and induces the mind into a state of contemplative absorption. The object of this meditation is the supreme and pervasive divinity of the pure subject whose true nature is known to none but himself alone (svasatpvedya). The yogi attains him by merging into the constant flow of awareness that streams into the Light which illumines his own nature. Contemplation (Samadhi). The yogi rises to the level of contemplation when the awareness he has of himself and the things around him become one and he realises his own identity with Siva, **(or Rama the omniscient: cosmic general ledger)** the sole reality. The aim of this Yoga in all its phases is to achieve the Fourth State of consciousness (turiya) beyond the three states of waking, dreaming and deep sleep and to then ultimately reach the liberated state Beyond the Fourth (turiyatita). These five states correspond to: (a) Siva's activity (vyapara), that is, His power of action; (b) Siva's Lordship (adhipatya), which is His power of knowledge; (c) the absence of these two, which corresponds to Siva's power of will; (d) His exertion (prerakatva), which contains all the cycles of creation and destruction and, (e) the rest Siva enjoys in His own nature, which is His power of consciousness. The first three states, when divorced from the last two, belong to the sphere of transmigratory existence. The Fourth and Beyond the Fourth on the other hand are higher, **supramundane (alaukika)** states of consciousness in which the yogi **enjoys bliss and repose (visranti) in his own nature by penetrating (samavesa) into the universal consciousness of the Self**, through which he ultimately becomes liberated (jivanmukta). Beyond the Fourth is the state of awareness ParamaSiva Himself enjoys when duality has entirely disappeared and everything is realised to be one with consciousness. The Fourth is the state of awareness of the yogi who, catching hold of the pure subjectivity (upalabdhrta) flowing through the lower three states, is still actively eliminating his sense of duality. While the former is the supreme subject as T consciousness (aham), the latter is the pure awareness (prama) or **I-nesses (ahanta)** of the subject which encompasses the lower states, giving them life and **uniting them** together. As such, the Fourth State is the reflective awareness of one's own nature shining in all three states at one with them. The fact that we recall that we slept well is proof that this state of consciousness persists even in deep sleep. Indeed, if the flow of Turiya could somehow be brought to a halt, all the other states of consciousness would come to an end in the absence of the pure subjectivity which makes them, and their contents, manifest. The states of waking, dreaming and deep sleep correspond to the form of awareness consciousness **assumes when it it predominantly manifest as the object, means of knowledge and individual subject, respectively.** Turiya is the pure awareness (prama) that both transcends them and merges them all into itself. 300 As such, it appears as the triad of deed, means and agent in the pure act (vyapara) of consciousness unsullied by any outer reality. Abhinava explains: **{Turiya) transcends the three aspects of *form\ 'sight' and T consisting as it does of the pure act of 'seeing'; therefore any means [by which this state could be realised] has merely a [provisional] instrumental value. It is, in other words, pure subjectivity of the nature of absolute freedom, independent of all external means. This is the state of consciousness called Turiya, luminous with its own light. Turiya is thus not just a psychological state but the supreme creative power (para sakti) of consciousness, the Goddess (samviddevi) who**

generates and withdraws the entire universe of subject, object and means of knowledge. In the Heart of Recognition Kshemaraja explains: Whenever the extroverted [conscious] nature rests within itself, external objectivity is withdrawn [and consciousness] is established in the inner abode of peace which threads through the flux of awareness in every [externally] emanated [state]. Thus Turiya, the Goddess of **Consciousness, is the union of creation, persistence and destruction. *Read as cosmic consciousness; Kshemaraja does not distinguish like we have done, individual, collective and cosmic and of three states*** She emanates every individual [cycle] of creation and withdraws it. Eternally full [of all things] and [yet] void [of diversity] She is both and yet neither, shining radiantly as non-successive [consciousness] alone. The yogi is fully absorbed in this state of consciousness and takes possession of its power when he is able to rise from contemplation (samadhi) carrying with him the abiding awareness of Turiya throughout his waking, dreaming and deep sleep. When he achieves this constantly, he continues to experience these states individually, but they no longer obscure the insight (pratibha) he has acquired because he realises that they are all aspects of the bliss of Turiya. Thus, while the common man calls this state the 'Fourth' (turiya) because he cannot experience it directly and knows only that it is beyond the other three, the yogi calls it 'Beyond Form' (rupatita) because it transcends the detachment of the state of deep sleep which, devoid of objective content, is the naked form' of the individual subject tending towards the fullness of consciousness. Those who are on the path of knowledge (jnaniri) call it the 'Whole*' (pracaya) because, in this state, they see the entire universe gathered together in one place. **'Supra-mental Awareness' (manonmana)** is the name given to the experience of Turiya in the waking state. The yogi in this state moves and lives in the world of waking experience free of all disturbing thoughts while abiding in the transcendental silence beyond the activities of the mind. 'Infinite' is the name of the experience of Turiya while dreaming because, free of the limitations imposed upon the body by time and space, the yogi enjoys the unlimited expanse of the Self. **When Turiya is experienced in deep sleep, the yogi's state is called 'All things' (sarvdrtha) because in it he discovers his freedom from limitations in this, the most contracted state of human consciousness.** The yogi who manages to maintain Turiya- consciousness comes to experience the three states of waking, dreaming and deep sleep as the constant flow of the bliss of consciousness in which all traces of the relative distinction between these states and their contents is eradicated. Following the stream of Turiya to its highest level (para katfha), he reaches the state Beyond the Fourth (turiydrta), which is the universal consciousness (caitanya) of the Self. Here the yogi comes to **rest within his own nature.** Plunged in the vast, waveless ocean of the consciousness and bliss (ciddnanda) of the state Beyond the Fourth, the yogi becomes Siva, the Free One (svacchanda), and thus wanders freely, practising the Yoga of Freedom. Kshemaraja equates the Fourth State with the pure (suddha), innate (sahaja) knowledge that one's own conscious nature is all things. It is the Supra-mental State (unmana) in which Siva's pervasive presence is experienced once the Yoga practised at the Individual level attains fruition at the Empowered. What the yogi must do, once consciousness is elevated to grasp the Fourth State, is make it constant. He must forcefully lay hold of it within himself and not release his grip until it becomes permanent. Then he travels 'Beyond the Fourth' to enlighten- ment. Before this ultimate attainment the yogi inevitably falls. The forces operating within consciousness that limit and obscure it throw him down whenever they possibly can. The only way the yogi can defend himself against them is to maintain a constant attentive awareness of the Fourth State. He falls when he is distracted but when he attends carefully to his pure conscious nature, he realises that every aspect of his state of being, including the forces that lead him astray, are one with the pulsing flux of his own consciousness and so cannot affect him. These powers, which are the energies of Matrka we have already discussed, are not the only obstacles the yogi must overcome. He must, for example, also resist the temptation to rest content with the miraculous yogic powers (siddhi) he acquires in the course of his spiritual develop- ment. Again to do this he must practice Yoga. Similarly, in order to pervade the Fourth State gradually through the other states in the manner proper to practice at the Individual level, the method is the same. He must practise the higher yoga of the Tantras which, turning his mind inwards and freeing it from discursive representations, allows him to penetrate into the Supreme Principle. Once the yogi has attained this contemplative state, his main problem is to make it permanent. In the introverted state the gross external movement of the breath is suspended and with it the activity of the intellect, mind, individualised consciousness, powers of the senses and the ego. 313 When the yogi rises out of this state, he is liable to fall again into the lower order of creation generated by Maya if he does not maintain his awareness of the higher reality he has experienced and allows his awakened, illumined insight to be obscured by the

dream-like vision of thought-constructs. 314 Naturally, the yogi must rise out of the introverted condition of suspension. It is inherent in the very nature of reality that it should move out of itself. 315 **Pure, universal consciousness initially transforms itself into the vital breath 316 charged with the impression (iyasana) of the power of awareness attained through introversion.** By attending to the pulse (spanda) of the breath as it moves out of the absolute, the yogi can develop an intuitive sense of the inherent unity of all he will perceive in the mental and physical spheres created by the outpouring of consciousness. In this way he realises that his own nature is everywhere present in all he perceives and that all things thus reside within him. Blessed with this insight his consciousness remains free and unlimited even at the individual level where the breath, mind, senses and body are active. If the yogi fails to do this, he finds himself once again beset by the strictures of his embodied existence and must, as before, try to pervade all his other states of consciousness with the aesthetic delight (rasa) and wonder (camatkara) of the Fourth State he experienced in contemplation. Again this means that he must strengthen his pure, empowered awareness that his universal nature manifests as all things. In this way he discovers Siva's presence in every sphere of individualised consciousness ranging from the breath to outer objectivity. The yogi's mind then becomes tranquil and undistracted because wherever it may wander, the yogi perceives only Siva, his authentic nature. **Consciousness is thus freed of all external referents and the yogi's subjectivity is purified of all identification with the body or anything else that belongs to the objective sphere.** The yogi then becomes detached from the opposites of pleasure and pain and is transcendently free (kevalin). The yogi is again, however, liable to fall if he allows himself to **get entangled in the play of opposites.** This fall is more serious than the others because, although he is caught by the confining restrictions of individualised consciousness as before, he is now also affected by karma. Fleeing from pain in the pursuit of pleasure he is bound to act (karma) to minimise one and maximise the (a **maximization problem is mentioned elsewhere about the svabhava of human beings to enhance pleasure reduce deprivation**) other and so is thrown down to the lowest level of embodied subjectivity (sakala). In order to regain his lost state, he must ascend gradually, by Siva's grace, from one order of subjectivity to the next and so free himself progressively of the limitations of the lower levels to gain the greater freedom and expansion of the higher. As he progresses, the objective sphere also evolves from the grossest perceptions of physical objects outside the lowest order to subjectivity, through to the subtler inner, mental perceptions to finally reach the order of subjectivity that contains objectivity within itself and is free to externalise it at will. The degree to which this process develops depends, as before, on the yogi's awareness of the Fourth State. In consonance with the general principle that the remedy should suit the defect, the yogi is instructed to seek this higher state of consciousness in the wonder (camatkara) or delight (ananda) he feels in moments of intense physical pleasure. At first he experiences this subtle consciousness **for an instant in the subjective sphere.** If he manages to catch hold of it, it becomes more intense as the cognitive and objective spheres are also gradually pervaded and vitalised by it. Occasions for this practice are, for example, the sense of satisfaction one feels after a good meal or the aesthetic delight one experiences when listening to good music or the pleasure of sexual union with the Tantric consort or even solitary sexual excitation. In these moments of delight the yogi can penetrate momentarily into his own authentic Siva-nature (sambhava) through the empowered contact (saktasparsa) he makes with it in the freedom of the pure subjectivity of the Fourth State. If the yogi develops his awareness of this higher level of consciousness and maintains it, he eventually experiences it constantly. Clearly, what prevents the yogi from attending to his state of consciousness rather than the circumstances which induce it is the craving for pleasure (abhilasha) born of ignorance — the source of every impurity which clouds consciousness. Craving directs the yogi's attention towards outer, worldly things and so he is caught in the **net of thought-constructs.** To free himself of his worldly desires and reverse this binding extroversion, the yogi must eradicate its cause. To be freed of all the ups and downs of the path and no longer be tormented by the possibility of a fall, the yogi must see reality perfectly and completely. This insight is itself liberation and the moment it dawns the yogi is instantly freed. This sudden realisation is the goal of Tantric Yoga. Accordingly the Tantra declares: **"He, who perceives reality directly, even for the brief moment it takes to blink, is liberated that very instant and never reborn again."** Although the yogi's body and mind continue to function as before, they are like mere outer coverings which contain, but do not obscure, the mighty, universal consciousness which operates through them. The yogi's body is the universe, the senses the energies that vitalise it, his mind Mantra, the rhythm of his breath the pulse of time and his inner nature pure, dynamic consciousness. Raised above

all practice, and hence all possibility of falling to lower levels, the yogi realises that he has always been free and that his journey through the dark land of Maya was nothing but a dream, a construct of his own imagination. (Models are given in one of the papers of the series due to spatial constraints) Relationship betwixt string theory (note the spade remarks above) and holographic principle has been enucleated and expatiated in literature. Some exemplar work is sententiously mentioned in the following: The ratio of shear viscosity to volume density of entropy can be used to characterize how close a given fluid is to being perfect. Using string theory methods, we show that this ratio is equal to a universal value of $\hbar/4\pi k_B$ for a large class of strongly interacting quantum field theories whose dual description involves black holes in anti-de Sitter space. **P K. Kovtun, D. T. Son, and A. O. Starinets** provide evidence that this value may serve as a lower bound for a wide class of systems, thus suggesting that black hole horizons are dual to the most ideal fluid **Viscosity in Strongly Interacting Quantum Field Theories from Black Hole Physics Phys. Rev. Lett. 94, 111601 – Published 22 March 2005 P K. Kovtun, D. T. Son, and A. O. Starinets** According to 't Hooft the combination of quantum mechanics and gravity requires the three-dimensional world to be an image of data that can be stored on a two-dimensional projection much like a holographic image. The two-dimensional description only requires one discrete degree of freedom per Planck area and yet it is rich enough to describe all three-dimensional phenomena. After outlining 't Hooft's proposal we give a preliminary informal description of how it may be implemented. One finds a basic requirement that particles must grow in size as their momenta are increased far above the Planck scale. The consequences for high-energy particle collisions are described. The phenomenon of particle growth with momentum was previously discussed in the context of string theory and was related to information spreading near black hole horizons. The considerations of this paper indicate that the effect is much more rapid at all but the earliest times. In fact the rate of spreading is found to saturate the bound from causality. Finally **Leonard Susskind** considers string theory as a possible realization of 't Hooft's idea. The light front lattice string model of Klebanov and Susskind is reviewed and its similarities with the holographic theory are demonstrated. The agreement between the two requires unproven but plausible assumptions about the nonperturbative behavior of string theory. Very similar ideas to those in this paper have long been held by Charles Thorn. © 1995 American Institute of Physics **The world as a hologram Leonard Susskind J. Math Phys. 36, 6377 (1995); <http://dx.doi.org/10.1063/1.531249> Ofer Aharony, , Steven S. Gubser, , Juan Maldacena, c, , Hiroshi Ooguri, e, , Yaron Oz, , review the holographic correspondence between field theories and string/M theory, focusing on the relation between compactifications of string/M theory on Anti-de Sitter spaces and conformal field theories. They review the background for this correspondence and discuss its motivations and the evidence for its correctness. They describe the main results that have been derived from the correspondence in the regime that the field theory is approximated by classical or semiclassical gravity. We focus on the case of the supersymmetric gauge theory in four dimensions, but discuss also field theories in other dimensions, conformal and non-conformal, with or without supersymmetry, and in particular the relation to QCD. We also discuss some implications for black hole physics. **Physics Reports Volume 323, Issues 3–4, January 2000, Pages 183–386 Large N field theories, string theory and gravity Ofer Aharony, , Steven S. Gubser, , Juan Maldacena, c, , Hiroshi Ooguri, e, , Yaron Oz, , doi:10.1016/S0370-1573(99)00083-6 T. Banks, W. Fischler, S. H. Shenker, and L. Susskind** suggest and motivate a precise equivalence between uncompactified 11-dimensional M theory and the $N=\infty$ limit of the supersymmetric matrix quantum mechanics describing D0 branes. The evidence for the conjecture consists of several correspondences between the two theories. As a consequence of supersymmetry the simple matrix model is rich enough to describe the properties of the entire Fock space of massless well separated particles of the supergravity theory. In one particular kinematic situation the leading large distance interaction of these particles is exactly described by supergravity. The model appears to be a nonperturbative realization of the holographic principle. The membrane states required by M theory are contained as excitations of the matrix model. The membrane world volume is a noncommutative geometry embedded in a noncommutative spacetime. DOI: <http://dx.doi.org/10.1103/PhysRevD.55.5112> **M theory as a matrix model: A conjecture Phys. Rev. D 55, 5112 – Published 15 April 1997 T. Banks, W. Fischler, S. H. Shenker, and L. Susskind** These TASI lectures review the Holographic principle. The first lecture describes the puzzle of black hole information loss that led to the idea of Black Hole Complementarity and subsequently to the Holographic Principle itself. The second lecture discusses the holographic entropy bound in general space-times. The final two lectures are**

devoted to the ADS/CFT duality as a special case of the principle. The presentation is self contained and emphasizes the physical principles. Very little technical knowledge of string theory or supergravity is assumed. Report number: SU-ITP 99-14, KUL-TF-2000/03 Cite as: arXiv: hep-th/0002044 **TASI lectures on the Holographic Principle Daniela Bigatti, Leonard Susskind** The notion of a space-time uncertainty principle in string theory is clarified and further developed. The motivation and the derivation of the principle are first reviewed in a reasonably self-contained way. It is then shown that the nonperturbative (Borel summed) high-energy and high-momentum transfer behavior of string scattering is consistent with the space-time uncertainty principle. It is also shown that, as a consequence of this principle, string theories in 10 dimensions generically exhibit a characteristic length scale which is equal to the well-known 11 dimensional Planck length $g_s^{1/3} \ell_s$ of M-theory as the scale at which stringy effects take over the effects of classical supergravity, even without involving D-branes directly. The implications of the space-time uncertainty relation in connection with D-branes and black holes are discussed and reinterpreted. Finally, we present a novel interpretation of the Schild-gauge action for strings from the viewpoint of noncommutative geometry. This conforms to the space-time uncertainty relation by manifestly exhibiting a noncommutativity of quantized string coordinates between, dominantly, space and time. We also discuss the consistency of the space-time uncertainty relation with S and T dualities. Copyright (c) 2000 Progress of Theoretical Physics **String Theory and the Space-Time Uncertainty Principle Tamiaki Yoneya Oxford Journals Science & Mathematics Progress of Theoretical Physics Volume 103, Issue 6Pp. 1081-112** In this Letter a recently proposed gravity dual of noncommutative Yang-Mills theory is derived from the relations between closed string moduli and open string moduli recently suggested by Seiberg and Witten. The only new input one needs is a simple form of the running string tension as a function of energy. This derivation provides convincing evidence that string theory integrates with the holographical principle and demonstrates a direct link between **noncommutative Yang-Mills theory and holography**. DOI: <http://dx.doi.org/10.1103/PhysRevLett.84.2084> **Holography and Noncommutative Yang-Mills Theory Phys. Rev. Lett. 84, 2084 – Published 6 March 2000 Miao Li and Yong-Shi Wu Petr Hořava** suggests that M theory could be nonperturbatively equivalent to a local quantum field theory. More precisely, we present a “renormalizable” gauge theory in eleven dimensions, and show that it exhibits various properties expected of quantum M theory, most notably the holographic principle of ’t Hooft and Susskind. The theory also satisfies Mach’s principle: A macroscopically large space-time (and the inertia of low-energy excitations) is generated by a large number of “partons” in the microscopic theory. **Petr Hořava** argues that at low energies in large eleven dimensions, the theory should be effectively described by eleven-dimensional supergravity. This effective description breaks down at much lower energies than naively expected, precisely when the system saturates the Bekenstein bound on energy density. He shows that the number of partons scales like the area of the surface surrounding the system, and discuss how this holographic reduction of degrees of freedom affects the cosmological constant problem. proposal is put forth for the holographic field theory as a candidate for a covariant, nonperturbative formulation of quantum M theory. DOI: <http://dx.doi.org/10.1103/PhysRevD.59.046004> **M theory as a holographic field theory Phys. Rev. D 59, 046004 – Published 26 January 1999 Petr Hořava** The recent proposal on the correspondence between the super-Yang–Mills theory and string theory in the Penrose limit of the $AdS_5 \times S^5$ geometry involves a few puzzles from the viewpoint of holographic principle, especially in connection with the interpretation of times. To resolve these puzzles, we propose to interpret the PP-wave strings on the basis of tunneling null geodesics connecting boundaries of the AdS geometry. Our approach predicts a direct and systematic identification of the S-matrix of Euclidean string theory in the bulk with the short-distance structure of correlation functions of super-Yang–Mills theory on the AdS boundary, as an extension of the ordinary relation in supergravity–CFT correspondence. Holography requires an infinite number of contact terms for interaction vertices of string field theory and constrains their forms in a way consistent with supersymmetry. Copyright © 2003 Elsevier B.V. All rights reserved. **Nuclear Physics B Volume 665, 18 August 2003, Pages 94–128 Holographic reformulation of string theory on $AdS_5 \times S^5$ background in the PP-wave limit Suguru Dobashi, Hidehiko Shimada, Tamiaki Yoneya doi:10.1016/S0550-3213(03)00460-7** The strongest adversary in quantum information science is decoherence, which arises owing to the coupling of a system with its environment¹. The induced dissipation tends to destroy and wash out the interesting quantum effects that give rise to the power of quantum computation², cryptography² and simulation³. Whereas such a statement is true for many forms of dissipation, **Frank**

Verstraete1, Michael M. Wolf2 & J. Ignacio Cirac3 show here that dissipation can also have exactly the opposite effect: it can be a fully fledged resource for universal quantum computation without any coherent dynamics needed to complement it. The coupling to the environment drives the system to a steady state where the outcome of the computation is encoded. In a similar vein, they show that dissipation can be used to engineer a large variety of strongly correlated states in steady state, including all stabilizer codes, matrix product states⁴, and their generalization to higher dimensions⁵. **Letter abstract Nature Physics 5, 633 - 636 (2009) Published online: 20 July 2009 | doi: 10.1038/nphys1342** Quantum computation and quantum-state engineering driven by dissipation **Frank Verstraete1, Michael M. Wolf2 & J. Ignacio Cirac3** Nonlocal twist operators are introduced for the $O(n)$ and **Q-state Potts models in two dimensions which count the numbers of self-avoiding loops (respectively, percolation clusters) surrounding a given point. Their scaling dimensions are computed exactly. This yields many results: for example, the number of percolation clusters** which must be crossed to connect a given point to an infinitely distant boundary. Its mean behaves as $(1/33\sqrt{\pi}) |\ln(p_c - p)|$ as $p \rightarrow p_c^-$. As an application **John Cardy** computes the exact value $3\sqrt{2}$ for the conductivity at the spin Hall transition, as well as the shape dependence of the mean conductance in an arbitrary simply connected geometry with two extended edge contacts. DOI: <http://dx.doi.org/10.1103/PhysRevLett.84.3507> **Linking Numbers for Self-Avoiding Loops and Percolation: Application to the Spin Quantum Hall Transition Phys. Rev. Lett 84, 3507 – Published 17 April 2000 John Cardy Ferenc Iglói, Loïc Turban, Dragi Karevski, and Ferenc Szalma** consider the Ising model and the directed walk on two-dimensional layered lattices and show that the two problems are inherently related: zero-field thermodynamical properties of the Ising model are contained in the spectrum of the transfer matrix of the directed walk. The critical properties of the two models are connected to the scaling behavior of the eigenvalue spectrum of the transfer matrix which is studied exactly through renormalization for different self-similar distributions of the couplings. The models show very rich bulk and surface critical behaviors with nonuniversal critical exponents, coupling-dependent anisotropic scaling, first-order surface transition, and stretched exponential critical correlations. It is shown that all the nonuniversal critical exponents obtained for the aperiodic Ising models satisfy scaling relations and can be expressed as functions of varying surface magnetic exponents. DOI: <http://dx.doi.org/10.1103/PhysRevB.56.11031> **exact renormalization-group study of aperiodic Ising quantum chains and directed walks Phys. Rev. B 56, 11031 – Published 1 November 1997 Ferenc Iglói, Loïc Turban, Dragi Karevski, and Ferenc Szalma** Using exact expressions for the persistence probability and for the leading eigenvalue of the Fokker-Planck operator of a random walk in a random environment, we establish a fundamental relation between the statistical properties of anomalous diffusion and the critical and off-critical behavior of random quantum spin chains. Many exact results are obtained from this correspondence, including the space and time correlations of surviving random walks and the distribution of the gaps of the corresponding Fokker-Planck operator. In turn **Ferenc Iglói and Heiko Rieger** derives analytically the dynamical exponent of the random transverse-field Ising spin chain in the Griffiths-McCoy region. DOI: <http://dx.doi.org/10.1103/PhysRevE.58.4238> **Anomalous diffusion in disordered media and random quantum spin chains Phys. Rev. E 58, 4238 – Published 1 October 1998 Ferenc Iglói and Heiko Rieger** To gain deeper insight into the dynamics of complex quantum systems we need a quantum leap in computer simulations. We cannot translate quantum behaviour arising from superposition states or entanglement efficiently into the classical language of conventional computers. The solution to this problem, proposed in 1982 (ref. 1), is simulating the quantum behaviour of interest in a different quantum system where the interactions can be controlled and the outcome detected sufficiently well. **A. Friedenauer1, H. Schmitz1, J. T. Glueckert, D. Porras & T. Schaetz** study the building blocks for simulating quantum spin Hamiltonians with trapped ions². They experimentally simulate the adiabatic evolution of the smallest non-trivial spin system from paramagnetic into ferromagnetic order with a quantum magnetization for two spins of 98%. We prove that the transition is not driven by thermal fluctuations but is of quantum-mechanical origin (analogous to quantum fluctuations in quantum phase transitions³). We observe a final superposition state of the two degenerate spin configurations for the ferromagnetic order ($|\uparrow\uparrow\rangle_{\text{right fence}} + |\downarrow\downarrow\rangle_{\text{right fence}}$), corresponding to deterministic entanglement achieved with 88% fidelity. This method should allow for scaling to a higher number of coupled spins², enabling implementation of simulations that are intractable on conventional computers. **Letter Nature Physics 4, 757 - 761 (2008) Published online: 27 July 2008 | doi:10.1038/nphys1032** Subject Categories:

Quantum physics | Atomic and molecular physics Simulating a quantum magnet with trapped ions A. Friedenauer¹, H. Schmitz¹, J. T. Glueckert, D. Porras & T. Schaetz This article gives a comprehensive description of the fractal geometry of conformally-invariant (CI) scaling curves, in the plane or half-plane. It focuses on deriving critical exponents associated with interacting random paths, by exploiting an underlying quantum gravity (QG) structure, which uses **KPZ maps relating exponents in the plane to those on a random lattice, i.e., in a fluctuating metric. This is applied to critical models, like O(N) and Potts models, and to the Stochastic Loewner Evolution (SLE). The multifractal (MF) function $f(\alpha, c)$ of the harmonic measure near any CI fractal boundary, is given as a function of the central charge c of the associated CFT. The Hausdorff dimensions $D_{\{H\}}$ of a non-simple scaling curve or cluster hull, and $D_{\{EP\}}$ of its external perimeter or frontier, are shown to obey the duality equation $(D_{\{H\}}-1)(D_{\{EP\}}-1)=1/4$, valid for any c .** The universal mixed MF spectrum $f(\alpha, \lambda; c)$ describing the local spiralling rate λ and singularity exponent α of the potential near any CI scaling curve is given. The duality between simple and non-simple random paths is established via symmetry of the KPZ quantum gravity map. An extended dual KPZ relation is introduced for the $SLE_{\{\kappa\}}$, which commutes with the κ to $\kappa'=16/\kappa$ duality. This gives the SLE exponents from simple QG rules, established from the general structure of correlation functions of arbitrary interacting random sets on a random lattice. Journal reference: {\it Fractal Geometry and Applications: A Jubilee of Benoît Mandelbrot} (M. L. Lapidus and M. van Frankenhuysen, eds.), Proc. Symposia Pure Math. vol. 72, Part 2, 365-482 (AMS, Providence, R.I., 2004) arXiv: math-ph/0303034 (or arXiv: math-ph/0303034v2 for this version) **Conformal Fractal Geometry and Boundary Quantum Gravity Bertrand Duplantier** Hume, For Deleuze, Considers The Mind To Be A System Of Associations Alone, A Network Of Tendencies (ES 25): “We Are Habits, Nothing But Habits – The Habit Of Saying ‘I’. Perhaps There Is No More Striking Answer To The Problem Of The Self.” (ES X.) The Mind, Affected By The Natural Principle Of Association, Becomes Human Nature, From The Ground Up: Empirical Subjectivity Is Constituted In The Mind Under The Influence Of The Principles Affecting It; The Mind Therefore Does Not Have The Characteristics Of A Preexisting Subject. (ES 29) These Associations Account Not Only for Experience in the Basic Sense, But up To the Highest Level of Social and Cultural Life: This Is The Basis For Hume’s Rejection (E) Of A Social Contract Model Of Society (Such As Hobbes’), In Favour Of Convention Alone. Morals, Feelings, Bodily Comportment, All Of These Elements Of Subjectivity Are Explained, Not By Transcendental Structures, Such As Kant Will Propose, But By The Immanent Activity Of Association. Once This Habitual Structure Of The Self Is In Place, Deleuze Suggests, The Humean Concept Of Belief Comes Into Play , Which Is Resolutely A Central Part Of Human Nature. It Describes The Particularly Human Way Of Going Beyond The Given. When We Expect The Sun To Come Up Tomorrow, We Do Not Do So Because We Know That It Will, But Because Of A Belief Based On A Habit. This In Turn Reverses The Hierarchy Of Knowledge And Belief, And Results, For Deleuze, In A, “Great Conversion Of Theory To Practice.” (PI 36) Every Act Of Belief Is A Practical Application Of Habit, Without Any Reference To An A Priori Ability To Judge. Not Only Is The Human Being Thus Habitual, On Deleuze’s Reading, But Also Creative Even In The Most Mundane Moments Of Life. Finally, Deleuze Insists That One Of Hume’s Greatest Contributions To Modern Philosophy Is His Insistence That All Relations Are External To Their Terms: This Is The Essence Of Hume’s Anti-Transcendental Stance. Human Nature Cannot Unite Itself, There Is No ‘I’ Which Stands Before Experience, But Only Moments Of Experience Themselves, Unattached And Meaningless Without Any Necessary Relation To Each Other. A Flash Of Red, A Movement, A Gust Of Wind, Dashing Of A Woman Against You, Obscene Gestures, Spitting At You, Talking Extempore About You (My Addition) And These Elements Must Be Externally Related To Each Other To Create The Sensation Of A Tree In Autumn. In The Social World, This Externality Attests To The Always-Already Interested Nature Of Life: No Relation Is Necessary, Or Governed By Neutral Laws, So Every Relation Has A Localised And Passional Motive. The Ways In Which Habits Are Formed Attests To The Desires At The Heart Of Our Social Milieu. Subjectivity, As Deleuze Describes It Through His Reading Of Hume, Is A Practical, Passional, And Empiricist Concept, Immediately Located At The Heart Of The Conventional, Which Is To Say The Social. (Stanford Encyclopedia of Philosophy). The Buddha Always Used The Terms Void, No Rising And Falling, Calmness And Extinction To Explain The Profound Meaning Of Sunyata And Cessation. For Example, Sunyata And The State Of Nirvana Where There Is No Rising Nor Falling, Are Interpreted By Most People As A State Of Non-Existence And Gloom. They

Fail To Realise That Quite The Opposite, Sunyata Is Of Substantial And Positive Significance. The Sutras Often Use The Word "Great Void" To Explain The Significance Of Sunyata. In General, We Understand The "Great Void" As Something That Contains Absolutely Nothing. However, From A Buddhist Perspective, The Nature Of The "Great Void" Implies Something Which Does Not Obstruct Other Things, In Which All Matters Perform Their Own Functions. Materials Are Form, Which By Their Nature, Imply Obstruction. The Special Characteristic Of The "Great Void" Is Non-Obstruction. The "Great Void" Therefore, Does Not Serve As An Obstacle To Them. Since The "Great Void" Exhibits No Obstructive Tendencies, It Serves As The Foundation For Matter To Function. In Other Words, If There Was Neither "Great Void" Nor Characteristic Of Non-Obstruction, It Would Be Impossible For The Material World To Exist And Function. The "Great Void" Is Not Separated From The Material World. The Latter Depends On The Former. We Can State That the Profound Significance of Sunyata and the Nature of Sunyata in Buddhism Highlight The "Great Void's" Non-Obstructive Nature. Sunyata Does Not Imply The "Great Void". Instead, It Is The Foundation Of All Phenomena (Form And Mind). It Is The True Nature Of All Phenomena, And It Is The Basic Principle Of All Existence. In Other Words, If The Universe's Existence Was Not Empty Nor Impermanent, Then All Resulting Phenomena Could Not Have Arisen Due To The Co-Existence Of Various Causes And There Would Be No Rising Nor Falling. The Nature Of Sunyata Is Of Positive Significance. Calmness And Extinction Are The Opposite Of Rising And Falling. They Are Another Way To Express That There Is No Rising And Falling. Rising And Falling Are The Common Characteristics Of Worldly Existence. All Phenomena Are Always In The Cycle Of Rising And Falling. However, Most People Concentrate On Living (Rising). They Think That The Universe And Life Are The Reality Of A Continuous Existence. Buddhism On The Other Hand, Promotes The Value Of A Continuous Cessation (Falling). This Cessation Does Not Imply That It Ceases To Exist Altogether. Instead, It Is Just A State In The Continuous Process Of Phenomena. In This Material World, Or What We May Call This "State Of Existence", Everything Eventually Ceases To Exist. Cessation Is Definitely The Home Of All Existences. Since Cessation Is The Calm State Of Existence And The Eventual Refuge Of All Phenomena, It Is Also The Foundation For All Activities And Functions. Teachings In Chinese Buddhism (6) Sunyata (Emptiness) In The Mahayana Context (Wikipedia) Studies by Eric V. Linder about geometric dark energy bring out the quintessential prerequisites about the extreme physics ad dark matter. The acceleration of the expansion of the universe arises from unknown physical processes involving either new fields in high energy physics or modifications of gravitation theory. It is crucial for our understanding to characterize the properties of the dark energy or gravity through cosmological observations and compare and distinguish between them. In fact, close consistencies exist between a dark energy equation of state function $w(z)$ and changes to the framework of the Friedmann cosmological equations as well as direct spacetime geometry quantities involving the acceleration, such as "geometric dark energy" from the Ricci scalar. We investigate these interrelationships, including for the case of super acceleration or phantom energy where the fate of the universe may be gentler than the Big Rip. DOI: <http://dx.doi.org/10.1103/PhysRevD.70.023511>

Probing gravitation, dark energy, and acceleration Phys. Rev. D 70, 023511 – Published 28 July 2004 Eric V. Linder. In the same vein he continues to question the number of parameters that exists in dark energy .How many dark energy parameters? For exploring the physics behind the accelerating Universe a crucial question is how much we can learn about the dynamics through next generation cosmological experiments. For example, in defining the dark energy behavior through an effective equation of state, how many parameters can we realistically expect to tightly constrain? Through both general and specific examples (including new parametrizations and principal component analysis) we argue that the answer is two. Cosmological parameter analyses involving a measure of the equation of state value at some epoch (e.g. w_0) and a measure of the change in equation of state (e.g. w') are therefore realistic in projecting dark energy parameter constraints. More elaborate parametrizations could have some uses (e.g. testing for bias or comparison with model features), but do not lead to accurately measured dark energy parameters. DOI: <http://dx.doi.org/10.1103/PhysRevD.72.043509> **Phys. Rev. D 72, 043509 – Published 11 August 2005 Eric V. Linder and Dragan Huterer D.F. Mota, C. van de Bruck Oxford Univ** study the spherical collapse model in dark energy cosmologies, in which dark energy is modelled as a minimally coupled scalar field. We first follow the standard assumption that dark energy does not cluster on the scales of interest. Investigating four different popular potentials in detail, we show that the predictions of the spherical collapse model depend on the potential

used. We also investigate the dependence on the initial conditions. Secondly, they investigate in how far perturbations in the quintessence field affect the predictions of the spherical collapse model. In doing so, we assume that the field collapses along with the dark matter. Although the field is still subdominant at the time of virialisation, the predictions are different from the case of a homogeneous dark energy component. This will in particular be true if the field is non--minimally coupled. **D.F. Mota, C. van de Bruck** conclude that a better understanding of the evolution of dark energy in the highly non--linear regime is needed in order to make predictions using the spherical collapse model in models with dark energy. *Astrophys J* 421 (2004) 71-81 DOI:10.1051/0004-6361:20041090arXiv:astro-ph/0401504 **on the spherical collapse model in dark energy cosmologies D.F. Mota, C. van de Bruck (Oxford Univ)** Studies by Eric V. Linder about geometric dark energy bring out the quintessential prerequisites about the extreme physics ad dark matter. The acceleration of the expansion of the universe arises from unknown physical processes involving either new fields in high energy physics or modifications of gravitation theory. It is crucial for our understanding to characterize the properties of the dark energy or gravity through cosmological observations and compare and distinguish between them. In fact, close consistencies exist between a dark energy equation of state function $w(z)$ and changes to the framework of the Friedmann cosmological equations as well as direct spacetime geometry quantities involving the acceleration, such as "geometric dark energy" from the Ricci scalar. We investigate these interrelationships, including for the case of super acceleration or phantom energy where the fate of the universe may be gentler than the Big Rip. DOI: <http://dx.doi.org/10.1103/PhysRevD.70.023511> **Probing gravitation, dark energy, and acceleration Phys. Rev. D 70, 023511 – Published 28 July 2004 Eric V. Linder. In the same vein he continues to question the number of parameters that exists in dark energy .**How many dark energy parameters? For exploring the physics behind the accelerating Universe a crucial question is how much we can learn about the dynamics through next generation cosmological experiments. For example, in defining the dark energy behavior through an effective equation of state, how many parameters can we realistically expect to tightly constrain? Through both general and specific examples (including new parametrizations and principal component analysis) we argue that the answer is two. **All Existences Exhibit Void-Nature And Nirvana-Nature. These Natures Are The Reality Of All Existence. To Realise The Truth, We Have To Contemplate And Observe Our Worldly Existence. We Cannot Realise The Former Without Observing The Latter. Consider This Heart Sutra Extract, "Only When Avalokiteshvara Bodhisattva Practised The Deep Course Of Wisdom Of Prajna Paramita Did He Come To Realise That The Five Skandhas (Aggregates, And Material And Mental Objects) Were Void." Profound Wisdom Leads Us To The Realisation That All Existences Are Of Void-Nature. The Sutras Demonstrate That The Profound Principle Can Be Understood By Contemplating And Observing The Five Skandhas. We Cannot Realise The Truth By Seeking Something Beyond The Material And Mental World. The Buddha, Using His Perfect Wisdom, Observed Worldly Existence From Various Implications And Aspects, And Came To Understand All Existences. Teachings In Chinese Buddhism (6) Sunyata (Emptiness) In The Mahayana Context (Wikipedia) .But From The Profound Contemplation And Wisdom Of The Buddha And Mahabodhisattvas, We Know There Is No Such Reality. Instead, Egolessness (Non-Self) Is The Only Path To Understand The Reality Of The Deluded Life. All Existences Are Subject To The Law Of Causes And Conditions. These Include The Smallest Particles, The Relationship Between The Particles, The Planets, And The Relationship Between Them, Up To And Including The Whole Universe! From The Smallest Particles To The Biggest Matter, There Exists No Absolute Independent Identity. Egolessness (Non-Self) Implies The Void Characteristics Of All Existence. Egolessness (Non-Self) Signifies The Non-Existence Of Permanent Identity For Self And Existence (Dharma). Sunyata Stresses The Voidness Characteristic Of Self And Existence (Dharma). Sunyata And Egolessness Possess Similar Attributes. As We Have Discussed Before, We Can Observe The Profound Significance Of Sunyata From The Perspective Of Inter-Dependent Relationships. Considering Dharma-Nature And The Condition Of Nirvana, All Existences Are Immaterial And Of A Void-Nature. Then We See Each Existence As Independent Of Each Other. But Then We Cannot Find Any Material That Does Exist Independent Of Everything Else. So Egolessness Also Implies Void-Nature! From The Observation Of All Existences, We Can Infer The **Theory Of Nirvana And The Complete Cessation Of All Phenomena. From The Viewpoint Of Phenomena, All Existences Are So Different From Each Other, That They May Contradict Each Other. They Are So Chaotic. In Reality, Their Existence Is Illusionary And****

Arises From Conditional Causation. They Seem To Exist On One Hand, And Yet Do Not Exist On The Other. They Seem To Be United, But Yet They Are So Different To One Another. They Seem To Exist And Yet They Do Cease! Ultimately Everything Will Return To Harmony And Complete Calmness. This Is The Nature Of All Existence. It Is The Final Resting Place For All. If We Can Understand This Reality And Remove Our Illusions, We Can Find This State Of Harmony And Complete Calmness. Teachings In Chinese Buddhism (6) Sunyata (Emptiness) In The Mahayana Context (Wikipedia) All Our Contradictions, Impediments And Confusion Will Be Converted To Equanimity. Free From Illusion, Complete Calmness Will Be The Result Of Attaining Nirvana. The Buddha Emphasised The Significance Of This Attainment And Encouraged The Direct And Profound Contemplation On Void-Nature. He Said, "Since There Is No Absolute Self-Nature Thus Every Existence Exhibits Void-Nature. Because It Is Void, There Is No Rising Nor Falling. Since There Is No Rising Nor Falling, Thus Everything Was Originally In Complete Calmness. Its Self-Nature Is Nirvana." Teachings In Chinese Buddhism (6) Sunyata (Emptiness) In The Mahayana Context (Wikipedia). The Scholars Of Tien Tai Called It The "Embodied Nature". (This Is The Buddha-Nature That Includes Both Good And Evil.) The Scholars Of Xian Shou Say, "It Is Arising From Primal Nature", And The Scholars Of Chan (Zen) Say, "It Is Nature That Causes The Rising Of Things". All Dharma Is Dharma-Nature. It Is Not Different From Dharma-Nature. Dharma And Dharma-Nature Are Not Two Separate Identities, "Phenomena" And "Nature" Is Also Not Distinguishable Either. In Other Words, There Is No Difference Between Principle (Absolute) And Practice (Relative). Teachings In Chinese Buddhism (6) Sunyata (Emptiness) In The Mahayana Context (Wikipedia). In Other Words, All Dharma Arises From Causes And Conditions, And All Dharma Is Empty In Nature. The Law Of Dependent Origination (Existence) And The Nature Of Emptiness Is Neither The Same Nor Different. They Exist Mutually. The Truth Of "Sunyata" And "Existence", And "Nature" And "Phenomena" Are Not In Conflict With Each Other. Unlike The Scholars Of The Dharmalakshana Sect Who Explain The Dharma Only From The Aspect Of Dependent Origination, Or The Scholars Of Dharma-Nature That Explain The Existence Of Dharma Only From The Aspect Of Dharma-Nature, The Scholars Of Madhyamika Explain The Truth Of The Dharma From Both Aspects. Hence This Is Called The Middle Path Which Does Not Incline To Either Side. Teachings In Chinese Buddhism (6) Sunyata (Emptiness) In The Mahayana Context (Wikipedia) **C S Lent, P D Tougaw, W Porod and G H Bernstein** formulate a new paradigm for computing with cellular automata (CAS) composed of arrays of quantum devices-quantum cellular automata. Computing in such a paradigm is edge driven. Input, output, and power are delivered at the edge of the CA array only; no direct flow of information or energy to internal cells is required. Computing in this paradigm is also computing with the ground state. The architecture is so designed that the ground-state configuration of the array, subject to boundary conditions determined by the input, yields the computational result. proposal is put forth for a specific realization of these ideas using two-electron cells composed of quantum dots. The charge density in the cell is very highly polarized (aligned) along one of the two cell axes, suggestive of a two-state CA. The polarization of one cell induces a polarization in a neighboring cell through the Coulomb interaction in a very non-linear fashion. Quantum cellular automata can perform useful computing. The authors show that AND gates, OR gates and inverters can be constructed and interconnected. **C S Lent et al 1993 Nanotechnology 4 49 doi:10.1088/0957-4484/4/1/004 Quantum cellular automata C S Lent, P D Tougaw, W Porod and G H Bernstein** The basic features of the dynamics of open quantum systems, such as the dissipation of energy, the decay of coherences, the relaxation to equilibrium or non-equilibrium stationary state, and the transport of excitations in complex structures are of central importance in many applications of quantum mechanics. The theoretical description, analysis and control of non-Markovian quantum processes play an important role in this context. While in a Markovian process an open system irretrievably loses information to its surroundings, non-Markovian processes feature a flow of information from the environment back to the open system, which implies the presence of memory effects and represents the key property of non-Markovian quantum behaviour. **Heinz-Peter Breuer** review recent ideas developing a general mathematical definition for non-Markovianity in the quantum regime and a measure for the degree of memory effects in the dynamics of open systems, which are based on the exchange of information between system and environment. **Heinz-Peter Breuer** further study the dynamical effects induced by the presence of system-environment correlations in the total initial state and design suitable methods to detect such correlations through local measurements on the open system. **Heinz-Peter Breuer 2012 J. Phys. B: At. Mol. Opt. Phys. 45**

154001 doi:10.1088/0953-4075/45/15/154001 Foundations and measures of quantum non-Markovianity A fundamental step towards atomic- or molecular-scale spintronic devices has recently been made by demonstrating that the spin of an individual atom deposited on a surface¹, or of a small paramagnetic molecule embedded in a nanojunction², can be externally controlled. An appealing next step is the extension of such a capability to the field of information storage, by taking advantage of the magnetic bistability and rich quantum behaviour of single-molecule magnets^{3, 4, 5, 6} (SMMs). Recently, a proof of concept that the magnetic memory effect is retained when SMMs are chemically anchored to a metallic surface⁷ was provided. However, control of the nanoscale organization of these complex systems is required for SMMs to be integrated into molecular spintronic devices^{8, 9}. Here we show that a preferential orientation of Fe⁴ complexes on a gold surface can be achieved by chemical tailoring. As a result, the most striking quantum feature of SMMs—their stepped hysteresis loop, which results from resonant quantum tunnelling of the magnetization^{5, 6}—can be clearly detected using synchrotron-based spectroscopic techniques. With the aid of multiple theoretical approaches, **M. Mannini et al** relate the angular dependence of the quantum tunnelling resonances to the adsorption geometry, and demonstrate that molecules predominantly lie with their easy axes close to the surface normal. Findings of **M. Mannini et al** prove that the quantum spin dynamics can be observed in SMMs chemically grafted to surfaces, and offer a tool to reveal the organization of matter at the nanoscale. **Quantum tunnelling of the magnetization in a monolayer of oriented single-molecule magnets M. Mannini et al** Nature 468, 417–421 (18 November 2010) doi: 10.1038/nature09478 Supposing That There Were Indeed An "Energy Of Contraction" Constant In All Centers Of Force Of The Universe, It Remains To Be Explained Where Any Difference Would Ever Originate. It Would Be Necessary For The Whole To Dissolve Into An Infinite Number Of Perfectly Identical Existential Rings And Spheres, And We Would Therefore Behold Innumerable And Perfectly Identical Worlds COEXISTING [Nietzsche Underlines This Word Twice] Alongside Each Other. Is It Necessary For Me To Admit This? Is It Necessary To Posit An Eternal Coexistence On Top Of The Eternal Succession Of Identical Worlds? The Eternal Return: Genesis And Interpretation BY PAUL D'ITORIO "The Cyclical Hypothesis, So Heavily Criticized By Nietzsche (VP II 325 And 334), Arises In This Way." In Fact, Nietzsche Was Not Criticizing The Cyclical Hypothesis But Only The Particular Form Of That Hypothesis Presented In Vogt's Work. All Of **Nietzsche's Texts Without Exception Speak Of The Eternal Return As The Repetition Of The Same Events Within A Cycle Which Repeats Itself Eternally. If Deleuze's Interpretation Holds That The Eternal Return Is Not A Circle, Then What Is It? A Wheel Moving Centrifugally, Operating A "Creative Selection," "Nietzsche's Secret Is That The Eternal Return Is Selective"** Says Deleuze: The Eternal Return Produces Becoming-Active. It Is Sufficient To Relate The Will To Nothingness To The Eternal Return In Order To Realize That Reactive Forces Do Not Return. However Far They Go, However Deep The Becoming-Reactive Of Forces, Reactive Forces Will Not Return. The Small, Petty, Reactive Man Will Not Return. Affirmation Alone Returns, This That Can Be Affirmed Alone Returns, Joy Alone Returns. Everything That Can Be Denied, Everything That Is Negation, Is Expelled Due To The Very Movement Of The Eternal Return. We Were Entitled To Dread That The Combinations Of Nihilism And Reactivity Would Eternally Return Too. The Eternal Return Must Be Compared To A Wheel; Yet, The Movement Of The Wheel Is Endowed With Centrifugal Powers That Drive Away The Entire Negative. Because Being Imposes Itself On Becoming, It Expels From Itself Everything That Contradicts Affirmation, All Forms Of Nihilism And Reactivity: Bad Conscience, Ressentiment..., We Shall Witness Them Only Once. [...] The Eternal Return Is The Repetition, But The Repetition That Selects, The Repetition That Saves. Here Is The Marvelous Secret Of A Selective And Liberating Repetition. There Is No Need To Remind The Reader That Neither The Image Of A Centrifugal Movement Nor The Concept Of A Negativity-Rejecting Repetition Appears Anywhere In Nietzsche's Writings, And Indeed Deleuze Does Not Refer To Any Text In Support Of This Interpretation. Further, One Could Highlight That Nietzsche Never Formulates The Opposition Between Active And Reactive Forces, Which Constitutes The Broader Framework Of Deleuze's Interpretation. For Some Years, Marco Brusotti Has Called Attention To The Fact That Deleuze Introduced A Dualism That Does Not Exist In Nietzsche's Writings. To Be Sure, The German Philosopher Describes A Certain Number Of "Reactive" Phenomena (For Example, In The Second Essay Of The Genealogy Of Morality, § 11, He Talks About "Reactive Affects" [Reaktive Affekte], "Reactive Feelings" [Reaktive Gefühlen], Reactive Men [Reaktive Menschen]); But These Are Nonetheless The Result Of Complex Ensembles Of Configurations Of Centers Of Forces That Remain In

Themselves Active. Neither The Word Nor The Concept Of "Reactive Forces" Ever Appears In Nietzsche's Philosophy. The Eternal Return: Genesis And Interpretation BY PAUL D'ORIO Some Parts Are Deleted Due To Spatial Constraints. Please Pardon Me For The Deletion. Deleuze Opposes The Historical Course Of The Hegelian Notion That Confronts Struggles And Finally Dialectizes The Negative And Results In A Consoling Teleology Leading To The Triumph Of The Idea Or The Liberation Of The Masses With The Centrifugal Movement Of The Wheel, Which Simply Ejects The Negative. It Is Still A Case Of A Consoling And Optimistic Teleology, Which, Instead Of Confronting The Weight Of History, The Grief And The Negative, Makes It Disappear In One Centrifugal Stroke Of A Magic Wand. There Is Reason To Worry That This Be A Case Of Repression, Which, Unable To Dialectize Or Accept The Negative, Simply Seeks To Exorcise It In One Gesture Of "Creative Selection." But Exorcism Is A Feat Of Magic And Not Of Philosophy: It Is Unfortunately Not Enough To Make The Negative Disappear. In All Probability, The Negative Will Come Back With A Vengeance. In Contrast To Deleuze's "Affirmation Of Affirmation", Which Affirms Only Affirmation, Nietzsche Conceives Of The Eternal Return From A Rigorously Non-Teleological Perspective As The Accomplishment Of A Philosophy Strong Enough To Accept Existence In All Its Aspects, Even The Most Negative, Without Any Need To Dialecticize Them, Without Any Need To Exclude Them By Way Of Some Centrifugal Movement Of Repression. It Denies Nothing And Incarnates Itself In A Figure Similar To The One Nietzsche, In Twilight Of The Idols, Draws Of Goethe: Such A Spirit, Who Has Become Free Stands In The Middle Of The World With A Cheerful And Trusting Fatalism In The Belief That Only The Individual Is Reprehensible, That Everything Is Redeemed And Affirmed In The Whole-He Does Not Negate Anymore. Such A Faith However, Is The Highest Of All Possible Faiths: I Have Baptized It With The Name Of Dionysus. The Eternal Return: Genesis and Interpretation BY PAUL D'ORIO The Most Complete Theory Of Mind That Comports With The Holographic Principle Of Mind, As Developed Here, Is The Microgenetic Theory Of Neurologist Jason Brown. Brown (2005) Presents An Evolutionary Theory Of Values, Morals, And Ethics, Building On The Microgenetic Theory And Process Theory Of His Earlier Works (1988, 1991, 1996, 1998, And 2000). Brown Is Work Has Been A Progression That Began With The Study Of Brain Processes In Neurological Subjects With Brain Injuries Or Lesions. In His Earlier Work (Brown, 1988; 1991) He Described The Process Of Microgenesis As A Process Of Elaboration Of Mental Contents In An Evolutionary And Developmental Hierarchy Of Brain Structures Within The Process Of The Genesis Of The Mental State, And Related The Hierarchy Of The Genesis Of The Mental State To Disturbances Of Language Comprehension Or Expression (Aphasia), Of Knowledge (Agnosia), And Or Purposeful Movement (Apraxia). The Fundamental Neurology Of These Disturbances Had Been Described In Brown's Previous Work (1972, 1988), And In The Seminal Work Of The Russian Neurologist And Neuropsychologist, Alexander Luria (1966). Brown (1988, 1991, 1996, 1998) Created A Process Theory Of Mind On The Basis Of His Neurological Observations, And Went On From His Neurological Work To Incorporate Process Theory In Philosophy, As Developed Principally By Henri Bergson (1911/1998). And Alfred North Whitehead (1925/1953, 1929/1978). Bergson Had Pioneered The Concept Of An Irreducible Duration Of Experience. This Concept Was Elaborated By Whitehead, In Keeping With Discontinuities Of Particles, As "Actual Entities" Of A Rudimentary Sort, As They Make "Quantum Leaps" Along Their Trajectories, Halting Or Persisting At A Given Location For A Short Period Of Time Or Duration Before Leaping To The Next. Whitehead Employed These "Quantum Leaps" To The Process By Which Experience Arises In "Actual Entities" At The Most Fundamental Level Of Actuality, Through The Epochal Or Halting Duration Of A Discontinuous Process (Whitehead, 1925/1953, 1929/1978). This Halting Allows Actual Entities To Participate In A Process Of Internal Relations, Relations On The "Inside" Of Things. According The Holographic Principle, This "Inside" Is In The Holographic Surface Of The Relevant Region Of Spacetime Of The Actual Entity, And Within Other Surfaces That Are Non-Locally Connected With That Surface. The Holographic Surface Is Thus The Locus Of Non-Local Information And Experience, In Our Interpretations Of Whitehead Is Metaphysics. Internal Relations Occur Through Feeling, Or "Prehension," Which Is Nothing More Than The Subjective Quality Of Non-Local Experience. The Internal Relations Of The Becoming Entity Occur In A Seriality Of Prior Becomings Of The Actual Entity, With The Entity Inheriting All Of Its Causal Past As Well As Its Relevant Feelings Or Prehensions From Prior "Occasions" Or Quanta Of Experience. This Process Is, In The Holographic Model, The Process By Which Former Surfaces, Now-Pasts, Are Elaborated On To The Now-Present. Prehensions Also Occur With Other Actual Entities

Throughout The Universe, With Which They Are Non-Locally Connected, And With Neo-Platonic "Eternal Objects," Existing In An Eternal Heaven. The "Ingression" Of Eternal Objects Is Fundamental To Each Duration Of Becoming Of The Entity, And, On This Basis, Whitehead Formulated The Notion Of An Eternal Heaven In Constant Intercourse All Of Its Creation. There Are Clear Parallels Between Whitehead's Cosmology And The Cosmology Introduced Here According To The Holographic Principle, In That The "Heaven" May Be Regarded As The Holographic Boundary, In Constant Intercourse With The Relevant Volume Of Spacetime. This Interconnection Of Feelings Or Prehensions, According To Their Relevance, Produces Intensities And Contrasts Which Give Rise To The Creative Advance, Which, For Whitehead (1929/1978) Is The Process Which Impels The Movement From The Physical To The Mental Pole Of Process. With Each Halting, Internally-Timeless Duration There Is A Concrecence Or Integration Of These Feelings, Which, When Complete, Constitutes A New Actual Entity Or Actual Occasion (They Are The Same Thing) At A New Locus In Spacetime. This Movement To A New Locus, Or Transition, Is, Fundamentally, The Quantum Leap. MICROGENESIS AND THE HOLOGRAPHIC PRINCIPLE

The Holographic Principle Theory Of Mind MARK GERMINE Institute For Psycho Science If the Holographic Principle is true, than it must be the fundamental principle of mind. The brain has no way around the Holographic Principle. For the reductionist, the Holographic Principle is the ultimate reduction. It applies to the most minuscule level of what we can observe, and beyond. For the Universalist, the Holographic Principle gives us the ultimate universal. It extends to the limit of our Universe, the Universal holographic boundary, and beyond. For the Phenomenalist, the Holographic Principle gives us the ultimate ground of our phenomenal perception, our grounding in the Universal "now," in the now-present of the Universal holographic boundary, moving outward from now-pasts to now-futures. Wheeler (1988) has said that all of reality is information, and that other physical quantities are "mere incidentals." Information monism is gaining popularity in physics. By breaking the dichotomy of between information and experience, we find a deep connection between Wheeler is monism and the experiential monism of Whitehead, sometimes called panexperientialism, which relates to the later theological concept of his student, Charles Hartshorne, panentheism, God inside of everything (Hartshorne, 1964). There thus seems to be a convergence of the concepts of information, experience, and spirituality. There is not one universe, but many parallel universes, or, more accurately, a vast superposition of universes called the multiverse (Penrose, 2004). However, as explained in our treatment of the double-slit experiment, in the observation of events on the quantum level, the individual observer sees only one of these vast superpositions, not a summation of a vast number of potentials. This has been explained in terms of the many worlds or many minds theories, in which there are multiple copies of the same observing individual, but this particular issue remains unresolved, leaving physics ungrounded in reality (Penrose, 2004). The concept of One Mind is not only more parsimonious than that of many minds, but also lets us out of the bizarre and counterintuitive idea that there are multiple copies of our own selves, which exist as mere potential, and are thus not actual. There are many possible universes, but there is only One Mind, which determines events on the quantum level, and thus creates our Universe. As we had discussed previously, the quantum level provides the essential ground of the Holographic Principle, such that quantum-level holographic surfaces are elaborated at higher levels, manifesting higher orders of information from the quantum "world." In this process, there is a reduction of the wave function or potentialities of quantum fields. At the level of consciousness, this entails freedom to choose which observations we make (Stapp, 1997), which gives us the capacity to think and to make decisions. These capacities are the basis of individuality, self-determination, judgment, and values. The progressive evolution of the manifestation of Mind through higher orders of experience, leading to higher orders of consciousness, is entailed by the Holographic Principle. Again, we are dealing here with levels of description, with the multiverse of all potentials fundamentally supporting the single Universe we collectively observe. The multiverse is the wave function of the Universe (Penrose, 2004). The recursive integration of nested hierarchies of holographic surfaces brings out a single actuality in consciousness from a wave function that is unconscious. Consciousness thus gives us information at a level of experience that is causal. In this sense, we partake in the creation of the Universe. We participate in creation, and this participation, when fully realized, leads us to higher levels of consciousness and of realization. As a system, the biosphere that we live in has a holographic surface, creating a deep sense of ecology as we collectively move toward a planetary consciousness. It is only when the Universal Holographic Boundary reaches the information storage and processing capacity needed for the requisite

biological and biochemical complexity that consciousness evolves in living things. This evolution is natural and spontaneous, since consciousness gives rise to what is actual, as opposed to what is merely potential. In this sense, consciousness is still in the process of creating our Universe, and levels of higher consciousness will continue to evolve. We are all in the same "now," and that now is defined by the present Universal Boundary. This assures us that our experience is Universal, and does not pass with time. This is the fundamental basis of memory and of cognition. The identity of mind and brain is a myth. We have a continual, internal or non-local relation with the Universe, as it has with us. Once this mystery is resolved, the myth is no longer needed, and there is a confluence of science and spirituality. Experience is primary, information is secondary. We can only gather information from experience, whether it is in the laboratory or in life. We cannot measure the information on the surfaces of systems. The physicists that have formulated the Holographic Principle for all systems are quite aware of this, or else the principle would have been established or discredited. We cannot measure what we experience. It is intangible, yet it is all we know to be actual. Everything else is inferred. Because it cannot be measured, it has been fundamentally disregarded by mainstream science. Materialism is considered scientific, while idealism is considered unscientific. But aren't ideas, fundamentally, made of information? Brain science has mistaken the representation of information for information itself, and has tied those representations to matter and energy. Consciousness, the highest order of information, has generally been regarded as superfluous, something that needs to be explained away, or altogether ignored. Yet it is the only "thing" that reaches our awareness. Consciousness comes at a price. For everything that becomes conscious, there must be something that becomes conscious. Consciousness is certainty, and its complement, unconsciousness uncertainty. Consciousness is the particle nature of experience. It has definiteness about it. The unconscious is the wave nature of experience, it is like the metaphysical cloud of unknowing. If we are the most conscious of animals, then we must also be the most unconscious. Perhaps this is the predicament of humankind. The Holographic Principle of Mind leads us naturally from the most fundamental experiences, existing as quantum potentials from the conformations of proteins down through the fields of electrons, through their manifestation upward through a recursion or successive applications of the same holographic process, through higher levels of experience, to the emergence of consciousness as higher and higher orders of experience. The quantum Holographic Principle Mind does not require anything more quantum than is obviously present at the microscopic and submicroscopic levels, as it represents successive orders of manifestations from these levels. Recursion also applies here in the sense that it is used in computer science, in which the function of the part depends on the function of the whole. A program, as a part, cannot work without a functioning whole, the operating system. Recursive wholes which are, for us, supra-conscious, are on the group, species, planetary, and universal levels. As individuals, our consciousness cannot function without recursion to the Universal Consciousness, even though we may be unaware that such Universal Consciousness exists. Reaching upward to these supra-conscious levels is a spiritual process, making transcendence the ultimate solution to the unconscious human predicament. What was once supra-conscious becomes unconscious through a process of conditioning? We are born as creatures of the Earth and of the Universe, as evidenced by the beliefs and practices of "primitive peoples." There is evidence from cave paintings that our hominid ancestors experienced a kind of holographic perception (Combs, 1996), which could constitute our early connection with the holographic subtext of reality, and which we might then propose existed in our animal ancestors, and in extant animal species. If this is the case, then microgenesis would entail the recapitulation of this holographic experience as it progresses through our ancestral past. The supra-conscious mind seems to envelope perinatal experiences, and Stanislov Grof (1994) has developed techniques to gain access to these experiences, as well as to the earlier experiences of our human and animal ancestral lineage, and of our universal history. Grof (1994) concludes: "Our consciousness seems to have the amazing capacity to directly access the earliest history of the universe and witnessing dramatic sequences of the Big Bang, the formation of the galaxies, the birth of the solar system, and the early geophysical processes on this planet billions of years ago." The Holographic Principle Theory Of Mind MARK GERMINE Institute For Psycho Science. There Are Similarities Between The Patterns Of Holography And Of Psychological Transference, Where Holography Is The Process Of Recording And Reconstructing Holograms. Employing A Theoretical Perspective Using A Hermeneutic Method, This Dissertation Parallels Holography With Transference, Offering Another Way To Encounter Transference By Showing Similarities Between The Processes Of Each And The Results Of Each. Though Complex, Infinitely Varying, And

Unique, Their Patterns Are Clearly Identifiable. Thus They Are A Metaphorical Fit To The Concept Of Strange Attractors In Physics And A More Literal Fit To The Concept Of Archetypes In Depth Psychology Or Dynamic Psychology, Psychology Which Attends To The Living, Autonomous Unconscious. This Study Explains How Holography Models Transference, What A Hologram Is And How It Works, And How Depth Psychology Understands Of The Interaction Between Consciousness And The Unconscious Is Related To The Hologram. It Describes Transference And Related Psychological Processes As Understood In Six Different Schools Of Depth Psychological Thought. It Shows That The Underlying Pattern Or Strange Attraction Between Transference And Holography Extends To Other Processes Both Within And Outside The Field Of Psychology, Processes Such As Projection, Projective Identification, Splitting, Memory, Biology, Creative Discovery, Theology, Synchronicity, Chaos, And Nonlocality. By Identifying The Similar Patterns Of These Processes, This Study Demonstrates The Existence Of An Underlying Holographic Archetype In Which Essential Qualities Of The Whole Are Present In Each Of The Parts Of The Whole: The Visual Image Of The Overall Hologram Is Present In Each Component Part Of The Hologram, The Autonomy Of The Overall Human Is Present In Each Conscious And Unconscious Component Part Of The Human Psyche. By Noting Differences As Well As Similarities In These Processes, It Suggests An Inventory Of The Qualities Of The Holographic Archetype. This Study Furthers Understanding Of The Pervasiveness, Force, And Autonomy Of The Unconscious Acting Through Transference And Projection By Identifying A Group Projection Of Domestic Violence Lying At The Core Of The Christian Myth. This Study Also Furthers Understanding Of The Concept Of Transference By Providing A Reflection Hologram Of The Human Psyche As An Artistic Work And As A Visual Metaphor Of Transference. Strange Attractors: Transference, Holography, And An Archetype Burke, J. (2003). Strange Attractors: Transference, Holography, And An Archetype (Doctoral Dissertation, Pacifica Graduate Institute, 2003). Onticology, Like All Variations Of Object-Oriented Ontology, Is Realist In Its Orientation. In Defending A Realist Ontology Onticology Holds That The Vast Majorities Of Objects, Actants, Beings, Or Entities Are Independent Of Humans And Are What They Are Regardless Of Whether Any Humans Regard Them Or Register Them. In Short, Onticology Rejects Any Anthropomorphic, Idealist, Or Anti-Realist Thesis To The Effect That To Be Is To Be The Correlate Of Mind, Spirit, The Body, The Human, And Language Or Otherwise. While It Is Certainly The Case That Knowledge Is Necessarily Dependent On The Object To Which It Relates, The Converse Does Not Hold True. Objects Are Not Dependent On Being Known, Regarded, Perceived, Or Spoken About. As Such, And To Put It In Aristotelean Terms, Knowledge Is An Accident Of Objects, Not Objects An Accident Of Knowledge. As Althusser So Nicely Puts It, “[N]O Doubt There Is A Relation Between Thought-About-The-Real And This Real, But It Is A Relation Of Knowledge, A Relation Of Adequacy Or Inadequacy Of Knowledge, Not A Real Relation, Meaning By This A Relation Inscribed In That Real Of Which The Thought Is The (Adequate Or Inadequate) Knowledge” (Reading Capital, 96). Althusser Goes On To Remark That “[T]he Distinction Between A Relation Of Knowledge And A Relation Of The Real Is A Fundamental One: If We Did Not Respect It We Should Fall Irreversibly Into Either Speculative Or Empiricist Idealism” (Ibid.). Onticology Categorically Endorses Althusser’s Verdict It Is A Fundamental Necessity To Distinguish Between Those Relations That Belong To The Object And Those That Belong To Knowledge. Contemporary Philosophy Continuously Confuses These Two Very Different Sorts Of Relations. Naturally The Question Arises Of How It Is Possible To Surmount Our Relation To The Object So As To Determine Whether Objects Themselves Possess The Properties We Encounter In Relating To Objects. In Other Words, Given That We Can Only Ever Relate To The Object In Relating To The Object How Is It Possible To Surmount This Relation To Get At The Being Of The Object Itself? Much More Will Have To Be Said About This Later– And The Answers Will Be Surprising With Respect To Standard Prejudices About Realism –However, For The Moment It Can Be Said That Onticology Takes Its Epistemological Inspiration From The Transcendental Realism Of Roy Bhaskar. Among Other Things, Bhaskar Sought To Provide A Transcendental Grounding For The Sciences. Insofar As Onticology Defends The Thesis That The Field Of Being Is Much Vaster Than The Field Of Objects Investigated By The Natural Sciences, It Parts Way With The Thesis That The Domain Of Being Is Exhausted By The Domain Of Natural Objects. However, The General Form Of Bhaskar’s Argument Holds For Our Realist Purposes. A Transcendental Argument Seeks To Elucidate The Conditions Under Which Certain Acknowledged Practices And Forms Of Cognition Are Possible. Kant, For Example, Asked What Must Be The

Case For Mathematical Judgments To Be Possible. How Is It Both That We Are Able To Extend Our Knowledge, As If By Magic, Through Mathematical Judgments And, More Significantly, That These Judgments Are Able To Provide Genuine Knowledge Of The World Despite The Fact That These Forms Of Reasoning Are Not Based On Experience? Part Of Kant's Argument Consisted In Claiming That Mind Imposes The Forms Of Space And Time On The Data Of Experience. In Other Words, Space And Time Are Not Attributes Of Being Itself But Rather Of The Mind That Regards Being. Insofar, Kant Argues, As Mathematics Is Ultimately A Rumination On The Nature Of Space And Time Taken In Their Most Abstract Form And Insofar As The Mind Imposes Space And Time On The Manifold Of Sensation, It Thus Follows That A Priori Judgments About The Nature Of Spatio-Temporal Relations Are Possible That Anticipate The Structure Of Actual-Space Times Without Directly Experiencing These Space-Times. Why? Because Any Manifold Of Sensation Must Necessarily Be Structured By These Forms Imposed By Intuition Ontology— A Manifesto For Object-Oriented Ontology Part I Posted By Larval Subjects Under Object-Oriented Philosophy. it is necessary to distinguish the being of objects from the manifestation of objects. While objects are acts, these acts are not identical with their performance in either nature (events where no humans are about to perceive them) or with their performance for humans. Rather, the proper being of the object is not its performance or manifestation, but the generative mechanism that serves as the condition under which these performances or manifestations are possible. As Graham Harman will argue— though in a very different theoretical constellation —the being of objects is essentially withdrawn or hidden. No one has ever perceived a single object, but we do perceive all sorts of effects of objects. Traditional epistemology has confused these effects with the objects themselves. Fortunately we do occasionally manage to cognize objects through a sort of detective work that infers these generative mechanisms from their effects; without, for all this, ever exhausting the infinity of a single object. At any rate, if objects were not withdrawn in this way, the practice of experiment would be unintelligible. This leads to Bhaskar's answer to the first question: What must the world be like for science to be possible? Note, this is not a question about mind or culture, but about the world itself, regardless of whether or not humans exist. Once again, knowledge is an accident of objects, not objects an accident of consciousness or cognition. If science is to be possible— and I would argue, if any human practice is to be possible —then the world must be structured and differentiated. The world must have joints or, as Harman puts it, the world must be composed of “chunks”. Why is this case? Let us return to the question of experimentation and the conditions under which experiment is possible. We will adopt two possible hypotheses pertaining to the ontological nature of the world independent of humans: 1) As certain mystics and contemporary crypto-mystics would have it, the world is an undifferentiated One-All that is only subsequently segmented or partitioned into discrete beings by some form of human agency whether this be through cognition or language (in the case of language we might think of Saussure's and Hjelmslev's undifferentiated “sonorous matter”). 2) Entities are the sum totality of their relations to all other entities in the universe. The first hypothesis is easily dispatched on two grounds: First, this hypothesis fails to register the contradiction in its own utterance. At the level of explicit content, it claims that the world is an undifferentiated One-All that is only subsequently segmented into discrete beings, yet what it misses at the unconscious level of its own utterance is that it registers at least one structured differentiation that is not undifferentiated within this One-All: Namely, the agency through which the One-All subsequently comes to be differentiated. Certain anti-realist, transcendental philosophers will, in a gesture that is all too cute, claim that the agency by which the world is segmented cannot properly be said to exist, thus attempting to resolve this contradiction through a sleight of hand. However, as Meillassoux has shown in *After Finitude*, all attempts on the part of transcendental anti-realist philosophies to treat the transcendental subject as a non-existing or non-objectile agency that does not itself exist end up, all too clearly, attaching that conception of finitude and the segmentary work with which it is charged to the body (a differentiated being or generative mechanism). Second, suppose the anti-realist transcendental philosophers were to convince us through his appeals to the non-existence of the transcendental, would we still encounter problems? Like Atlas, transcendental subjectivity is charged with the monumental task of segmenting the formless Apeiron of the One-All from out of primal chaos into a supremely segmented world. But this world appears to us to be too slippery for even a titan like Atlas to grasp. Were the world, prior to and independent of humans genuinely a formless Apeiron it would contain no differences providing hand-holds for Atlas to grasp in his segmenting activity. Consequently, no segments could ever come into being. Yet

everywhere we encounter segments, so the world must not be a formless One-All, but must rather be structured and differentiated even if structure and differentiation are transformed in their encounter with the human.

Onticology– A Manifesto For Object-Oriented Ontology Part I Posted By Larval Subjects Under Object-Oriented Philosophy. The neural networks of the human brain act as very efficient parallel processing computers co-ordinating memory related responses to a multitude of input signals from sensory organs. Information storage, update and appropriate retrieval are controlled at the molecular level by the neuronal cytoskeleton which serves as the internal communication network within neurons. Information flow in the highly ordered parallel networks of the filamentous protein polymers which make up the cytoskeleton may be compared to atmospheric flows which exhibit long-range spatiotemporal correlations, i.e. long-term memory. Such long-range spatiotemporal correlations are ubiquitous to real world dynamical systems and are recently identified as signature of self-organized criticality or chaos. The signatures of self-organized criticality i.e. long-range temporal correlations have recently been identified in the electrical activity of the brain. A recently developed non-deterministic cell dynamical system model for atmospheric flows predicts the observed long-range spatiotemporal correlations as intrinsic to quantum-like mechanics governing flow dynamics. The model visualizes large scale circulations to form as the result of spatial integration of enclosed small scale perturbations with intrinsic two-way ordered energy flow between the scales. Such a concept maybe applied for the collection and integration of a multitude of signals at the cytoskeletal level and manifested in activation of neurons in the macroscale. The cytoskeleton networks inside neurons may be the elementary units of a unified dynamic memory circulation network with intrinsic global response to local stimuli. Cite as: arXiv: chaodyn/9809003. The neural network of the human brain responds as a unified whole memory bank to a multitude of input signals from the environment and functions with a high degree of robustness and stability. The three aspects of neural networks memory bank are, storage, real-time update and retrieval. The memory is believed to be embedded in the strength of the numerous connections or synapses in the network. Sensory inputs (electrical) produce particular patterns of activity in groups of neurons which then trigger optimal response to the input signal. The cooperative response of millions of neurons to a multitude of input signals has been compared to a very efficient parallel processing computer with neurons and their synaptic connections as fundamental units of information processing, like switches within computers. However, recent studies by Hameroff et al [1,2] and Rasmussen et al [3] show that neurons and synapses are extremely complex and resemble entire computers, rather than switches. The interiors of neurons (and other eucaryotic cells) are now known to contain highly ordered parallel networks of filamentous protein polymers collectively termed the cytoskeleton. Information storage, update and appropriate retrieval are controlled at the molecular level by the neuronal cytoskeleton which serves as the internal communication network within neuron. Organization of information at the molecular level in the cytoskeletal network contributes to the overall response of each neuron and the collective activity pattern of neurons then governs the response to the environmental stimuli. The general awareness or consciousness of the individual to environment may also be governed by the overall background activity pattern of the neurons and their cytoskeletal networks. Coherent signal flow patterns in neural networks may form the basis for general consciousness and response to stimuli (external or internal). Inputs signals trigger spontaneous appropriate coherent pattern formation in the activity of the neurons with implicit spatial correlations in the activity pattern. The time variation of electrical activity of the brain as recorded by the Electro Encephalogram (EEG) exhibits fluctuations on all scales of time, i.e. a broadband spectrum of periodicities (frequencies) contribute to the observed fluctuations [4]. Power spectral analysis which is used to resolve the component frequencies (f) and their intensities shows that the intensity (power) of the component frequencies follows the inverse power law form $1/f^B$ where B is the exponent. Inverse power law form for power spectra of temporal fluctuations imply long-range temporal correlations, i.e. long - term memory of short - term fluctuations. The signatures of short - term fluctuations are carried as internal structures of long - term fluctuations. Time variation of spatial activity pattern in neural networks therefore has inbuilt long - term memory. Neural network activity patterns therefore exhibit long - range spatial and temporal correlations. Such non-local connections in space and time are ubiquitous to time evolution of spatially extended dynamical system in nature and are recently identified as signature of self-organized criticality [5]. Examples of dynamical systems, i.e. systems which change with time include atmospheric flows, electrical activity of the brain, heart rhythms, stock market price fluctuations, etc. Extended dynamical systems in nature have self similar fractal geometry. Self similarity implies

that submits of a system resemble the whole in shape. The world fractal coined by Mandelbrot [6] means fractional or broken Euclidean geometry appropriate for description of non-Euclidean structure generic to natural phenomena.

The fractal dimension D is given by $d \ln M / d \ln R$ where M is the mass contained within a distance R from a point within the extended object. A constant value for D implies uniform stretching on logarithmic scale for length scale range R . Objects in nature exhibit multifractal structure, i.e. the fractal dimension D varies with length scale R . Fractal architectures generic to nature support functions which exhibit fluctuations on all time scale, i.e. the fluctuations are irregular (nonlinear) and apparently chaotic. The association of fractal structures with chaotic dynamics has been identified in all dynamical systems in nature. Fractals, chaos and nonlinear dynamics or Chaos Science is now an area of intensive research in all branches of science [7]. Incidentally, Chaos Science began in 1963 with identification of sensitive dependence on initial conditions resulting in chaotic solution for computer realizations of deterministic nonlinear mathematical model of atmospheric flows and named appropriately deterministic chaos. The computed trajectory of time evolution exhibits fractal geometry. The discipline of nonlinear dynamics and chaos began with investigation of universal characteristics of deterministic chaos in nonlinear mathematical models of dynamical systems in all branches of science. In mathematics, the Cantorian fractal space-time is now associated with reference to quantum mechanical objects [8,9,10]. Further, El Naschie has shown that fractal structures (space-time) incorporate the golden mean equal to $(1 + \sqrt{5})/2 = 1.618$ in their architecture signifying ordered signal /information flow in the fractal network. The golden mean is incorporated in the fractal architecture of the cycloskeleton network [11] which plays a very important role in sub-consciousness to consciousness process integration [12,13]. Surprisingly similar chaotic behavior in space and time was found to be exhibited by all real world dynamical systems. Fractal structure to the spatial pattern concomitant with chaotic (irregular) dynamics has now been identified to be intrinsic to physiological and biological systems [14,15]. The branching interconnecting networks of neurons and intra-neuronal cytoskeleton networks are fractal structures which generate electrical signal pattern with self-similar fluctuations on all scales of time characterised by $1/fB$ power law behavior for the power spectrum. Such inverse power law form for spectra of temporal fluctuations implies long-range temporal correlations, i. e., long term memory of short term fluctuations or events. Fractal architecture of neural networks supports and coordinates information (fluctuations) flow on all time and space time scales in a state of dynamic equilibrium, now identified as self-organized criticality, is ubiquitous to natural phenomena (living and non-living) and is independent of the exact details of the dynamical processes governing the space-time evolution. The physics of self-organized criticality or deterministic chaos is not yet identified. The physical mechanism governing self-organized criticality should be universally applicable to diverse biological, physical, chemical and other dynamical systems. In this paper a universal cell dynamical system model for self-organized criticality applicable to neural networks of the brain is summarised [16, 17, and 18]. This model was originally developed to explain the observed self-organized criticality in atmospheric flows [19, 20, and 21]. Therefore a brief description of the model with respect to atmospheric flows in first described followed by application to neural networks. Atmospheric flows exhibit self-organized criticality or long-range spatiotemporal correlations manifested in the self similar fractal geometry to the global cloud cover pattern concomitant with inverse power law form $1/fB$ for power spectrum of temporal fluctuations in meteorological variables such as temperature, pressure, etc. documented by Tessier et al [22]. The co-operative existence of fluctuations ranging in size (duration) from the turbulence scale of millimetres (seconds) to the planetary scale of thousands of kilometres (years) contribute to coherent weather pattern in atmospheric flows. Townsend [23] postulated that large eddies (waves) form in atmospheric flows as a chance configuration (envelope) of enclosed turbulent (small scale) eddies. A hierarchical continuum of eddies is therefore generated with larger eddies enclosing smaller eddies. Since large eddy is but the integrated mean of enclosed turbulent eddies, atmospheric eddy energy (kinetic) distribution follows normal distribution characteristics according to the Central Limit Theorem in Statistics. The eddy kinetic energy represented by square of eddy amplitude then represents the probability density. Such a result that the additive amplitudes of eddies, when squared, represent probability densities is observed in the subatomic dynamics of quantum system such as the electron or photon. Atmospheric eddy energy spectrum therefore follows quantum-like mechanical laws [19,20,21]. Condensation of water vapour in updraft regions of large eddies give rise to cloud formation while adjacent downdraft regions are associated with evaporation and cloud dissipation, thereby accounting for the discrete cellular

structure to cloud geometry.\ Under the head of applications authors' describe the following: Frohlich [24] had described analogous self-organization of vibrational modes of all frequencies triggering coherent activity in biological functions. Insinna [25] has summarized Froehlich's coherent excitation concept as follows. More than 20 years ago Frohlich [24,26-28] introduced the concept of cooperative vibrational modes between proteins. Coherent oscillations in the range of 1010- 1012 Hz involving cell membranes, DNA and cellular proteins could be generated by interaction between vibrating electric dipoles contained in the proteins as a result of nonlinear properties of the system. Through long-range effects proper to Froehlich's nonlinear electrodynamics a temporospatial link, is, in fact, established between all molecules constituting the system. Single molecules may thus act in a synchronized fashion and can no longer be considered individual. New unexpected features arise from such a dynamic system, reacting as a unified whole entity [25]. Coherent Frohlich oscillations may be associated with the dynamical pattern formation of intraneuronal cytoskeletal architecture which coordinates and integrates information flow into the neuron and generates output signal. Hameroff and colleagues [1,29-31] have simulated such interaction in their cellular automata model. Conclusion they draw in is herein below: **Fractal architecture to information flow path results in spatiotemporal integration of signals so that the fractal system responds as a unified whole to a multitude of input signals. Two disparate examples for such self-organized information flow networks are atmospheric flows and the neural networks of the human brain.** **Cantorian Fractal Spacetime and Quantum-like Chaos in Neural Networks of the Human Brain** **A. M. Selvam (for reference s please see the original article. Models are given elsewhere in the papers series .** If The Holographic Principle Is True, Than It Must Be The Fundamental Principle Of Mind. The Brain Has No Way Around The Holographic Principle. For The Reductionist, The Holographic Principle Is The Ultimate Reduction. It Applies To The Most Minuscule Level Of What We Can Observe, And Beyond. For The Universalist, The Holographic Principle Gives Us The Ultimate Universal. It Extends To The Limit Of Our Universe, The Universal Holographic Boundary, And Beyond. For The Phenomenalist, The Holographic Principle Gives Us The Ultimate Ground Of Our Phenomenal Perception, Our Grounding In The Universal "Now," In The Now-Present Of The Universal Holographic Boundary, Moving Outward From Now-Pasts To Now-Futures. Wheeler (1988) Has Said That All Of Reality Is Information, And That Other Physical Quantities Are "Mere Incidentals." Information Monism Is Gaining Popularity In Physics. By Breaking The Dichotomy Of Between Information And Experience, We Find A Deep Connection Between Wheeler Is Monism And The Experiential Monism Of Whitehead, Sometimes Called Panexperientialism, Which Relates To The Later Theological Concept Of His Student, Charles Hartshorne, Panentheism, God Inside Of Everything (Hartshorne, 1964). There Thus Seems To Be A Convergence Of The Concepts Of Information, Experience, And Spirituality. There Is Not One Universe, But Many Parallel Universes, Or, More Accurately, A Vast Superposition Of Universes Called The Multiverse (Penrose, 2004). However, As Explained In Our Treatment Of The Double-Slit Experiment, In The Observation Of Events On The Quantum Level, The Individual Observer Sees Only One Of These Vast Superpositions, Not A Summation Of A Vast Number Of Potentials. This Has Been Explained In Terms Of The Many Worlds Or Many Minds Theories, In Which There Are Multiple Copies Of The Same Observing Individual, But This Particular Issue Remains Unresolved, Leaving Physics Ungrounded In Reality (Penrose, 2004). The Concept Of One Mind Is Not Only More Parsimonious Than That Of Many Minds, But Also Lets Us Out Of The Bizarre And Counterintuitive Idea That There Are Multiple Copies Of Our Own Selves, Which Exist As Mere Potential, And Are Thus Not Actual. There Are Many Possible Universes, But There Is Only One Mind, Which Determines Events On The Quantum Level, And Thus Creates Our Universe. As We Had Discussed Previously, The Quantum Level Provides The Essential Ground Of The Holographic Principle, Such That Quantum-Level Holographic Surfaces Are Elaborated At Higher Levels, Manifesting Higher Orders Of Information From The Quantum "World." In This Process, There Is A Reduction Of The Wave Function Or Potentialities Of Quantum Fields. At The Level Of Consciousness, This Entails Freedom To Choose Which Observations We Make (Stapp, 1997), Which Gives Us The Capacity To Think And To Make Decisions. These Capacities Are The Basis Of Individuality, Self-Determination, Judgment, And Values. The Progressive Evolution Of The Manifestation Of Mind Through Higher Orders Of Experience, Leading To Higher Orders Of Consciousness, Is Entailed By The Holographic Principle. Again, We Are Dealing Here With Levels Of Description, With The Multiverse Of All

Potentials Fundamentally Supporting The Single Universe We Collectively Observe. The Multiverse Is The Wave Function Of The Universe (Penrose, 2004). The Recursive Integration Of Nested Hierarchies Of Holographic Surfaces Brings Out A Single Actuality In Consciousness From A Wave Function That Is Unconscious. Consciousness Thus Gives Us Information At A Level Of Experience That Is Causal. In This Sense, We Partake In The Creation Of The Universe. We Participate In Creation, And This Participation, When Fully Realized, Leads Us To Higher Levels Of Consciousness And Of Realization. As A System, The Biosphere That We Live In Has A Holographic Surface, Creating A Deep Sense Of Ecology As We Collectively Move Toward A Planetary Consciousness. It Is Only When The Universal Holographic Boundary Reaches The Information Storage And Processing Capacity Needed For The Requisite Biological And Biochemical Complexity That Consciousness Evolves In Living Things. This Evolution Is Natural And Spontaneous, Since Consciousness Gives Rise To What Is Actual, As Opposed To What Is Merely Potential. In This Sense, Consciousness Is Still In The Process Of Creating Our Universe, And Levels Of Higher Consciousness Will Continue To Evolve. We Are All In The Same "Now," And That Now Is Defined By The Present Universal Boundary. This Assures Us That Our Experience Is Universal, And Does Not Pass With Time. This Is The Fundamental Basis Of Memory And Of Cognition. The Identity Of Mind And Brain Is A Myth. We Have A Continual, Internal Or Non-Local Relation With The Universe, As It Has With Us. Once This Mystery Is Resolved, The Myth Is No Longer Needed, And There Is A Confluence Of Science And Spirituality. Experience Is Primary, Information Is Secondary. We Can Only Gather Information From Experience, Whether It Is In The Laboratory Or In Life. We Cannot Measure The Information On The Surfaces Of Systems. The Physicists That Have Formulated The Holographic Principle For All Systems Are Quite Aware Of This, Or Else The Principle Would Have Been Established Or Discredited. We Cannot Measure What We Experience. It Is Intangible, Yet It Is All We Know To Be Actual. Everything Else Is Inferred. Because It Cannot Be Measured, It Has Been Fundamentally Disregarded By Mainstream Science. Materialism Is Considered Scientific, While Idealism Is Considered Unscientific. But Aren't Ideas, Fundamentally, Made Of Information? Brain Science Has Mistaken The Representation Of Information For Information Itself, And Has Tied Those Representations To Matter And Energy. Consciousness, The Highest Order Of Information, Has Generally Been Regarded As Superfluous, Something That Needs To Explained Away, Or Altogether Ignored. Yet It Is The Only "Thing" That Reaches Our Awareness. Consciousness Comes At A Price. For Everything That Becomes Conscious, There Must Be Something That Becomes Conscious. Consciousness Is Certainty, And Its Complement, Unconsciousness Uncertainty. Consciousness Is The Particle Nature Of Experience. It Has Definiteness About It. The Unconscious Is The Wave Nature Of Experience, It Is Like The Metaphysical Cloud Of Unknowing. If We Are The Most Conscious Of Animals, Then We Must Also Be The Most Unconscious. Perhaps This Is The Predicament Of Humankind. The Holographic Principle Of Mind Leads Us Naturally From The Most Fundamental Experiences, Existing As Quantum Potentials From The Conformations Of Proteins Down Through The Fields Of Electrons, Through Their Manifestation Upward Through A Recursion Or Successive Applications Of The Same Holographic Process, Through Higher Levels Of Experience, To The Emergence Of Consciousness As Higher And Higher Orders Of Experience. **The Quantum Holographic Principle Mind Does Not Require Anything More Quantum Than Is Obviously Present At The Microscopic And Submicroscopic Levels, As It Represents Successive Orders Of Manifestations From These Levels.** Recursion Also Applies Here In The Sense That It Is Used In Computer Science, In Which The Function Of The Part Depends On The Function Of The Whole. A Program, As A Part, Cannot Work Without A Functioning Whole, The Operating System Recursive Wholes Which Are, For Us, Supra-Conscious, Are On The Group, Species, Planetary, And Universal Levels. As Individuals, Our Consciousness Cannot Function Without Recursion To The Universal Consciousness, Even Though We May Be Unaware That Such Universal Consciousness Exists. Reaching Upward To These Supra-Conscious Levels Is A Spiritual Process, Making Transcendence The Ultimate Solution To The Unconscious Human Predicament. What Was Once Supra-Conscious Becomes Unconscious Through A Process Of Conditioning? We Are Born As Creatures Of The Earth And Of The Universe, As Evidenced By The Beliefs And Practices Of "Primitive Peoples." There Is Evidence From Cave Paintings That Our Hominid Ancestors Experienced A Kind Of Holographic Perception (Combs, 1996), Which Could Constitute Our Early Connection With The Holographic Subtext Of Reality, And Which We Might Then Propose Existed In Our Animal Ancestors, And In Extant Animal Species. If This Is The Case, Than

Microgenesis Would Entail The Recapitulation Of This Holographic Experience As It Progresses Through Our Ancestral Past. The Supra-Conscious Mind Seems To Envelope Perinatal Experiences, And Stanislov Grof (1994) Has Developed Techniques To Gain Access To These Experiences, As Well As To The Earlier Experiences Of Our Human And Animal Ancestral Lineage, And Of Our Universal History. Grof (1994) Concludes: "Our Consciousness Seems To Have The Amazing Capacity To Directly Access The Earliest History Of The Universe Ñ Witnessing Dramatic Sequences Of The Big Bang, The Formation Of The Galaxies, The Birth Of The Solar System, And The Early Geophysical Processes On This Planet Billions Of Years Ago." The Holographic Principle Theory Of Mind MARK GERMINE Institute For Psycho Science There Are Similarities Between The Patterns Of Holography And Of Psychological Transference, Where Holography Is The Process Of Recording And Reconstructing Holograms Employing A Theoretical Perspective Using A Hermeneutic Method, This Dissertation Parallels Holography With Transference, Offering Another Way To Encounter Transference By Showing Similarities Between The Processes Of Each And The Results Of Each. Though Complex, Infinitely Varying, And Unique, Their Patterns Are Clearly Identifiable. Thus They Are A Metaphorical Fit To The Concept Of Strange Attractors In Physics And A More Literal Fit To The Concept Of Archetypes In Depth Psychology Or Dynamic Psychology, Psychology Which Attends To The Living, Autonomous Unconscious. This Study Explains How Holography Models Transference, What A Hologram Is And How It Works, And How Depth Psychology Understands Of The Interaction Between Consciousness And The Unconscious Is Related To The Hologram. It Describes Transference And Related Psychological Processes As Understood In Six Different Schools Of Depth Psychological Thought. It Shows That The Underlying Pattern Or Strange Attraction Between Transference And Holography Extends To Other Processes Both Within And Outside The Field Of Psychology, Processes Such As Projection, Projective Identification, Splitting, Memory, Biology, Creative Discovery, Theology, Synchronicity, Chaos, And Nonlocality. By Identifying The Similar Patterns Of These Processes, This Study Demonstrates The Existence Of An Underlying Holographic Archetype In Which Essential Qualities Of The Whole Are Present In Each Of The Parts Of The Whole: The Visual Image Of The Overall Hologram Is Present In Each Component Part Of The Hologram, The Autonomy Of The Overall Human Is Present In Each Conscious And Unconscious Component Part Of The Human Psyche. By Noting Differences As Well As Similarities In These Processes, It Suggests An Inventory Of The Qualities Of The Holographic Archetype. This Study Furthers Understanding Of The Pervasiveness, Force, And Autonomy Of The Unconscious Acting Through Transference And Projection By Identifying A Group Projection Of Domestic Violence Lying At The Core Of The Christian Myth. This Study Also Furthers Understanding Of The Concept Of Transference By Providing A Reflection Hologram Of The Human Psyche As An Artistic Work And As A Visual Metaphor Of Transference.

Strange Attractors: Transference, Holography, And An Archetype Burke, J. (2003). **Strange Attractors: Transference, Holography, And An Archetype (Doctoral Dissertation, Pacifica Graduate Institute, 2003).** A host of observed, but very basic human phenomena, including consciousness itself, have eluded rigorous scientific description by all disciplines of science. This is true, not because of insufficient evidence for a particular phenomenon's existence, but rather for lack of a theoretical construct, which could fit within the prevailing paradigms of science. For the past century eminent men and women of science have accumulated thousands of pages of data on mind/mind and mind/matter interactions. Many of the most telling experiments have been criticized, perfected and repeated numerous times during the past five decades, using increasingly sophisticated technologies. **Meta analysis of these experiments produce accumulated probabilities against chance occurrences exceeding trillions to one (Radin, 1997).** It has required, however, that quantum science mature for seventy-five years and during that period, test, validate and synthesize a number of seemingly outrageous physical concepts arising from quantum theory, before testable theories could arise which offer hope that anomalous mind and consciousness data can be explained (Mitchell, 1996). The missing concepts that prevented the earliest investigators of consciousness from succeeding in their quest were 1) a **generalized theory of information, and 2) quantum science itself, with the associated phenomena of non-locality, the zero point energy field and the quantum hologram.** These associated phenomena are still not well understood but are sufficiently validated today by both theory and experiment to provide a basis for postulating a necessary condition for the existence of consciousness phenomena, as experienced in the observable four dimensional space/time universes. A third concept, chaos theory, is also necessary to understand the nonlinear

evolutionary processes that caused consciousness to evolve toward the anthropic consciousness experienced by humans. In particular, chaos theory maps far from equilibrium systems and demonstrates the irreversibility of nonlinear processes and thus the irreversibility of time in the macro-scale universe. Another class of phenomena, including normal sensory perception and evolution, to cite but two, have explanatory theories in classical science, but which in view of current developments in late quantum physics and in chaos theory may be incomplete approximations to the correct theory. Information concepts have been examined by Weiner, von Neuman and Shannon in well-known seminal works and by Frieden more recently (1998) to produce theories useful to physics, to computation and to communications technologies. These theories, although accurate and mathematically useful in their domains, fall short of being sufficiently encompassing when considering the problem of consciousness, its evolution and its associated phenomena. Even relatively simple perceptual organisms utilize patterns of energy, that is, information, not completely described by existing mathematical theories. Theory and experimental evidence for the zero point energy field has been published by many authors, but I shall cite Haisch, Rueda and Puthoff, (1997, 1998), as the most contemporary and relevant work for this paper. Theory and experimental evidence concerning the quantum hologram has been developed by Schempp (1992, 1993) and Marcer (1996, 1997, 1998), separately and jointly, based upon a new understanding of quantum mechanics. (See previous work by Cramer [1986], Berry [1988], Anandan [1992] and Resta [1997]). Non-locality, although predicted by the earliest work in quantum theory and decisively demonstrated by Aspect in 1982, has been thought to be a curious property of particle physics but of little relevance to macro-scale reality until discovery of the quantum hologram. Further, it is widely believed that non-local quantum information represented by entanglement of particles could not be recovered locally as useable information (Eberhard's theorem). However recent work both in theory and experiment (e.g. see Nature, 1997, 11th December, vol. 390, Sudbury T pp 551-552, and Bouwmeester D. et al pp 575-579) is in line with the work by Berry, Resta, Schempp and Marcer and makes it clear that this is not the general case for quantum information processing and communication. It has been widely accepted in science, until recently, particularly in the field of artificial intelligence, that the brain was likely a complex classical computer, incapable of supporting quantum processes. The work by Hammeroff (1994) and Penrose in isolating and describing microtubules in brain tissue have caused a re-examination of this dogma, and renewed interest in uncovering the quantum processes involved. Based upon this earlier work I postulate and examine the evidence in this paper for the following theories:

1. The basis of subjective experience is rooted in the quantum attribute of nature called non-locality. I will use the word "perception" in its most generic sense to denote a basic subjective experience at all levels of complex matter. Thus the non-local quantum correlation between entangled quantum particles is considered the root cause of the phenomenon experienced as perception in more complex matter, but the non-local quantum hologram is the non-local carrier of information for molecular and larger scale matter. Thus, perception is not an object but rather the label for a nonlinear process involving an object, a percipient and information.
2. The experience of humans is that they sometimes, perhaps often, perceive information from or about physical objects that is not available through normal, local, sensory mechanisms, nor classical space/time information. Objective testing data in overwhelming abundance provides evidence that this is true, though an explanatory mechanism has until contemporary times remained elusive. I shall call this intuitive information or intuitive perception. I postulate that a quantum hologram is the source of this intuitive perception and that the percipient is at that time in phase-conjugate-adaptive-resonance (pcar) with the entity or object associated with the quantum hologram.
3. The phenomenon of "learning" in humans is a subjective process that involves perception, memory, intentionality, and evaluation of outcome and behavior change. This may be viewed as a classical nonlinear feedback loop. Although we cannot know precisely the subjective experience of another entity, presumably in the successful training of animals, an analogous subjective process is in effect. Sheldrake (1981) has published a successful theory of morphic resonance related to animal learning based upon non-local information. Marcer has published papers (1996, 1997) theorizing a mechanism by which the quantum hologram causes learning to take place in both DNA molecules and prokaryote cells as an adaptation process of environmental resonance, rather than mutation and adaptation solely by random processes. I postulate that Marcer's concept can be generalized to nature at large and that the quantum hologram is the information structure suitable to explain Sheldrake's morphic resonance. The non-local quantum correlations observed in particles, and the non-local quantum hologram associated with molecular and larger scale objects, serve the purpose of providing

information at all scale sizes to guide evolutionary processes. That is to say, that quantum non-locality is the basis of perception, and thus fundamental and necessary to the complex organizations of matter and information in the universe. Further, since learning is an observed property of complex systems such as animals and, via the quantum hologram, is theorized to be a property of simple cells and molecules, one can also postulate the generalization that nature evolves through a learning process rather than because of random mutations. 4. Marcer (1997) has proposed that the condition of phase-conjugate-adaptive-resonance (pcar) is a necessary condition for an object in three-dimensional reality to be perceived as it really is. That is, resonance requires a virtual path mathematically equal but opposite to the incoming sensory information about the object. Further, that it is the incoming electromagnetic (space/time) information (visual, acoustic, etc), which decodes the information of the quantum hologram and establishes the condition of pcar so that accurate three-dimensional perception is possible. That is to say, both quantum information and space/time information are used in the act of perception by organisms having sensory preceptors. I propose that the two equal but opposite paths required by the pcar condition are the mathematical equivalent of perception and attention (or intention). (I shall distinguish between attention and intention in following pages.) Discussion The anecdotal evidence for humans perceiving non-local information dates to prehistory. The data were sufficiently robust that both experiencers and philosophers, from Plato and Aristotle forward, accepted that both physical and non-physical realms of reality must exist. Non-physical was thought to explain the subtle, ephemeral and mystical subjective experiences ubiquitously reported in human culture. After Descartes and Newton, however, classical western science rapidly discarded the non-physical hypothesis and systematically began to ignore all evidence for perception of non-local information. Field theories and point particles were created to preserve the concept of physical contact between particles and to explain obvious examples of "spooky action at a distance" such as gravitation and electromagnetic interactions. Information, broadly defined as patterns of energy, reemerges however, in non-local form in the mysterious quantum spin correlations of double slit experiments, although it has been widely believed that such non-local information could not be recovered and utilized by sensory systems. With validation of theory and experiments concerned with the non-local quantum hologram, information, including non-local information, suddenly acquires a more important status in physical theory, a status as important as energy itself. This is true because information is the basis of the cognition and knowing by which creatures perceive reality, and non-local information can now be seen as a ubiquitous and useful property of the cosmos, rather than a unique attribute of particles (and human animals). It is likely that most, if not all, subtle, ephemeral and unexplained phenomena associated with subjective experience are connected, directly or indirectly, with the phenomenon of non-locality. The brain is clearly a quantum computer (Schempp & Marcer, 1996) which utilizes both quantum and space/time information. This discovery alone almost certainly sets a necessary, but not sufficient condition, for intelligent life to have arisen in the cosmos, wherever environmental conditions permit. Many volumes have been written in this century by scientists experimenting with remote viewing, ESP, telepathy, clairvoyance, precognition, etc. Police agencies routinely use "psychics" to assist in criminal cases often with success. Intelligence agencies of governments have clandestinely utilized the findings to successfully gain information about an adversary. Many reports of these activities have been recently declassified and printed in open professional journals, even though no explanatory physical mechanism has yet been reported which is acceptable to mainstream science. The most succinct modern summary of this activity and analysis of results have been published by Radin (1997). Quantum Holography Non-locality and the non-local quantum hologram provide the only testable mechanism discovered to date which offer a possible solution to the host of enigmatic observations and data associated with consciousness and such consciousness phenomena. Schempp (1992) has successfully validated the concept of recovery and utilization of non-local quantum information in the case of functional Magnetic Resonance Imaging (fMRI) using quantum holography. Marcer (1995) has made compelling arguments that a number of other chemical and electromagnetic processes in common use have a deeper quantum explanation that is not revealed by the classical interpretation of these processes. Hameroff (1994) and Penrose have presented experimental data on microtubules in the brain supporting quantum processes. The absorption/re-emission phenomena associated with all matter is well recognized. That such re-emissions are sufficiently coherent to be considered a source of information about the object is due to the theoretical and experimental work of Schempp and Marcer, based upon the transactional interpretation of quantum mechanics of Cramer (1986), the Berry geometric phase analysis of information (Berry, 1988; Anandan,

1992) and the ability of quantum phase information to be recovered and utilized (Resta, 1997). The mathematical formalism appropriate to these analyses is consistent with standard quantum mechanical formalism and is defined by means of the harmonic analysis on the Heisenberg nilpotent Lie group G , algebra \mathfrak{g} and nilmanifold (see Schempp (1986) for a full mathematical treatment). The information carried by a quantum hologram encodes the complete event history of the object with respect to its three dimensional environment. It evolves overtime to provide an encoded non-local record of the "experience" of the object in the fourdimensional space/time of the object as to its journey in space/time and the quantum states visited. The question of the brain's ability, as a massively parallel quantum processor, to decode this information is addressed by Marcer and Schempp in "Model of the NeuronWorking by Quantum Holography" (1997) and "The Brain as a ConsciousSystem" (1998). They argue that an organism's ability to perceive objects as they are and where they actually are in three-dimensional reality requires the phase conjugaterelationship provided by quantum holography. It is not sufficient for the incomingelectromagnetic illumination (or acoustic signal) carrying object information to present tothe brain a wave front in the manner presented to a flat photographic plate. Rather, a virtual signal as mapped by the phase conjugation of quantum holographic formalism is required to decode the information in order for perception and cognition to exist as weexperience it in three dimensional realities. The percipient and the source of information are in a resonant relationship for the information to be accurately perceived. Manyinvestigators have proposed a holography mechanism as a basis for brain functioning,beginning with Pribram, and indeed, others have proposed holography as a construct forthe universe itself, but discovery of the non-local quantum hologram created by theabsorption/remission phenomenon and characteristic of all physical objects provides thefirst quantum physical mechanism compatible with macro-scale three dimensional worldas we experience it. The existence of a quantum hologram associated with each physical object provides eachphysical object with the non-local waveform predicted by quantum theory's wave/particleduality and extends quantum theory to all physical matter. It allows, for the first time, apossible approach for understanding the mysterious world of consciousness. Postulating that this is globally true, we inhabit a quantum world where non-local effects should be expected at all levels of functioning, not just as a curious artifact of the subatomic level ofreality. Existence of the non-local quantumhologram suggests that nature has utilized non-local information from the big bangforward, throughout its evolutionary history; and long before planetary environments selforganized to permit living matter and complex space/time sensory systems to evolve. Thepapers of Marcer and Schempp on learning inherent in DNA and prokaryote cells usingquantum holography, when generalized, helps explain the ubiquitous appearance in nature across distances, scale sizes and species, of similar processes, organs and sensory systems. This certainly conforms to the fractal geometry of chaos theory. Certainly the similaritiesof DNA, cell structure, organs and brains across species are easier to reconcile with a non-local learning process than with a theory of localized random mutation and natural adaptation. It is important to observe that in standard particle physics experiments the object is todiscover the quantum characteristics of the individual types of particles, and the conditions under which they split and recombine. In quantum holography the object is totreat the entire group of re-emitted quanta as a whole, and as in laser holography, toexamine the information carried in the interference pattern and phase relationships. These represent two quite different levels of approach to quantum information. In particle experiments, it is considered that the eigenvalues of the applicable matrix represent measurable values; and that information is lost during measurement due to decoherence of the particles and energy exchange. But in the quantum holographic formalism, the information is carried in the phase relationships, which are represented by off-diagonal terms in the matrix, and the information is recoverable under the proper conditions as Berry and Resta have predicted, and as Schempp has demonstrated with fMRI. The quantum mathematics is consistent with standard quantum theory in both cases. In decoding the quantum holographic information, however, the energy exchange is insignificant. The similarity of the mathematical treatment in these various experiments is important tothe thesis of this paper. In examining the quantum non-locality of particles it is spin numbers and/or polarization that are the parameters of interest. A standard technique of analysis is to use the Fourier transform to map the state of the particles into the frequency domain. In the formalism of the quantum hologram, mapping into the frequency domain is also fundamental; however, the requirement for pcar assures that the phase relationships are matched so that the percipient (sensory system) is able to decode the information carried in the phase relationships. It is precisely the pcar requirement that permits the encoded

holographic information to be decoded by the percipient. Mathematically, decoding is simply reversing the rotation of the phase vector in phase space. Physically, it is matching the frequencies and phase of the information such that resonance results. Frequency, phase matching and resonance are an operational characteristic of every type receiver technology. Pribram's earliest proposal that the brain stored information encoded as in a hologram and mapped by the Fourier transform is in complete agreement with the evidence presented by quantum holographic mathematical formalism. It is the spin and polarization attributes of particles (both are mapped by wave mathematics) that represent the puzzling non-local property of subatomic matter. It is the phase relationships that carry the information in holography (again mapped by waveform mathematics). And it appears that the brain stores and manages information not as a classical digital machine, but rather as an **analog device using non-local properties of the quantum hologram, which can be analyzed by wave form mathematics (harmonic analysis on the Heisenberg Lie group)**. In the cosmological evolutionary scheme of things this similarity of appropriate mapping techniques is too bizarre a coincidence to be ignored as a cosmic accident. Thus there is ample evidence that the **non-local attribute of** nature is much more than just a curious artifact of subatomic particle interactions, but rather is a more fundamental phenomenon that appears at all scale sizes and is, in particular, associated with the utilization of information in nature, and associated with the fact that information has a causal effect independent of distance. It is precisely information, however, that is the basis of the phenomena of perception, cognition, memory, learning etc, that is to say, consciousness and the subjective experience. Though the evidence is quite ample to postulate that non-locality is the unique, universal basis for perception and the subjective experience, the evidence though compelling is not sufficient to be conclusive that such is indeed the case. The next steps are to validate more completely with experimental evidence that non-locality plays a major role at all scale sizes and that all physical objects are quantum objects and thus interconnected by information in this strange way. **Non-Locality in Nature:** There is experimental evidence to strongly suggest that simple organisms perceive and respond to information non-locally as well. Cleve Backster was perhaps the earliest to experiment with plants and simple life forms in electromagnetic isolation in late 1960's and early 1970's. His work was not confirmed through replication by others at that time. Other investigators have had mixed results replicating non-local information perception by simple organisms and living tissues. In the area of human experimentation, results likewise have been mixed and controversial for three-quarters of a century. However Meta analysis by Radin (1997) and independently by Utts (1991) across a large and appropriate spectrum of experiments demonstrates compelling statistics that the **perception of non-local information exists and is real**. Perhaps was there a larger body of experimental evidence for simple life forms, similar Meta patterns would emerge. Failure to replicate results in well constructed experiments does not, in the case of subtle consciousness phenomena, prove that the phenomenon is missing but rather that a **hidden mechanism** below the threshold of classical measurement is operating. For example, the most telling experimental evidence to explain the sometimes inconsistent results relates to direct non-local observer and/or experimenter effects. Gertrude Schmedler isolated the **"sheep/goat"** effect in human experimentation decades ago (1972). Experimenters and/or participants in a human telepathy (or similar non-local) experiments exhibited results statistically above or below chance results depending upon their subjective bias toward the experiment. (In other words, 100% wrong answers would be as statistically significant as 100% correct answers in such tests, and in addition betrays the mind set or intention of the subject; whereas only chance results would be inconclusive.) More recently, a series of experiments by Marilyn Schlitz (1997) investigating "intentionality" clearly demonstrated that experimenter bias (intentionality) affected the outcome even of double blind experiments. Thus, in the subtle realms of mind and consciousness studies, bias, belief and intention clearly have an effect. The lack of an existing theoretical structure in classical science to support any type perception of non-local information, much less to support bias, belief or intention as having a non-local effect, when in fact it does have a non-local effect, is quite sufficient to account for anomalous results in many scientific experiments. Further validation and acceptance of the non-local thesis will have strong positive repercussions for the prevailing scientific paradigm and particularly the theory of measurement. **Attention and Intention:** A powerful and telling series of experiments conducted by Dean Radin (1997) at University of Nevada at Las Vegas following a decade long set of equally significant experiments by Brenda Dunne and Robert Jahn at Princeton University (1988) provide insight as to the subtleties involved in this level of mind/brain functioning. Jahn and Dunne provided overwhelming evidence that **subjects could intentionally produce statistically skewed results** in mechanical

processes normally thought to be driven by random processes. Radin went further; he discovered that audiences watching a stage performance would skew the output of nearby random number generators during periods of high emotional content in the stage performance. Further, in a wide-ranging audience participation experiment, he recorded the output of computer random number generators during the television broadcasts of the O.J. Simpson murder trial. Most television media reported this event for weeks on end and tens of millions of humans were watching the results. Again, the results of the random number generators were skewed corresponding to emotional peaks during the trial drama and corresponding to the number of people watching television. The thesis in the Princeton experiments was that participant intentionality **created a non-random effect to bias the skewed distribution**. In the Radin experiments the results were not intentional, as the participants were unaware of the experiment, but the hypothesis was that attention (in particular, rapt attention) drove the system away from chaos (randomness) and toward greater order (reduced entropy). These results suggest that attention and intention provide closely correlated outcomes, further, that randomness may not be a general property of nature, but that what is perceived as random noise in a system may be information (a pattern of energy) that is not in resonance at that moment with the particular perceptual system. William Tiller, emeritus professor at Stanford also has performed experiments (1997) that are consistent with these results, though his interpretation of the operating mechanism is somewhat different. These different types of mind/mind, mind/matter experiments have been rigorously and routinely conducted for decades with statistically compelling results but just as routinely dismissed or ignored by main stream science because the implications of non-local action are so foreign to the classical paradigm. However, if we consider that the condition of **phase-conjugate-adaptive-resonance** is necessary to completely specify the act of perception as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as locked in a resonant feedback loop. The incoming wave front carrying information may be labeled as **"perception"** from the point of view of the percipient, and the return path required by the resonant relationship may be labeled **"attention"** (or for subsequent discussion, "intention"). It is a well established principle in the meditative practices of esoteric disciplines that prolonged focused attention on a object of meditation causes the percipient and the object to appear to merge so that a deeper level of information about the object is obtained; information such as history or internal functioning, that would not be available through classical space/time information. The concept of the quantum hologram adequately and completely describes how this phenomenon might take place. Further, it is accepted that the mind/brain is a massively parallel processor, capable of performing many tasks simultaneously and subconsciously (in the right, intuitive part of the brain). Attention (meaning conscious, focused attention) is a unique and singular task that must take place sequentially, mostly in the left cognitive part of the brain. The condition of attention deficit disorder (ADD) is precisely the problem of a percipient being unable to maintain a singular focus for sufficient time to complete a desired task or observation. Thus, the **action of focusing attention by apercipient may be construed as a necessary condition for pcar to be established with the perceived object**. Non-Locality, Near and Far Marcer has presented the case for the pcar requirement in normal sensory perception (visual and acoustic). A frequent modality used by psychic sensitive individuals to gain information is to physically touch an object. Touching an object satisfies the pcar requirement and presumably allows the percipient access to information about the object not available from space/time information. Police agencies frequently use this modality with psychic sensitives to gain information about a crime scene, much as they utilize bloodhound to track the scent of an individual, often with considerable success. If, as in the theory of the quantum hologram, the object has been in the presence of the individual about whom information is desired, the event history of the object and that of the individual intersect. **The Berry phase information of the object contains its journey in three dimensional space and time, as well as the quantum states through which it has passed on this journey. The sensitive individual, with a honed talent, seems often able to decode useful Berry phase information from the object about the individual sought. It may also be the case with the blood hound, which additional non-local information has been gained about the subject, even though the classical explanation is that the animal is operating only with heightened olfactory sensing.** Although perception in the three dimensional world requires and utilizes pcar, most humans, however, do not bring to conscious awareness non-local information when we are routinely operating in three-dimensional reality. We perceive objects as presented by space/time information, that is, shape, color, function (tree, chair, table, etc) but are not usually

aware of the additional non-local information. It takes training as provided by many of the esoteric traditions and/or certain naturally sensitive individuals to routinely perceive the non-local holographic information associated with a particular object. There is massive evidence to suggest, however, that the brain has these latter capabilities at birth. Suppression by cultural conditioning in childhood and subsequent lack of practice cause the natural ability for conscious, intuitive perceptions to atrophy. Particularly in western tradition, educational interest has been on the left brain, rational functions rather than right brain, intuitive functions. However, mystic adepts and natural psychics' routinely demonstrate that non-local information is perceptible from physical objects by focusing attention, quieting the left brain and allowing intuitive perceptions to appear. It is the left brain cognitive ability in humans that provides canonical labeling of the intuitive and artistic processes taking place in the right brain. The fact that with training and practice, individuals can recover, deepen and label their individual cognitive access to intuitive, **on-local information demonstrates that learning is taking place within the whole brain itself and involves enhanced coherence and coordination between the hemispheres**. This process is different and distinct from the left brain function of extending and extrapolating factual data and logical deduction to leap to an "intuitive" conclusion, while omitting the intermediate steps leading to that conclusion. Experimental protocols for remote viewing normally provide clues to the location of the object such as a description, a picture, or location by latitude and longitude, that is to say, an icon representing the object. These clues seem to be sufficient for the percipient to establish a resonance with the object. Normal space/time information (visual, acoustic, tactile) about the object is not being directly perceived by the percipient, nor does the object usually appear at its physical location in space/time like a photograph or map in the mind. Rather, the information is perceived and presented as internal information and the percipient must associate the perceptions with his/her internal data base of experience in order to cognize and to describe the object's perceived attributes. In the case of complex objects being remotely viewed, the perceived information is seldom so unambiguous as to be instantly recognizable as correct. Sketches, metaphors and analogies are usually employed to cognize and communicate the non-local information. A considerable amount of training, teamwork and experience are necessary to reliably and correctly extract complex non-local information from a distant location. The information appears to the percipient as sketchy, often dream-like, and wispy, subtle impressions of the remote reality. Very skilled individuals may report the internal information as frequently vivid, clear and unambiguous. The remote viewing information, being strictly non-local, and in this hypothesis, the information perceived by quantum holography, is missing the normal space/time components of information necessary to completely specify the object. It has been demonstrated that this intuitive mode of perception can be trained in most individuals. Perhaps additional training and greater acceptance of this capability will allow percipients to develop greater detail, accuracy and reliability in their skill. In principle, training will not only increase the skill and accuracy but should cause the appropriate neural circuitry to become more robust as well. In the absence of space/time (electromagnetic) signals to establish the pear condition and to provide a basis for decoding the quantum hologram, an icon representing an object seems to be sufficient to allow the brain to focus on the object and to establish the pear condition. However, a reference signal is also required to provide decoding of the encoded holographic phase dependent information. Marcer (1998) has established, using Huygen's principle of waves and secondary sources, which any waves reverberating through the universe remain coherent with the waves at the source, and are thus sufficient to serve as the reference to decode the holographic information of any quantum hologram emanating from remote locations. The **Zero Point Field: The results of the Michelson/Morley experiment banished the concept of aether from early twentieth century physics**. However, it left a void as the nature of interstellar space and nothing for propagating waves to wave in. Quantum physics reincarnated the aether as the zero point fields, a seething cauldron of quantum potential and unmanifest energy where particles and antiparticles spontaneously arise and then disappear. The very fabricant structure of space/time itself is again in question; its structure and its metric under intense investigation with far more questions than answers having emerged to date. For the purposes of this paper the relevant issues are two: 1) the emission/absorption phenomenon, and 2) the structure and mechanics of non-locality. Zero point (zero degrees Kelvin) emission and absorption of quanta from all physical objects is a well established phenomenon. It is our view that the zero point fields are the plenum (or cauldron) which supports this absorption and re-emission, and makes the phenomenon of the quantum hologram possible at all temperatures. Although particle experiments are carried out under rigid conditions of temperature and

pressure, Schempps experimental work with the fMRI requires no such constraints. There are deep and difficult questions yet to be answered about how the information of the quantum hologram maintains its integrity and is propagated, about how resonance takes place at extremely large distances. There is considerable evidence that intuitively perceived information is truly non-local. It does not obey the inverse square law for space/time energy propagation; it is time independent and cannot be shielded by electromagnetic shielding. Such characteristics are the mark of non-locality. But understanding the mechanics of non-locality (or a visual picture) is missing from standard models. Some physicists turn to superluminal speed of propagation, others to the zero point as a zero dimension, which is resonant with all parts of the universe simultaneously. The issue of instantaneous communication (or at least superluminal communication) of non-local effects on a cosmic scale remains a problem, even though the phenomenon itself is well validated. Perhaps it is a problem of topology. What shape can the universe have such that one point can be in simultaneous contact with all other points? In this regard it is clear that certain problems between quantum mechanics, special and general relativity remain inexistence. Haisch, Puthoff and Rueda continue to investigate the **metrics of the zero point fields** with regard to better defining the unanswered questions about, mass, gravitation and inertia. Perhaps these investigations will also bring answers for how phase related information is propagated non-locally, likely within the zero point fields, and thereby unveil the mechanics of the resonance phenomenon. Further, new investigations reported by Van Flandern (1998) on measurements from orbiting Global Positioning System (GPS) clocks indicate the predictions from Lorentzian relativity to be approximately four times more accurate (0.7% opposed to 3%) than predictions of special relativity. If these measurements are further validated, it implies that Lorentzian relativity with a Hubble absolute rest reference frame, an aether (zero point field), and instantaneous propagation of non-local effects may be the preferred one. If this is the case, then many questions about non-locality would be resolved. Intentionality I have argued that by establishing pcar between a **percipient and an object, the phase conjugate (equal but opposite) paths connecting the two can be labeled "perception" and "attention"**. In the case where the object is a simple physical object (rock, flower, etc.), our interest is on the non-local information perceived by the percipient about the object. However, from the point of view of the object, information about the percipient is also available to the object. The pcar condition is a reciprocal relationship, mathematically. Quantum holographic formalism predicts that the history of events of quantum objects is carried in the quantum hologram, thus we must conclude that the **"attention" focused upon the object causes that event to be recorded in that object's quantum hologram**. Although we cannot query the object about its experience perhaps an experiment such as one utilizing the Aharonov-Bohm effect would detect a phase shift in the object's holographic field. (In this discussion I use anthropic labeling as we are discussing human perception. The phenomena however, are rooted in natural (and primitive) non-local physical processes, which are fundamental. The evolved complexities of perception, cognition, etc, associated with a brain obviously, as yet, have no analogous label to describe the experience of simple objects.) Once the pcar condition is established, the percipient can change its mind state with regard to the object. The perceived information can be operated upon by the brain's function so that cognition occurs with respect to the perceived information and meaning assigned. Cognition and meaning require finding a relationship between the perceived information and the information residing in the percipient's memory. The percipient can then form intent with respect to the object. In such case the path I have labeled "attention" could

Figure 1 In self aware animals (those with a brain) cognition, meaning and intent with respect to an object can often be described in simple terms, for example: enemy, fight or flight; food, eat; friend, greet, etc. The non-local component of information, although present and creating effect, is operating below the level of conscious perception in humans and results in "instinctual" subconscious behaviors in all animals. The brain as a massive parallel computer is simultaneously performing numerous tasks to accomplish the desired intention. Classical modelings of this autonomous activity describe it only in terms of classical information and energy flow in the central nervous system and the brain. However, if non-locality is operating at all levels of activity, as this theory suggests, certainly there are resonances involving non-local information operating throughout the body of an organism in parallel with classical space/time functions. Subsequent experimental work will surely uncover these quantum processes where non-local resonance is involved in the functioning of an animal's internal processes. In the case of non-local effects at a distance, outside the body, simple spin correlations of entangled particles is the most basic. The spin coherence is reciprocal. Action on one particle creates an effect on other entangled particles. The non-local information is

causal of affects at large distances. It is no less important for macro-scale objects. Sheldrake (1995) proposed and others conducted experiments with dogs whereby the animals correctly anticipated their owners' departure from work to return home. He proposed other successful experiments where rats learning a new maze benefited non-locally from the experience of others that had previously learned the maze, in the total absence of classical space/time information. It is not surprising then, that human's exhibit an even wider range of reactions to non-local information. The evidence suggests that humans can perceive, cognize, and give meaning to non-local information across a range of complexity, from inanimate objects, simple organisms, animals and other humans. The existence of quantum holography provides an adequate informational structure to permit a theory for the observed results. The case is a classic case in phenomenology, where results are repeatedly observed over time that fall outside the prevailing paradigm, and must await new developments in science before an explanation is forthcoming. The results for intentional effects of non-locality should be no more difficult to accept than the results for perception. The pcar relationship implies symmetry, that is, information flows in both directions between object and percipient such that each is object and each is percipient. Only the complexity of the more ordered perceiving system suggests a non-symmetrical relationship. We humans have great difficulty in accepting that thoughts, specifically intentionality, can cause action at a distance. Yet, it has been observed for centuries and in recent decades subjected to scientific scrutiny. Were not prayer to have produced some positive results, religion would have been abandoned centuries ago. That cause was ascribed to supernatural agency rather than non-locality is simply, again, phenomenology needing to wait while science caught up. Modern studies by Dossey (1993), Byrd (1988) plus many others have attempted to document the efficacy of prayer, particularly healing prayer. The results in most cases are very suggestive of non-local effects, and some claim they establish the case for healing prayer. However, the difficulties of controlling all variables in such clinical studies leave many avenues for valid criticism. The fact that Radin's several studies (1997) demonstrated that attention alone produced non-local results in machines, i.e., reduced randomness (increases order) does confirm that information has non-local effect and may be correctly formulated as negentropy. These results apply directly to healing prayer as well. The case for pcar conditions to create remote effects by transfer of non-local information between equally complex percipients, humans for example, is not difficult to understand. Indeed, hundreds of successful experiments establish the case. In these cases no energy transfer is required, only non-local information, as each percipient/object has access to its own energy source. The case for intentionally creating remote physical effects in inanimate objects is more puzzling. Even though teleportation of quantum states has been successfully accomplished for particles, and numerous studies (Radin, 1997; Dunne and Dunne, 1988) show that macro-scale objects can also be changed or moved, the energy transfer mechanism by which the classical states of a remote object are affected remains elusive. Conclusions The case for mind/mind and mind/matter interactions is impressively well documented over many decades as studies in phenomenology, with staggering probabilities against chance having produced the results. The discovery of the non-local quantum hologram, which is theoretically sound and experimentally validated in at least one application, the fMRI, is sufficient to postulate that the quantum hologram is a solution to the foregoing enigma. Further, recognition that the quantum hologram is a macro-scale, non-local, information structure described by the standard formalism of quantum mechanics extends quantum mechanics to all physical objects including DNA molecules, organic cells, organs, brains and bodies. The discovery of a solution which seems to resolve so many phenomena, and also that points to the fact that in many instances classical theory is incomplete without including the subtle non-local components involved, suggests a major paradigm change must be forthcoming. The papers already published by Marcer and Schempp proposing a learning model both for DNA and prokaryote cells, which uses quantum holography, suggests that evolution in general is driven by a learning feedback loop with the environment, rather than by random mutations. This solution to biological evolution was proposed by Lamarck in 1809 but discarded for the mechanistic solution of random mutations by the colleagues of Darwin. The fact that non-local correlations and non-local quantum information can now be seen as ubiquitous in nature leads to the conclusions that the **quantum hologram can properly be labeled as "nature's mind"** and that the intuitive function we label in humans as the "sixth sense" should properly be called the "first sense". The perception of non-local information certainly preceded and helped to shape, through learning feedback, the sensory systems that evolved in planetary environments, and which we currently label as the five normal senses. We must conclude that evolved, complex organisms, which can

form intent can produce and often **do produce non-local causal effects associated with that intent.** Further, that attention alone produces coherence in nature that in some measure reduces randomness. Finally, I conclude that the cited experiments and current understanding of non-locality innature is sufficient to postulate that **non-locality is the antecedent attribute of energy and matter which permits perception and is the root of the consciousness which manifests in the evolved organisms existing in three dimensional realities.** References Anandan, J. "The Geometric Phase"; Nature, 360,26, 307-313 (1992)Berry, M. V. "The Geometric Phase" Scientific American, December, 26-32 (1988)Byrd, R. C. "Positive therapeutic effect of intercessory prayer in a coronary care population "Southern Medical Journal 81 (7): 826-829 (1988)Dossey, L. "Healing Words" Harper San Francisco, (1993)Dunne, B. J., Nelson, R. D., and Jahn, R. G. "Operator-related anomalies in a random mechanical cascade" JSE 2:155-80 (1988)Frieden, B, Roy "Physics from Fisher Information", Cambridge University Press (1998) Haisch, B., Rueda, A., Puthoff, H.E., "Physics of the Zero-Point Field: Implications for Inertia, Gravity and Mass". Speculations in Science and Technology 20 99-114 (1997) Haisch, B., Rueda, A., Puthoff, H.E. "Advances in the Proposed Zero-Point Field Theory of Inertia", 34th AIAA Joint Propulsion Conference, paper AIAA 98 3143. (1998) Hammeroff, S.R. "Quantum coherence in microtubules: A neural basis for emergent consciousness?" Journal of Consciousness Studies 1:91-118 (1994) Marcer, P. J. "Getting Quantum Theory off the Rocks: Nature as we consciously perceive it, is quantum reality! Proc. 14 International Congress of Cybernetics, Namur, Aug 21-25 (1995)Marcer, P. J. & Schempp, W. "A mathematically specified template for DNA and the genetic code in terms of the physically realizable processes of quantum holography", Proc. The Greenwich Symposium on Living Computers, eds. Fedorec, AM, Marcer, PJ, pp. 45-62 (1996)Marcer, P. J. & Schempp, W. "Model of the Neuron Working by Quantum Holography "Informatica 21:519-534 (1997)Marcer, P. J. & Schempp W. The model of the prokaryote cell as an anticipatory system working by quantum holography, Proc. 1st Intern. Conf. Computing Anticipatory System (1997)Marcer, P. J. "The Jigsaw, the Elephant and the Lighthouse", Proceedings of ANPA 20, (1998) in press .Marcer P. J. & Schempp, W. "The brain as a conscious system" Inter Journal of General Systems,(1998)Mitchell, E. D. & Williams D. "The Way of the Explorer" Putnam, NY (1996)Puthoff, H. E. & Targ, R. "A perceptual channel for information transfer over kilometer distances: Historical perspective and recent research." Proceedings of the IEEE 64: 329-354 (1976)Puthoff, H. E., "CIA initiated remote viewing program at Stanford Research Institute" JSE 10:63-76 (1996)Radin, Dean; "The Conscious Universe", Harper, San Francisco (1997)Resta, R. "The Berry Phase", Europhysics News, 28, 19 (1997)Sheldrake, R. "A New Science of Life: the Hypothesis of Formative Causation", Tarcher, Los Angeles, (1981) Sheldrake, R. "Seven Experiments That Could Change the World", Riverhead Books, NY (1995)Schempp, W. "Harmonic analysis on the Heisenberg group with application in signal theory", Pitman Research Notes in Mathematics, Series 14, Longman Scientific and Technica, London(1986)Schempp, W. "Quantum Holography and Neurocomputer Architectures", Journal of Mathematical Imaging and Vision 2: 109-164 (1992)Schlitz, M. & Wiserman, R. "Experimenter Effects and the Remote Detection of Staring", Journalof Parapsychology Vol. 61, September, (1997)Schmeidler, G. R., Craig, J. G. "Moods and Esp Scores in Group Testing" Journal of ASPR vol66no. 3, pp. 280-287 (1972)Tiller, A. W. "Science and Human Transformation" Pavoir Publishing, California, (1997)Utts, J. M. "Replication and meta-analysis in parapsychology" Statistical Science 6: 363-382,(1991)Van Flandern, T. "What the Global Positioning System tells us about Relativity", Open Questions in Relativistic Physics, ed. by F. Selleri, Apeiton, Montreal (1998) http://www.cosmicdreaming.com/pdf2011/Nature's%20Mind%20the%20Quantum%20Hologram%20by%20Edgar%20Mitchell,%20Ph_D.pdf. Models given in one of the papers. Some sentences are deleted and reformulated for felicity of reading and spatial restrictions. Kindly pardon me on that count. National Institute for Discovery Science: Nature's Mind: the Quantum Hologram by Edgar Mitchell, Ph.D. National Institute for Discovery Science Home > Resources > Consciousness Studies Nature's Mind: the Quantum Hologram Edgar Mitchell, Sc.D. Institute of Noetic Sciences, Sausalito, Calif. Fax: 561-641-5242, edgarmitchell@msn.comIntroduction My Answer To Quora Question "If Consciousness Has No Evolutionary Advantage, Doesn't That Imply That It Is An Emergent Property? " **While Human Consciousness Has Certainly Been Shaped By Evolution, Which Does Not Mean That Consciousness Itself Could Have Evolved From Non-Consciousness. Whether We Are Talking About Other Species Of Animals Or Cells Or Organic Molecules, The Same Issues Which We Run Into In Explaining Human Consciousness Are Still Present At Any Scale.** The Issues

Of The Hard Problem Of Consciousness, Explanatory Gap, Binding Problem, And Symbol Grounding Problem Make The Mind-Body Split Just As Relevant With The 'Body' Is A Brain, Neuron, Or Subatomic Particle. No Matter What, You Have To Explain How An 'Interior World' Can 'Exist' In A Physical Structure Whose Behavior Is Causally Closed. Whatever Way You Slice It, If We Accept That T-Cells Can Be Effective In Detecting And Neutralizing Threats On A Cellular Level Without Having Consciousness, Or That DNA Can Create Cellular Machines Which Build A Brain Without Consciousness, And Then We Are Admitting That Consciousness Doesn't Make Sense As A Functional Adaptation. The Rest Of The Universe Already Works Too Well Without It. There Is Nothing Especially Interesting About A Hominid's Need For Food And Shelter Which Would Demand Rich Awareness To Develop Out Of Blind Reflex. Single Celled Organisms Chase Food, Avoid Danger, Etc Also. We Are Then Left With Considering That Either Consciousness Could Somehow Be An Accident Of Evolution, Or That Consciousness May Be Intrinsic To All Physical Phenomena In Some Sense (Panpsychism, Panexperientialism) Or Even That Consciousness Is The Universal Substrate Upon Which All Phenomena Depends (Idealism, Idealist Monism). If Consciousness Is A Mutation That Has No Functional Role (A Spandrel), We Have To Ask Why It Would Even Be A Possibility. Remember That If Consciousness Is A Mutation, We Are Assuming That There Is A Whole Universe Already In Place Which Is Overflowing With Processes, Biological And Otherwise, Which Are Perfectly Capable Of Directing Themselves Effectively While Being Unconscious. It's Actually A Radically Anthropocentric Cosmology Since We Are Privileging Our Tiny Piece Of History In The Universe As The Only Piece Which Is Not Devoid Of Experience. We Are Saying That Everything That Existed Before Humans Was Unconscious, Therefore An Invisible, Intangible, Silent Void With No Memory Etc. If We Are Not Intending That, And Prefer To Think That The Universe Looked, Sounded, Felt, And Tasted Just Like It Does For Homo Sapiens Since The Dawn Of Time, Then We Would Have To Ask Exactly What We Think Consciousness Is Adding To That Kind Of Eternal-Universal 'Unconsciousness'. If Consciousness Is Intrinsic To Physical Phenomena (As In Penrose-Hameroff's Microtubule-Based Quantum Consciousness) Or Is Intrinsic To Information Integration (As In Tononi-Koch's IIT), We Still Have The Same Kind Of Mind-Body Problem. A 'Body' Which Is A Statistical Function Rather Than A Literal Form In Space Is Still Falls Short Of Explaining Why And How There Is Any Such Thing As Consciousness. In My View, Only The Idealist Monist View Or What I Call Pansensitivity Makes Sense Ultimately As The Parent Of Both Physics And Information. Just As We Learn To Count On Our Fingers, All Forms Of Information Are Representations Of Experiences Which Have An Aesthetic Foundation – A Seeing, Feeling, Touching, Thinking, Etc. Without That Sensory-Motive Context From The Start, There Would Nothing To Evolve; Only Abstractions In The Dark (Or Not Even Dark). Once We Can Get Over Ourselves As A Species And Recognize That Consciousness Doesn't Begin And End With Us, I Think That Awareness Will Be Seen As The Container Of Relativity Itself, With Quantum Mechanics And Evolutionary Biology As A Consequence Of Deeper Stories Rather Than Their Originator. Multisense Realism A Cosmology Of Sense, Essence & Existence For Models See One Of The Papers In The Series Procrastinated Due To Constraints On Space. Events Occur In Time, But There Is No Time Except That Which Has Been Produced By The Big Bang. It Makes No Difference Whether The Big Bang Is The Only Beginning Of The Only Universe Or Just The Beginning Of One Of Many Universes, Either Way It Doesn't Explain The Beginning Of Existence. Some Say That Everything Following The Big Bang Is Random, Not Designed, But Why Is 'Designed' Even An Option? If That Word Means Anything, And Anything Has Ever Been Designed By Anything, Then What Difference Does It Make Whether That Capacity Of Intention Came Early On In The Universe Or More Recently? How Do We Know The Difference Between What We Design And What Is Random? How Do We Know That Randomness Even Exists? We Can't Generate True Randomness Computationally, We Can Only Grab Onto Some Pattern That Seems Nearly Random By Some Arbitrary Duration Of Measurement And Say That It Is Good Enough. Random Is A Concept, And The Difference Between Random And Intentional May Ultimately A Matter Of Perspective. Some Interesting Possibilities Arise When We Consider The Observation That The More Something Seems Intentional, The More It Is 'Like Me' And The More Unlike Me Something Appears, The More It Seems Mechanistic. In A Way That Is Similar To How Any Object Appears As A Dot Or Smudge If It Is Too Small Or Distant For Us To See, The Notion That Perceptual Relativity Dictates The Quality Of Intention And Unintention Is A Provocative Hypothesis Of MSR. Random Events Cannot Follow A Series Or Have Results. That Doesn't Mean That There Has To Be An Entity Making Everything Happen, But It

Suggests That The Universe Is A Phenomenon Of Appearance, And Part Of Appearance Is Oscillation Between Intentional And Unintentional Attributes. The Universe Speaks In Both Entropy And Significance. Randomness Doesn't Do Anything, Can't Be Anywhere Or Feel Anything. Randomness Is An Abstraction Derived From An Expectation Of The Contrary. By Random, All We Mean Is That It Lacks Pattern And Intent – Both Of Which Must Be Implicitly Present Before They Can Be Hypothetically Absent. **There Is Sense And There Is Non-Sense, Order And Dis-Order, It Is Not Non-Randomness And Dis-Chaos. Many People Believe That Physical Law Excludes The Possibility Of Free Will Because Of Strong Causal Closure.** By This, What Is Meant Is That There Is No Room In Physics For Any Force That Causes A Physical Effect That Has Not Been Accounted For, Whether Or Not This Is Actually True Or Not Is Questionable In The First Place. The Sudden Addition Of Dark Matter And Dark Energy In The 1990s, Which Together Must Account For 95.1% Of Mass-Energy Of The Universe, Doesn't Seem To Have Encountered Nearly As Much Resistance From The Scientific Community Than Conscious Intention Has. Even If It Were The Case, However, And The Typing And Reading Of These Words Will Be Someday Explained By Purely Neurochemical Mechanisms, The Question Of Why The Feeling Of Intention Exists, And How It Can Be Produced Will Remain Unanswered. My Free Will Demands That Your Free Will Admit That There Is No Free Will. In All Of The Contemporary Debates On Free Will There Seems To Be A Blind Spot, In Which Free Will Does Not Exist, Except Where It Concerns The Application Of The Results Of Scientific Experiments. There Is Always A Call For The Educated And Enlightened To Voluntarily Change Their Own Minds, To Be Persuaded By The Argument Of Their Own Free Will. The Hypocrisy Is Hard To Overlook Once You See It. It Seems That The Anti-Free Will Assertion Makes An Exception For People Who Are 'Right'. Being Right Seems To Give Us A Right To Have Opinions Which Others Cannot, Due To Their Enslavement To Physical Causes. An Issue Which Often Comes Up In Free Will Debates Is How Our Stance On Free Will Impacts Criminal Prosecution. Again, The Blind Spot Of The Anti-Free Will Philosopher Projects That While Criminals May Not Be Held Liable For Their Actions Because Their Neurology Is To Blame, Then We Can't Allow Society To Take Credit Or Blame For Its System Of Justice Either. If Guilt Or Innocence Is Irrational, Then Believing That We Can Modify Our Own Attitudes Toward Guilt And Innocence Must Also Be Irrational. What Seems More Irrational Is To Codify Law Into A Mechanistic Formula Which Diminishes Our Capacity For Thought And Feeling In Consideration Of The **Fate Of Others (Or Others Considering Our Fate).** Zero-Tolerance Determinism: Another Pillar Of The Anti-Free Will Position Is That It Must Be An All-Or-Nothing Phenomenon. This Seems To Be More Of A Hasty Generalization Than An Honest Look At The Phenomena. In A More Tolerant Analysis, It Should Be Clear That Free Will Doesn't Have To Be Absolutely Free. The Fact That Our Conscious Mind Thinks That It Was Presented With Certain Options Still Requires That We Choose Freely From Among Those Options. It's True That We Are Always, In A Sense, Only Able To Choose What Our Mind Thinks Is The 'Best' Option (Even If That Choice Is Based On Bad Advice From Our Stomach Or Emotions). But Still There Is An Intentional Participation Which Cannot Be Explained By Statistical Functions. No Matter How Constrained We Are By The Rules Of The Road, Our Car, The Limits Of Our Driving Skill, Etc, There Is Still A Difference Between Driving And Being Asleep At The Wheel, Or Between Driving And Being Forced To Drive Somewhere At Gunpoint. We Should Not Look Only At How Our Freedom Seems To Vanish Upon Thorough Inspection, We Must Also Look At How It Appears In The First Place – Unbidden And Self-Evident. Look At The Universal Appeal Of Freedom And The Universal Disparagement Of That Which Is Unfeeling, Robotic, And Mechanistic. One Thing That Is Seldom Mentioned In These Endless Debates Is Creativity. Were The Egyptians Destined To Build Pyramids And Not Megalithic Circles? Was It Inevitable That The Star Spangled Banner Was Written In Reference To The United States? Did Deterministic Processes Have No Choice But To Create The Illusion Of Free Will? Why? The Work Of Benjamin Libet Is Frequently (Compulsively?) Cited As Well, Work Which Even Libet Himself Later Made Clear Did Not Show That Free Will Didn't Exist. The Fact That A Neurological Signal Can Be Detected Before The Various Parts Of Us (The Self Who Makes The Decision, The Self Who Knows They Make The Decision, The Self Who Reports That They Make The Decision) Can Arrive At A Consensus Does Not Mean **That The Initial Impulse Doesn't Correspond To Conscious Experience. There Is Also The Matter Of Focusing On Repetitive, Reflex Actions Which Minimize Free Will And Maximize Predictive Expectation. These Kinds Of Experiments Are Like Proving That Chefs Lack Imagination By Studying The Fry Cooks At Mcdonalds. Since Physics And Neuroscience Has No Theory At All As To The Origin Or Utility Of**

Consciousness, We Cannot Give Inanimate Instruments The Benefit Of The Doubt When It Comes To Our Subjectivity. Because We Are A Single Zygote Which Has Reproduced Itself, In Some Sense, Every Cell In Our Body Is ‘Us’ Just As Much As Any Organ Or Process In Our Body. We Are Complex, But Not In The Way That A Machine Is Complex. We Are Not Assembled From Specialized Parts; We Are A Single Whole, Divided Into Relative Specialization. Just Because Every Part Of Us Doesn’t Know What Every Other Part Of Us Is Doing At All Times, Doesn’t Mean That It Isn’t All ‘Us’ Doing It Pansensitive Relativity: If The Perceptual Relativity Hypothesis Is Correct, And Pansensitivity Is The Engine Of Both Physics And Subjectivity, Then Cause Itself Is Relativistic. Intention Is Real And Primitively Creative In The First Person, And Unreal/Recombinant-Derivative In The Third Person...But Third Person Is Only Real Because There Is A First Person Participant Present. **Free Will, Determinism, And The Big Bang: Multi Realism Blog: Wikipedia** <http://multisenserealism.com/2012/08/27/deleuzes-the-logic-of-sense-part-i/>. From A Whiteheadian Perspective, However, There Are At Least Two Key Sources Of Misunderstanding Latent In Some Of These “Returns” To “The Body”, “The Object” And “Materiality”. There Is A Rather Too Hasty Dismissal Of The Concept Of Subjectivity As Such And There Is A Related Tendency To “Flatten Out” Any Would-Be Distinctions Between Human And Non-Human Entities. Such Positions Thus Risk A Return To A Bleak Anti-Subjectivism That Mocks Those Who Might Cling To The Idea That “Humans Are Very Different From Knives Or Paper” (Harman, 2002, Cited In Thrift, 2008). With Respect To The Latter, For Example, Nigel Thrift Invokes The ANT Principle Of The Democracy Of Things And States That In His Theory “Things [And Human Beings] Are Given Equal Weight, And I Do Mean Equal” (Thrift, 2008, P. 9). Much Of This Is Done In The Name Of A Whiteheadian Inspiration Since The Book Begins With Strong References To Whitehead, Including Key Concepts Such As The “Actual Occasion”. The Point Concerning The Dismissal Of Subjectivity Can Be Brought Out Most Starkly By Contrasting One Of Thrift's Statements About His Non-Representational Theory With A Statement From Whitehead Himself. “Thus Things”, Writes Thrift, “Are Not Just Bound By Their Brute Efficacy To The Visible Termini Of Humans In Some Form Of Latent Subjectivism Such As ‘Concern’ Or ‘Care’”. Now, It Is Certainly True That For Whitehead A “Thing” Is, By Definition, “Describable Without Reference To Its Entertainment” In An Occasion Of Experience (Whitehead, 1933/1935, P. 226). In This Sense, Things Are Not “Bound By Their Brute Efficacy To The Visible Termini Of Humans”. But It Is Certainly Problematic To Imply That His Work Is Based Upon A Move Away From The Subject–Object Relation As The Fundamental Structure Of Experience And Hence From Concepts Such As Concern And Care. On The Contrary, On This Matter Whitehead's Thinking Is Quite Comparable To That Of His Contemporary Martin Heidegger In Being Grounded In The Concept Of Concern. The Centrality Of This Concept Is Due To The Way In Which It Brings Together Subject And Object As Relative Terms In The Unity Of What He Calls An Actual Occasion Of Experience: Thus The Quaker Word “Concern”, Divested Of Any Suggestion Of Knowledge, Is More Fitted To Suggest This Fundamental Structure. The Occasion As Subject Has A “Concern” For The Object. And The “Concern” At Once Places The Object As A Component In The Experience Of The Subject, With An Affective Tone Drawn From This Object And Directed Towards It. With This Interpretation, The Subject–Object Relation Is The Fundamental Structure Of Experience. (Whitehead, 1933/1935, P. 226) Granted, Thrift Does State That He Wishes To “Temper... The More Extreme Manifestations Of This Lineage, Which Can End Up By Positing A Continuity Of And To Experience About Which I Am Sceptical” (2008, P. 6) However, One Wonders What Is Left Of Whiteheadian Cosmology If This Radical Extension Of The Concept Of Experience And Hence Of Empiricism Is Omitted. It Is A Little Like Being A Marxist Without The Dialectical Materialism. It Is Also Difficult Not To Be Sceptical About The Remaining Self-Consciously “Inhuman” Theoretical Framework “In Which Individuals Are Generally Understood As Effects Of The Events To Which Their Body Parts (Broadly Understood) Respond And In Which They Participate” (Thrift, 2008, P. 60). In Recent “Radical” Social Theory, It Seems The Baby Of Subjectivity Is At Risk Of Being Thrown Out With The Bathwater Of Representationalism, Leaving Only The Hollow Remainder Of A Reactive Ensemble Of “Body Parts”. Some Clarity Is Therefore Needed If We Are Not Simply Going To Recruit Whitehead Into Concerns Alien To His Own. His Chief Problem Was Not The Notion Of The Subject/Object Structure Of Experience Itself, But Its Too Rapid Identification With The Difference Between Knower And Known (Whitehead, 1933/1935, P. 225). It Is This Conflation Of The Deeper Subject/Object Relation With The More Superficial Distinction Between Knower And Known That Gives Rise To A



“Representational” Style Of Thinking And Its Interminable Debates. I Will Argue That It Is Precisely By Way Of The Subject/Object Relation Of The Actual Occasion That We Are Best Positioned To Conceive Of “The Cumulation Of The Universe And Not A Stage-Play About It” (Whitehead, 1927–1928/1985, P. 237). **Palgrave Communications | Palgrave Macmillan Journals Journal Home > Archive > Original Articles > Full Text Original Article Subjectivity (2008) 22, 90–109 Doi:10.1057/Sub.2008.4 A.N. Whitehead And Subjectivity Paul Stenner**¹ University Of Brighton, Brighton, UK Correspondence: Paul Stenner, School Of Applied Social Science, University Of Brighton, Falmer Campus, Mayfield House, Brighton BN9 1PH, UK. E-Mail: P.Stenner@Brighton.Ac.Uk (For Models Please See One Of The Papers In The Series Attributed To Spatial Restraints It Is Accommodated Where It Could Be)

[illegible]

Poems:

And indeed there will be time
For the yellow smoke that slides along the street,
Rubbing its back upon the window-panes;

There will be time, there will be time

To prepare a face to meet the faces that you meet;
There will be time to murder and create,
And time for all the works and days of hands
That lift and drop a question on your plate;
Time for you and time for me,
And time yet for a hundred indecisions,
And for a hundred visions and revisions,
Before the taking of a toast and tea.

In the room the women come and go
Talking of Michelangelo.

And indeed there will be time
To wonder, “Do I dare?” and, “Do I dare?”

Time to turn back and descend the stair,
With a bald spot in the middle of my hair —
(They will say: “How his hair is growing thin!”)
My morning coat, my collar mounting firmly to the chin,
My necktie rich and modest, but asserted by a simple pin —
(They will say: “But how his arms and legs are thin!”)

Do I dare

Disturb the universe?

In a minute there is time

For decisions and revisions which a minute will reverse.

For I have known them all already, known them all:

Have known the evenings, mornings, afternoons,

I have measured out my life with coffee spoons;

I know the voices dying with a dying fall

Beneath the music from a farther room.

So how should I presume?

Song of Alfred Prufrock, T.S.Eliot

A Dialogue of Self and Soul William Butler Yeats

I

My Soul

I summon to the winding ancient stair;
Set all your mind upon the steep ascent,
Upon the broken, crumbling battlement,
Upon the breathless starlit air,
"Upon the star that marks the hidden pole;

Fix every wandering thought upon

That quarter where all thought is done:

Who can distinguish darkness from the soul?

My Self

The consecrates blade upon my knees

Is Sato's ancient blade, still as it was

Still razor-keen, still like a looking-glass

Unspotted by the centuries;

That flowering, silken, old embroidery, torn

From some court-lady's dress and round

The wodden scabbard bound and wound

Can, tattered, still protect, faded adorn

My Soul

Why should the imagination of a man

Long past his prime remember things that are

Emblematical of love and war

Think of ancestral night that can,

If but imagination scorn the earth

And intellect is wandering

To this and that and t'other thing,

Deliver from the crime of death and birth.

Myself Montashigi, third of his family, fashioned it

Five hundred years ago, about it lie

Flowers from I know not what embroidery —

Heart's purple — and all these I set

For emblems of the day against the tower

Emblematical of the night,

And claim as by a soldier's right

A charter to commit the crime once more

My Soul

Such fullness in that quarter overflows

And falls into the basin of the mind

That man is stricken deaf and dumb and blind,

For intellect no longer knows

I, Is from the I, Ought, or I knower from the I Known —

That is to say, ascends to Heaven;

Only the dead can be forgiven;

But when I think of that my tongue's a stone.

II

My Self

A living man is blind and drinks his drop.

What matter if the ditches are impure?

What matter if I live it all once more?

Endure that toil of growing up;

The ignominy of boyhood; the distress

Of boyhood changing into man;

The unfinished man and his pain

Brought face to face with his own clumsiness;

The finished man among his enemies? —

How in the name of Heaven can he escape

That defiling and disfigured shape

The mirror of malicious eyes

Casts upon his eyes until at last

He thinks that shape must be his shape?

And what's the good of an escape

If honour find him in the wintry blast?

I am content to live it all again

And yet again, if it be life to pitch

Into the frog-spawn of a blind man's ditch,

A blind man battering blind men;

Or into that most fecund ditch of all,

The folly that man does

Or must suffer, if he woos

A proud woman not kindred of his soul

I am content to follow to its source

Every event in action or in thought;

Measure the lot; forgive myself the lot!

When such as I cast out remorse

So great a sweetness flows into the breast

We must laugh and we must sing,

We are blest by everything,

Everything we look upon is blest.

@@
@@
@@

Dairy Notes: In excited reveries: 9th December 2014 11.47 PM.....

thrilling excitement ,voluptuous melancholy, pure joyousness, serene exaltation, thrilling excitement, stirring heroic strenuousness, weirdness and mystery, imperious tragic grandeur, contingency enterprise, endangerment enterprise, jeopardizing adventure, jejune jeremiad, avoidable inertia, latent passiveness.....
Phenomenologists (especially with regard to its ethical implications)—the phenomenon of virtualization or virtuality
The term 'virtuality' is used here to refer to the mediation of interaction through an electronic medium between humans and humans as well as between humans and machines. The Internet is the most evident example of the

virtualization of interaction. The proponents of the virtualization of society (and its institutions) argue that virtuality extends the social in unprecedented ways (Fernback 1997, Rheingold 1993a, 1993b, Turkle 1995, 1996, Benedikt 1991, Horn 1998). They argue that it opens up an entirely new domain of social being. For example Rheingold (1993a) argues that it offers “tools for facilitating all the various ways people have discovered to divide and communicate, group and subgroup and regroup, include and exclude, select and elect. Turkle suggests that cyberspace makes possible the construction of an identity that is so fluid and multiple that it strains the very limits of the notion [of authenticity]. People become masters of self-presentation and self-creation. There is an unparalleled opportunity to play with one's identity and to ‘try out’ new ones. The very notion of an inner, ‘true self’ is called into question. An individual can literally decide to be who they wish to be. For example, the obese can be slender, the beautiful can be plain and the ‘nerdy’ can be elegant. The claims by Rheingold, Turkle and others are certainly bold. What about Platonism with Respect to Time — why would someone endorse that view? One reason is that the empty container metaphor has a lot of intuitive appeal. (This is no doubt true of both the temporal and spatial versions of Platonism.) And another reason is that some people do not find the main arguments against Platonism with Respect to Time compelling. For example, it has been suggested by Sydney Shoemaker that there are possible circumstances in which it would make perfect sense to posit periods of empty time, and even to claim to know just how long those periods are. **Reductionism and Platonism with Respect to Time (Mc Garette’s Theories Source: Stanford Encyclopedia.** That makes an insinuation and innuendo that whatever happens in space time affects time too. In fact that is what relationalist philosophy says which hawking mentions many a time. Questions About The Topology Of Time Appear To Be Closely Connected To The Issue Of Platonism Versus Reductionism With Respect To Time. For If Reductionism Is True, And Then It Seems Likely That Time's Topological Features Will Depend On Contingent Facts About The Relations Among Things And Events In The World, Whereas If Platonism Is True, So That Time Exists Independently Of Whatever Is In Time, Then Time Will Presumably Have Its Topological Properties As A Matter Of Necessity. But Even If We Assume That Platonism Is True, It's Not Clear Just What Topological Properties Should Be Attributed To Time. Reductionism And Platonism With Respect To Time (Mc Garette’s Theories Source: Stanford Encyclopedia).....every one seems to be island for themselves.....how fiercely they defend their theories..... master- expert, proficient-next, master commander, chieftain head, padrone commandant, sirdar senior, lord captain, sachem governor, lord paramount, chief sheik, ruler brilliant, champion distinguished, excellent expert, first-rate great, master outstanding, superb virtuoso, ace champion, genius master, pro star, virtuoso winner, wizard arch, principal- superior, accomplished champion, chief consummate, expert finished, first foremost.....fulminating avengers with crackling chemistry, seething intensity of a volcanic protagonist.....progressively demonstrate and pathologically apprehensive.....It seems mind has settled for a while..... having seen the tireless efforts to create an impression on impressionistic minds mind moves on with the same bumbling foolishness.....or is it a point of no return?deeds, actions, reactions...and more significantly reactionary potentialities.....tutored associations have become pathological.....there must be some difference between particle and anti particle.....manifested actions of latent subjective feelings are also of the wave form..... rage.....hate.....love.....passion.....speculative domains of mind indeed.....why do they not think of a particle going to a anti particle bath room.....things like abstract ideas become more amicable and understandable.....consciousness suddenly raises its head with glamorous showcasing, and pirouette dancers..... is it a case of glorification and mortification.....retreat villas and pedestrianian traffic.....wholesome mind gets either suffused with joy or despondency... depending on the glorification and mortification.....magical wizardly to a non conformist.....your quarrel with the world ends with an fight within.....character is mistaken for an actor...shadow for substance.....I donot know how to make amends for own mind.....sometimes one feels that mind is not at all based on reason.....it needs a crutch and that is enough to build castles in air.....Hidden variables do exist.....they are to be used for perception to get reality....perception is not reality..... It is frightening.....creating Prakruti Kshobha or Prakruti siddhi is so easy.....same questions are raised and similar answers are give.....involuntary ones with demonic prepossessions.....one does not live every moment.....one

always needs a stimulus.....some more salt....some more pepper..... an automorphism of display of same scenario, schottisches, spectre,.....well! What is mind after all.....acting on information (individual consciousness).....romps on the screen.....constantly enlightening and profane.....flaunty. Raunchy.....gratuitous both in thematic discursive action.....it is only with the soulful poems and images on the screen I become one with myself..... state of perception, perceiver, process of perception becoming one.....back to the space time it has produced more slumberingness and somnambulism.....that is when I feel mistakes are necessary part of expansion of individual consciousness.... in fact there are no mistakes.....or there are no rights.....dark energy.....expanding universe..... dark energy doesn't only have an energy density: it also has a negative pressure with very specific properties. As that negative pressure pushes outwards on space, it does negative work on the Universe, and the work it does is exactly equal to the increased mass/energy of whatever patch of space you're looking at (Ethan's blog).....pure consciousnessthere has to be impure consciousness.....well! That is what you are in!..... conjunction-contemporaneousness, harmony, order, peace, simultaneousness- synchronicity exciting optimism of youth, power obmbudsman,pulsations and throbbing of pop, hard luck stories of people, modernity and mischief, grimalkins of rurality, wobbling boats of lovers,.....seamlessly think of perception, reality, dreams, dreamlessness, turiya.....may be that is the abstract of all in nature's general ledger..... Analytical clarification, Explanatory judgement, Meaningful perception ,Readings and translations, Apprehension and assimilation, Interpretational solution, Definitional rendering , Interpretational rendition, Explanatory rationale, Renditional explication, Translational co significance, Exegetical exponible, Literal epexegetic, explanatory expository, metaphrastic equivalent.....all bogs me down..... Somewhere along their line of development they diverge and go off in different pathsthere seems to be no end for search..... or does the search itself ends..... Character existenceIdentification integrity, Personality- Status, Circumstantial coherence, Distinctiveness oneness, Particularity self, Singleness singularity, Uniqueness selfhood, Identity-convertibility, Oneself hom ousia, Equality identification, ipsissima verba (exactness),coincidence monotony, tautology (repetition).....where is the question of closing when you have not opened at all.....where is question of losing when you have not gained at all.....

What are doing?

Nothing?

Why?

It is not coming!

Crow is always a crow, whether it sits here or on top of tallest building in the world!

OK!

You called me a crow!

Much worse! You are lucubration cormorant, insectivorous vespertillion wormy verkrampte!

Some where I read that mass of hydrogen strand + mass of oxygen strand is equivalent to half the mass of the matter purported to be missing!

There is invisible Byronic matter the protons and neutrons that make up atoms!

It is not Byronic you fellow; It is baryonic!

Understand what I mean not what I say!

You are not Feynmann!

School girl knows it!

Listen!

I am fed up!

Can you not say that temperature of dark matter is getting reduced at the cost of temperature of the web made of strands of oxygen and hydrogen!

Yes!

There is a statutory warning which showed that tobacco kills!

OK!

Professor Gnanendra Prabhu a dear friend died of throat cancer!

Yes! I know that!

What is the insinuation?

Nothing!

Somehow I feel hurt!

Gratification-deprivation is always non conservative!

Do **not associate things the way** you are doing!

You keep quite when I associate temperature of unidentified dark matter with that of the temperatures of web made out of strands of oxygen and hydrogen!

You are not emotionally associating it!

What do you mean?

You do it as a matter of fact!

So on one hand you work as a machine apparatus-appliance, automobile, engine, gadget, instrument, motor, tool, vehicle, automaton, computer, contraption contrivance, implement, mechanism, robot, thingamabob, widget, accoutrement, appliance, black box, device, dingbat, doodad, doohickey, furnishings, gaff, gear, gimcrack, gimmick, gizmo, grabber, habiliments, idiot box, implement, jigger, paraphernalia, provisions, setup stuff, thingamajig, tools, utensils, whatchamacallit, buggy, bus, clunker, compact, convertible, conveyance, coupe, hatchback, heap, jalopy, jeep, junker, subcompact, contraption, contrivance, doohickey, thingamabob, thingamajig, accessory, agent, arrangement, contrivance, doohickey, equipment, expedient, material, thingamabob and on the other romanticist and idealist..... escapist, person who holds fancies in mind, who believes in perfection, dreamer, optimist, visionary, enthusiast, person who holds fancies in mind, who believes in perfection, Platonist, dreamer, enthusiast, escapist, stargazer, theorizer, transcendentalist, utopian, evader, classicist, impressionist (Roget's 21st Century Thesaurus, Third Edition Copyright © 2013 by the Philip Lief Group.)

Yes!

You also do that simultaneously!

Yes!

What guarantee is there you are not romanticizing science also..?

I do!

So Ψ in Dirac Brackets = + C (Machine) E (machine) in Dirac brackets + C (romanticist) E(romanticist) in Dirac brackets

Yes!

And you can use the ASCII computation for romanticist part.!

Possibilities of not being both?

Death like situation, where there neither romanticism not empiricism?

Yes!

So Ψ in Dirac Brackets = - C (Machine) E (machine) in Dirac brackets - C (romanticist) E(romanticist) in Dirac brackets

Yes!

So we are all gobbling up the mixture ravenously!

Yes!

Ugh! It is you in both the cases!

Subjectively objective?

Like a constrained man restraining his emotions, emotional intelligent person!

So Ψ in Dirac Brackets = - C (Machine) E (machine) in Dirac brackets + C (romanticist) E(romanticist) in Dirac brackets

Yes!

Objectively subjective?

So Ψ in Dirac Brackets = + C (Machine) E (machine) in Dirac brackets - C (romanticist) E (romanticist) in Dirac brackets

Anatol Rapoport's top dog- underdog syndrome, wherein underdog acts irrationally to get his due!

You said God is absolute subjectivity!

Yes!

You say we are objective!

There is no necessity to glorify objectification!

Why?

Because you are caught in the web of subjectivity!

Spider like web where ParaBrahaman waits to see your reactionary potentialities to systematize the punishments and punitive actions!

It is fun!

Leela!

Yes!

But why?

Show must go on!

Subject and Object were the same in the beginning!

God made man in his image!

You are stardust!

Nature is observer dependent!

Contradiction?

No!

Guru always hides an important secret from tutelage!

You are ensconced by Maya!

Infact they address most of the deities here as “maha mAyA, mahmayi”

Like mother sowing moon to a child and promising to get him for dinner!

We cannot be fools fundamentally!

I must be methinks!

Hunch back of Notre dame comes holding spectrum coloured clothes everyday when I go down the stairs!

Stop it!

Somehow the pretension is conspicuous!

Scientists are doing the same thing over and over again from time to time and time and again!

You think everything is discrete?

Yes!

How Einstein did suggest a continuum?

It was an equation!

But everyone has their own version and it is being portrayed in arrestingly captivating way!

Balls rolls over the rubber sheet which spasmodically turns and twists, warps and wefts due to the mass!

Donot you think mass of already existing matter must have curved it before to an extent that is indiscernible or taken as an initial condition?

May be!

Dangerously frightening, distressing disturbing, torturous traumatic, alarming racking, soaring tearing, tormenting torturing, heart-rending, nerve-racking, harrowing intense, racking struggling ambiguity and equivocality!

Large invisible matter exists!

Strands of half of the missing mass have been found as strands of hydrogen and oxygen!

Donot draw me in to will o' the wisp!

Strands form web like structure form the backbone of the universe!

You can apply your model now!

Can you not search for missing baryons or normal matter by some means?

I donot know!

Gamma rays!

I donot think so!

Quasars!

May be!

Why probe nature? Are not you disturbing nature?

Yes!

Ethical issues are involved!

What is nature?

"Nature "means: (Source: Wikipedia: Aristotle)

- (a) in one sense, the genesis of growing things — as would be suggested by pronouncing the ν of φύσις long—and
- (b) in another, that immanent thing from which a growing thing first begins to grow.
- (c) The source from which the primary motion in every natural object is induced in that object as such. All things are said to grow which gain increase through something else by contact and organic unity (or adhesion, as in the case of embryos). Organic unity differs from contact; for in the latter case there need be nothing except contact, but in both the things which form an organic unity there is some one and the same thing which produces, instead of mere contact, a unity which is organic, continuous and quantitative (but not qualitative). Again, "nature" means
- (d) the primary stuff, shapeless and unchangeable from its own potency, of which any natural object consists or from

which it is produced; e.g., bronze is called the "nature" of a statue and of bronze articles, and wood that of wooden ones, and similarly in all other cases. For each article consists of these "natures," the primary material persisting. It is in this sense that men call the elements of natural objects the "nature," some calling it fire, others earth or air or water, others something else similar, others some of these, and others all of them. Again in another sense "nature" means

(e) the substance of natural objects; as in the case of those who say that the "nature" is the primary composition of a thing, or as Empedocles says: Of nothing that exists is there nature, but only mixture and separation of what has been mixed; nature is but a name given to these by men. Hence as regards those things which exist or are produced by nature, although that from which they naturally are produced or exist is already present, we say that they have not their nature yet unless they have their form and shape. That which comprises both of these exists by nature; e.g. animals and their parts. And nature is both the primary matter (and this in two senses: either primary in relation to the thing, or primary in general; e.g., in bronze articles the primary matter in relation to those articles is bronze, but in general it is perhaps water—that is if all things which can be melted are water) and the form or essence, i.e. the end of the process, of generation. Indeed from this sense of "nature," by an extension of meaning, every essence in general is called "nature," because the nature of anything is a kind of essence. From what has been said, then, the primary and proper sense of "nature" is the essence of those things which contain in themselves as such a source of motion; for the matter is called "nature" because it is capable of receiving the nature, and the processes of generation and growth are called "nature" because they are motions derived from it. And nature in this sense is the source of motion in natural objects, which is somehow inherent in them, either potentially or actually.

— Metaphysics 1014b-1015a, translated by Hugh Tredennick, emphasis added

You mean Brahman experience is possible in space time?

Certainly!

It is just the identification with the object, and you becoming the object itself, may be your company, God, or another person!

Know completely. Apply in action!

OK!

It is at this stage both Terrestrial Brahman Anti Brahman and Celestial Brahman Anti Brahman meet!

Meet!

Yes if you think one is the subject and the other is the object!

But can it happen?

Man is essentially an evolutionary being and be it “good” or “bad” and tries to completely “become” role model!

No glorification or mortification please! I cannot stand!

It is just a procès of becoming! Like the identification with a film, company, or a job!

AL=GD?

Yes!

"The subjective thinker's form, the form of his communication, is his **style**. His form must be just as **manifold as are the opposites that he holds together**. The systematic eins, zwei, drei is an abstract form that also must inevitably run

into trouble whenever it is to be applied to the concrete. To the same degree as the subjective thinker is concrete, to the same degree his form must also be concretely dialectical. But just as he himself is not a poet, not an ethicist, not a dialectician, so also his form is none of these directly. His form must first and last be related to existence, and in this regard he must have at his disposal the poetic, the ethical, the dialectical, the religious. Subordinate characters, setting, etc., which belong to the well balanced character of the esthetic production, are in themselves breadth; the subjective thinker has only one setting—existence—and has **nothing to do with localities and such things**. The setting is not the fairyland of the imagination, where poetry produces consummation, nor is the setting laid in England, and historical accuracy is not a concern. The setting is inwardness in existing as a human being; the concretion is the relation of the existence-categories to one another. Historical accuracy and historical actuality are breadth." Søren Kierkegaard (Concluding Postscript, Hong p. 357–358) (Source: Wikipedia)

But to reach Turiya you have to cross the “Dwanda” the “Dvaita” or “Advaita”!

Yes!

How then can you have the general ledgers you have been ululating about!

Ugh! It is the principle of conservativeness that is emphasised! Zero is all in all!

It is question of: either” or “both” situation!

But no one dies here!

It is again the conservativeness of Assets and Liabilities and non conservativeness of Gratification and Deprivation that is embellished!

When the God-forsaken worldliness of earthly life shuts itself in complacency, the confined air develops poison, the moment gets stuck and stands still, the prospect **is lost, a need is felt for a refreshing**, enlivening breeze to cleanse the air and dispel the poisonous vapors lest we suffocate in worldliness. ... Lovingly to hope all things is the opposite of despairingly to hope nothing at all. Love hopes all things – yet is never put to shame. To relate oneself **expectantly to the possibility of the good is to hope**. To **relate oneself expectantly to the possibility of evil is to fear**. By the decision to choose hope one decides infinitely more than it seems, because it is an eternal decision. p. 246-250 (Kierkegaard: Either or: Wikipedia)

That is way of escaping reason and rationalism!

Leave that part of it! Take the quintessential part!

Aw!

But the grand design still exists with the contribution from individuals and collectivity!

Yes!

Is not that a contradiction?

Universal mind is not accepted by some people!

Do you accept?

I tend to believe that quantum information stays on before it reaches the Universal Mind and becomes one with it!

Well! That is what Geeta also says!

The universal mind appears as a Roman belief in Vergil as well:

"In the beginning, SPIRIT within strengthens Heaven and Earth,

The watery fields, and the lucid globe of Lina, and then --

Titan stars; and mind infused through the limbs

Agitates the whole mass, and mixes itself with GREAT MATTER"

(Virgil: "Aeneid, vi" 724 ff.)

Yes! Different paths! Single Truth!

Path is not the truth?

That is also there!

Path is also the destination!

Yes!

That is deductive reason!

"A truth that's told with bad intent

Beats all the lies you can invent."

— William Blake, *Auguries of Innocence*

You cannot live without a complaint!

Yes!

You have to live without high reactionary potential for issues!

Are not black holes consuming matter an issue?

Yes!

Then?

Why do you not arrest black hole?

Problem is they donot consider personalities, propositional ties and poisionalities as nature!

But they are!

So?

Live with what cannot be changed!

But there is a changeless substratum!

Faith!

Faith is then an Adhoc device given to train the mind!

You have to!

They ask you to do breathing exercises; eye exercises, and so on.....!

But then you would not know nature at all!

Training itself kills truth!

Then?

You have to live in despondency and disconcert and still not affected by it!

Sthithapragmata!

Yes!

You are serving the present purpose!

But the disease has to be checked!

Then?

You would not know where the disease would lead to!

Ask scientists!

Why?

They donot have any such constraints!

You want to have instant gratification!

Yes!

Then accept and move on!

If not?

Intuition and instinct, conservation and preservation of gratification and deprivation are quintessential!

You do it you are damned!

You donot do it! You are damned!

You start believing what others think you are!

There is no reality in that?

I donot think so!

You are then a vagabond without destination!

Who said there is one?

Path itself is destination!

That means the more charlatan you are, the more able you are to convince others, and you reap dividends!

Ethical question!

What is the Truth?

Progression of a point of view and its ultimate ramification is important!

Give reasons and take redressal measures!

Then you would not know where the 'positive' or 'negative' leads to!

Ugh! Too much!

Faith and patience are essential!

You are only aiming at self gratification and exaggeration!

Identity is essential!

But Brahman-AntiBrahman has no identity!

That is why they are one!

Dark matter, Dark energy?

May be!

You have to prove it!

Either they are one or not!

OK!

Infinite regresses can be stopped by temporarily freezing the object in question as some sort of reality in itself, not relying on a proposition. OK but only a thin sort of sense can then be extracted. As in black boxing? Or when a power relation forbids inquiry – hier IST nicht warum. We do learn something though – that actual events are largely 'sterile' when it comes to making sense (hence Deleuze's indifference to empirical inquiry?) (There is a reference to Husserl's indifference – as in the last bit of Cartesian Meditations – 'don't look outside, truth lies in the interior of mankind' etc). (LOS: Deleuze) Alexander I. Stingl's Blog A Nomadic Scholarship Entity NOTES ON: Deleuze, G (1990) The Logic Of Sense, Trans Mark Lester, Edited By Constantin Boundas, New York: Columbia University Press

Indifference arises out of either arrogance or ignorance!

Husserl or Deleuze were neither!

There are no power relations that forbid enquiry!

Yes!

Then?

May be social ostracism or societal constraints!

May be!

Explanation please!

Deleuze himself has this to say: Sense operates through a series of propositions, as the example of infinite regress shows. Quite often, the series becomes more abstract or general. The purest form, however, is where separate series are established, one involving denotation, and the other making sense. These different series operate in different ways, one at the surface, with denoted objects, and one inside propositions linking expressions, and also connecting them to denotations [pretty much like the way in which signifiers link together in propositions, and then are occasionally attached to actual referents which themselves develop sequences. Deleuze wants to develop the term signifier to mean anything which is an aspect of sense, and signified to mean that which is denoted, or realized. He wants to connect it back to the difference between events and states of affairs Notes that the signifier refers to the whole content of the proposition]. The issue is how can the series be joined? There are a number of possibilities. Mystifying literary examples follow 37-39 [I think what they are referring to is the way in which terms can both symbolise and denote literally. Clever writers such as Joyce or Robbe-Grillet have indicated this in particular ways, for example by using esoteric words at crucial points.] These writers are able to show that **there are genuine differences or displacements between the two series, and that both have their own momentum**. However the signifying series contain an excess of meaning, while the denotated features a lack of: it is this that articulates the two series and makes them make sense (Ibid)

Exemplification or illustration?

Indistinguishability!

Patience is the intuition, organisation and disorganization, and reorganization as it happens in the offices or Banks to get at the solution or the correct perspective!

Stochastic learning theory, response stimulus theory, linguistic structure theory, deductive reasoning model, non linearity of the language, social interactional dynamics.....!

Deconstruction of all these models is what could be called intuition!

Intuition and confidence in theory under whatever assumption is what makes the theory to converge!

Gravitational force can act as anti gravitational force too!

Ψ in Dirac Brackets = C (Gravitation) E (Gravitation) in Dirac brackets + C (anti gravitation) E (Anti Gravitation) in Dirac Brackets

Yes!

Brahman-Anti Brahman is responsible for the attraction of people towards me and revulsion and repulsion from me of the people!

What is said is such gravitation and antigravitation could be simulated!

How?

Shifting a bus stand!

People no more converge at that place!

Why then is the Black money –Dark energy syndrome?

With Black money AL Model does not tally, but it is only with dark energy that AL of nature tallies!

Balance sheet drawn up is correct!

That is because mass energy equivalence principle!

There is no principle that says financial energy is equivalent to the mass thereof for the inclusion of black money and then tally the AL sheet!

Black money doth expand inflate the money value. That is one of the reasons as to why all concerted efforts to reduce the lack money have remained futile!

Dark energy then seems to be responsible for the inflation of the universe!

So! So!

Aw!

All said and done space time itself is the product of the quantum information being projected by Celestial Brahman Anti Brahman on the screen of individual consciousness!

OK!

Some thing wrong with the projection!

Ugh! Continuity experienced is because of the continuity principle in projection of a film!

OK!

Are the basic forces also simulated by celestial Brahman nti Brahman?

Everything!

Then what are you seeing!

A dream!

Somehow you have the knack of getting me started from square zero every time!

Expansion is measured. Light from distant supernovae looks dimmer and redder than expected which shows that universe is expanding!

Donot associate anything! It could be misleading!

CMWB?

Yes!

But it changes!

Terrestrial Brahman Anti Brahman also changed the colours to give the subject a different perception!

Celestial Brahman Anti Brahman might not be doing that!

He is simulating!

ParaBrahman?

He is nothing! Zero!

Vacuum energy!

Yes!

Repulsion of dark energy gobbles up attraction of gravitational field which produces galaxies to drive apart!

Mentally is it possible to use to energy packet to get a Dirac an idea?

What can be done is to dissolve the “quantum clusters” with “hard” autosuggestion!

Tell the mind mildly!

Why?

It might retaliate and become recalcitrant and pathological!

Do you know you are advocating for yourself!

Yes! That is what I am, a polarised membrane!

Chit Shakti is the counterpart of mAyA Shakti!

So you are being misled?

I think we are getting everything wrong!

Remember there is no freewill to exercise in extant circumstances!

Thoroughbred information explosion and subjection thereof might constrain you accept one association after another!

Two documents open!

You must have double clicked twice!

No!

It has happened many times!

Identity less creatures’ simulating?

Do not associate every incident or accident with any thing!

So?

It happened!

Why do scientists do that!

They are on the journey of exploration!

You are on a journey to?

Imploration!

So!

It leads to implosion!

Their journey does not lead to explosion of information?

Everything is discrete!

There is an underlying substratum that encompasses all the things in the universe ,just everything!

Who told you that?

Vedas, Böhm's implicate order, Christ!

Oh! You irritable, testy, acrimonious, captious carping, caviling , childish churlish complaining crabbed cranky critical cross crotchety crust cussed fault-finding fractious fretful, fretting grouchy grousing growling grump huffy ill-natured mean morose obstinate ogre ornery, out-of-sort ,pertinacious ,petulant querulous short-tempered snappy, splenetic , sulky sullen surly tetchy touchy ,waspish, whining, acrimonious, acerbic, acid, angry, astringent, belligerent, biting, bitter, caustic, censorious, churlish, crabby, cranky, cross, cutting, indignant, irascible, irate, ireful, mad, mordant, peevish, petulant, rancorous, sarcastic, sharp, spiteful, splenetic, tart, testy, trenchant, wrathful, cantankerous, difficult, crabby, bad-tempered, bearish, captious, choleric, contrary, cranky, critical, cross, crotchety, crusty, disagreeable, dour, grouchy, grumpy, huffy, ill-humored, ill-natured, irascible, irritable, morose, obstinate, ornery, peevish, perverse, prickly, quarrelsome, snappish, sour, stuffy, testy, vinegarish, vinegary, carping, cavillous, contrary, crabby, cross, demanding, deprecating, disparaging, exacting, exceptive, fault-finding, finicky, hypercritical, irritable, nagging, nit-picking, overcritical, peevish, perverse, petulant, sarcastic, severe, testy, touchy, crabby/crabbed, awkward, blunt, brusque, captious, choleric, churlish, cranky, cross, crotchety, crusty, cynical, difficult, dour, fretful, gloomy, glum, grouchy, harsh, huffy, ill-humored, ill-tempered, irascible, irritable, misanthropic, morose, nasty-tempered, perverse, prickly, saturnine, snappish, surly, tart, testy, tough, trying, unsociable, cross, crotchety, crusty, fractious, fretful, grouchy, grumpy, ill-humored, ill-tempered, impatient, irascible, irritable, jumpy, querulous, quick-tempered, ratty, short, snappy, tetchy, touchy, vexed waspish, crotchety, cross-grained, crusty, curmudgeonly, difficult, disagreeable, eccentric, fractious, obstinate, obstreperous, odd, ornery, dour, gruff, harsh, ill-humored, scornful, short, short-tempered, snappish, snarling, snippety, snippy, bellicose, brusque, contentious, contrary, cross, difficult, disobliging, disputatious, eristic, grouchy, ill-natured, nasty, obnoxious, offensive, out of sorts, querulous, rude, snappy, ungracious, unlikable, unpleasant, uptight, waspish, whiny, disgruntled, discontented, griping, grouchy, grousing, kicking, kvetching, malcontented, demonstrative, edgy, emotional, enthusiastic, fidgety, fierce, fiery, galvanic, hasty, high-strung, hot-headed, hot-tempered, hysterical, impatient, impetuous, impulsive, inflammable, intolerant, irascible, mercurial, moody, nervous, neurotic, overzealous, passionate, peevish, quick, quick-tempered, rash, reckless, restless, sensitive, skittish, susceptible, temperamental, vehement, violent, volatile, volcanic, fractious, froward, huffy, indocile, indomitable, intractable, irritable, mean, ornery, recalcitrant, refractory, restive, scrappy, snappish, testy, thin-skinned, touchy, uncompliant. unruly, wayward, wild, contrary,, grumbling, truculent, huffy, exasperated, fractious, grumpy, huffish, moping, nettled, offended, riled, vexed, disputatious, dyspeptic, snappy, snarling, surly, brawling, tempestuous, thin-skinned, gruff, malevolent, malicious, dicey, jumpy, precarious, ticklish, tricky, wrung, bewailing, charging, critical, deploring, disapproving purulent ptomaine, sacrosuchus ray imperator! (Reference: Roger's Dictionary)

You See!

Tell me!

Our language affects how we perceive things:

Even comparatively simple acts of perception are very much more at the mercy of the social patterns called words than we might suppose. ...We see and hear and otherwise experience very largely as we do because the language habits of our community predispose certain choices of interpretation (p. 210). Supplement to Relativism The Linguistic Relativity Hypothesis (Stanford Encyclopedia)

OK! Go On!

Then language affects cognition!

Yes!

Affectation is restricted to cognition or does it have ramifications on anything else that is of importance to my study?

Cassirer tells us that:.....the distinctions which here are taken for granted, the analysis of reality in terms of things and processes, permanent and transitory aspects, objects and actions, do not precede language as a substratum of given fact, but that language itself is what initiates such articulations, and develops them in its own sphere (1946, p. 12). (Ibid)

The fact of the matter is that the 'real world' is to a large extent unconsciously built up on the language habits of the group. No two languages are ever sufficiently similar to be considered as representing the same social reality. The worlds in which different societies live are distinct worlds, not merely the same worlds with different labels attached (p. 209) (Ibid)

Ugh! What is the net result?

We are thus introduced to a *new principle of relativity*, which holds that all observers are not led by the same physical evidence to the same picture of the universe, unless their linguistic backgrounds are similar, or can in some way be calibrated. ...The *relativity of all conceptual systems*, ours included, and their dependence upon language stand revealed (1956, p. 214f, italics added).

We dissect nature along lines laid down by our native languages. The categories and types that we isolate from the world of phenomena we do not find there because they stare every observer in the face; on the contrary, the world is presented in a kaleidoscopic flux of impressions which has to be organized by our minds--and this means largely by the linguistic systems in our minds (p. 213).

That means there is no freewill to understand nature!

Yes!

Why do they work then?

Show must go on!

That means there is no objective reality only subjective experience!

Yes!

Subjectivism is the philosophical tenet that "our own mental activity is the only unquestionable fact of our experience" In other words, subjectivism is the doctrine that knowledge is merely subjective and that there is no external or objective truth. The success of this position is historically attributed to Descartes and his methodic doubt.

Subjectivism accords primacy to subjective experience as fundamental of all measure and law.[2] In extreme forms like Solipsism, it may hold that the nature and existence of every object depends solely on someone's subjective awareness of it. One may consider the qualified empiricism of George Berkeley in this context, given his reliance on God as the prime mover of human perception (Wikipedia)

Probabilities have no objective reality and Schrodinger's wave function might collapse in a singularity!

Yes!

I don't want any aggressive iconoclasts with serenading whimsicality and astute truculence!

I am telling what I felt!

What is "feeling?"

Subjective intuition!

Inspired guess?

Yes!

I felt because of the anagrammatic mind acting on individual consciousness which is information stored!

Why do get the "intuition" in the first place?

Well! I am unarmed, and look for some authority; solace so that I can be gratified!

Gratification is the main reason!

Rest?

Figment of imagination!

If you subscribe to perceptual relativism or cognitive relativism, then you say Truth is relativistic!

Yes!

That is what Dvaita says too!

Do you not think it is an act of self abnegation and revocation?

May be!

For a person in mental hospital world looks mad and he thinks he is the smartest ass going around!

So?

It is a case of social maladjustment!

Not community enlightenment?

No!

Then why use theories' to explain the behaviour?

Do not use!

No! That is escapist answer!

Madness is observer dependent?

I did not say that!

That is what is implicit!

In mathematics, more precisely in measure theory, a measure on the real line is called a discrete measure (in respect to the Lebesgue measure) if its support is at most a countable set. Note that the support need not be a discrete set. Geometrically, a discrete measure (on the real line, with respect to Lebesgue measure) is a collection of point masses. (Wikipedia)

That is the case of graph and bar charts!

Yes!

Denotation is there!

If H is a Hilbert space means just denotation and signification!

OK!

You are not bringing Hilbert space!

OK!

I will write not (something) when I don't know the answer!

Yes!

You can apply accentuation attrition models thereof?

Yes!

Then what is the problem in the concatenation of GTR and QM?

Nothing!

Why that everyone is cool as cucumber and dead as dodo?

Such things happen!

What then is the relationship between two persons say A and B?

It is like this!

Like?

Past transactions (net) of A (past transactions of B) (net) + (past positive transactions of A) (Past positive transactions of B divided by (present negative transactions of A) (present net negative transactions of B)

OK!

Gratification deprivation complex is never tallied individually but is conservative holistically! Yes! You see negative energy packets in some areas!

L>>>>>A (Bankruptcy)

D>>>>>>>>>G (madness)

So what to do to avoid madness?

Don't replicate thoughts!

But I am like that!

Rectify yourself! Exercise!

Why do they not go and sit watch and study big bangs taking place instead of LHC?

They can if they can!

What do you mean!

There is no AC room up there!

If they collect data, sign language, color language of nature?

they will be biased as much as anyone who reads them!

It changes I say!

Terrestrial Brahman Anti Brahman doth make small improvisation, notwithstanding the replication like all great scientists or artists!

After bloating do you think universe would survive?

It will burst sooner or later?

We have very small amount of knowledge of nature!

Look inward you would know the truth!

Death, damnation, disintegration is all part of life like revitalization, resurrection, rejuvenation is!

There is no way to change?

You need not!

Why?

Let posterity also get on the jamboree and bandwagon and get recognition, awards etc.,!

And get in to the same rigmarole!

Yes!

If it is repetitive do you not think everything is out of habit?

Yes!

Everything is and?

I think so!

Madness is a characteristic of space and time?

It is!

There is a legend where Krishna asks after coming in to space time ambit whether he could use a part of land belonging to Jambavantha!

Gravity is responsible for all the ills of segregating space in to zones...crime zone, silent zone, children zone,.....speed breakers, additional punitive justice.....?

I conjecture so!

Nothing specific?

You cannot see your own life or death I say!

But scientists' are seeing!

Simulation or dissimulation itself kills Truth!

Aw!

That means you are deducing the meaning out of the information stored in you! Individual consciousness!

Yes!

What then is science?

Collective or part of collective consciousness!

They have tested their theories!

What is testing!

Verification! They conduct experiment and find out the referential integrity of the theory!

Example?

Higgs Boson!

Say Celestial Brahman Anti Brahman has simulated the result!

What do you mean?

Everytime somebody conducts an experiment there will be same result!

Like whenever there is Meow! You are reminded of Schrödinger's cat or pirouette dancers!

Later is due to perilepsis baracudina doing the same thing and filling your consciousness out of it!

When atom bomb was dropped on Hiroshima and Nagasaki there was murder, mayhem , plunder, pillage, nemesis and apocalypses!

There is no denying it!

Pain is there, suffering is there!

It is because of your body consciousness and mind consciousness!

So?

You are neither body nor mind!

What?

You are neither ahamkara (ego) nor mamakara (what you think others think you are)!

There is pain!

There is neither pain nor pleasure!

That is the state of pure consciousness where all dualities cease!

But I am still in evolutionary stage!

So you suffer!

Evolution is suffering?

I spoke of stage not the process itself!

Oh! You are differentiating there too! You have a divide and rule policy!

It is clearcut ontological and phenomenological contradiction!

There is neither question nor answer!

Then?

Nothing! Buddha smiled!

I am not one!

That is why you are suffering!

OOOO.....!

**National Institute for Discovery Science: Nature's Mind: the Quantum Hologram by Edgar Mitchell, Ph.D.
Nature's Mind: the Quantum Hologram Fax: 561-641-5242, edgarmitchell@msn.com**

- (1) It takes training as provided by many of the esoteric traditions and/or certain naturally sensitive individuals to routinely perceive the non-local holographic information associated with a particular object.
- (2) There is massive evidence to suggest, however, that the brain has these latter capabilities at birth. **Suppression by cultural conditioning in childhood and subsequent lack of practice cause (eb) the natural ability for conscious, intuitive perceptions to atrophy.**
- (3) Particularly in western tradition, educational interest has been on the left brain, rational functions rather than right brain, intuitive functions. However, mystic adepts and natural psychics routinely demonstrate that non-local information is perceptible from physical objects by focusing attention, quieting the left brain and allowing intuitive perceptions to appear.
- (4) It is the left brain cognitive ability in humans that provides canonical labeling of the intuitive and artistic processes taking place in the right brain.
- (5) The fact that with training and practice, individuals can recover, deepen and label their individual cognitive access to intuitive, non-local information demonstrates that learning is taking place within the whole brain itself and involves enhanced coherence and coordination between the hemispheres. This process is different and distinct from the left brain function of extending and extrapolating factual data and

logical deduction to leap to (e&eb) an "intuitive" conclusion, while omitting the intermediate steps leading to that conclusion

OK!

Different spaces and times are coming out towards infinity!

Not even a blade of grass is cut without purpose!

Ok!

But life itself seems to be purposeless!

That is the beauty!

But lots of planning commissions, implementation offices are there!

Huh! Huh! Name sake!

Each action or transaction has a purpose towards contribution towards grand design, the cosmic general ledger!

Yes!

Like money may be a deposit or a credit!

Yes!

Why that is a virtue becomes vice and vice-versa as the time goes on!

First that is what polarisation leads to like Maxwell's demon!

Due to?

Terrestrial Brahman anti Brahman and celestial Brahman and Anti Brahman!

OK!

Is there a higher purpose or a lower purpose? No!

Good=evil!

$(\Psi) \text{ Dirac Brackets} = C(\text{good}) E(\text{good}) \text{ in Dirac brackets} + C(\text{evil}) E(\text{evil}) \text{ in Dirac brackets}$

Seems so!

This is order?

Yes!

No change necessary!

Time shall do that!

Motivation and demonization is not necessary?

No?

Somehow it sounds like Platonic philosophy, a depressive philosophy!

What ever be the appellation, nevertheless, it is the same by all measures and counts!

Could not every philosophy since Aristotle be characterized as an attempt to reverse Platonism (and not simply a footnote to Plato, as Whitehead once suggested)? 5 Plato, it is said, opposed essence to appearance, the original to the image, the sun of truth to the shadows of the cave, and to overturn Platonism would initially seem to imply a reversal of this standard relation: what languishes below in Platonism must be put on top; the **super-sensuous must be placed in the service of the sensuous**. But such an interpretation, as Heidegger showed, only leads to the quagmire of positivism, an appeal to the positum rather than the eidos. More profoundly, the phrase would seem to mean the abolition of both the world of essence and the world of appearance. Yet even this project would not be the one announced by Nietzsche; Deleuze notes that "the double objection to essences and appearance goes back to Hegel, and further still, to Kant" (LS 253). Essays on Deleuze Daniel W. Smith

Sounds quagmirish!

The path from myth to reason was not some sort of **inexplicable "miracle" or "discovery of the mind," they argue, but was conditioned historically by the social structure of the Greek polis, which "laicized" the mythic forms of thought characteristic of the neighboring empires by bringing them into the agonistic and public space of the agora**.

8 In Deleuze's terminology, imperial states and the Greek cities were types of social formations that "deterritorialized" their surrounding rural territories, but they did so according to two different models. The archaic States "overcoded" the rural territories by relating them to a superior arithmetic unity (the despot), by subordinating them to a transcendent mythic order that was imposed upon them from above. The Greek cities, by contrast, adapted the surrounding territories to a geometric extension in which the city itself became a relay-point in an immanent network of commercial and maritime circuits. These circuits formed a kind of international market on the border of the eastern empires, organized into a multiplicity of independent societies in which artisans and merchants found a freedom and mobility that the imperial states denied them (Ibid)

You mean you are a visitor and I am like a hotel, where you come and go!

Looks strange!

To whom?

Are you always with me?

Yes!

No mind condition?

Well! No more anagramatization!

What happens!

Thoughts come and go!

Nothing happens!

Nothing!

Aw!

But Aristotle interprets division as a means of dividing a genus into opposing species in order to subsume the thing being investigated under the appropriate species-hence the continuous process of specification in search for a definition of the angler's art! (Ibid)

Classification and denotation, signification, individuation seems to be the root cause of all problems!

At least Deleuze does not think so!

That should not matter!

The concept of the

Idea, in Deleuze's analysis, thus consists of three components:

I. the differential quality that is to be possessed or participated in (e.g., being just)

2. The pre-existent foundation or Idea that possesses it firsthand, as unparticipatable

(e.g., justice itself)

3- the rivals that lay claim to the quality (e.g., to be a just man) but can only possess it at a second, third, or fourth remove ... or not at all (the simulacrum) (WP 30). (Ibid)

Problem here is about the "differential quality"!

Your problem!

Not Deleuze's!

It is a question of information storing! It is you acting on that!

Derrida coins the concept of iterability in order to account for this possibility of non serious repetition. He says of the concept: "'iterability' does not signify simply ... repeatability of the same, but rather alterability of this same idealized in the singularity of an event, **for instance, in this or that speech act**" (L 119). **Iterability thus signifies, among other things, the** alterability of the concept of a promise in the singularity of an actual, occurring promise. This understanding of conceptual change is implied in earlier works by Derrida and indeed in 'Afterword ... ,' Derrida suggests that his attempt elsewhere to "think a difference which would be neither of nature nor of degree" was precisely an attempt to "think or deconstruct the concept of concept otherwise" (LII7). Thinking the Concept Otherwise: Deleuze and Expression PETER COOK

That is true! Moment you are repetitive your ego is boosted and you on terra firma!

You should be unstable?

Frivolous fiddle deddee! You start to see things!

But there is nothing!

Maya pervades!

Yes!

Then there are dual realities and virtualities?

I think so!

Accentuation coefficient and dissipation coefficient have to be used separately to get correct net result?

Yes! I think so!

Not sure?

No!

First you said what you see is not what you see! What you see is what you don't see! What you see is what you don't see! And what you don't see is what you see!

Add that to Peter Cook's quote and what do you get?

What is true is false, and what is false is true!

How can that be possible?

There are two options!

One you do not show what you are! Or you show what you are!

More often than not former reigns everywhere! Just watch all those videos! They seem to be too fastidious and fussy to impress others!

Yes! That is true!

But Derrida seems to think otherwise!

I told you it does not matter! They talk based on the storage of information they have!

Still read this on:

As regards this alteration, Derrida holds that the concept of iterability " ... entails the necessity of thinking at once both the rule and the event, concept and singularity" (L 119). For a concept must be considered not just as a general rule or definition, but rather as a general rule which is being applied to a particular and unique instance: as "this same idealized in the singularity of an event." In being applied to a particular which differs from the prior instances that constituted it, a concept effectively differs from itself. Derrida elsewhere speaks of such a concept being "Nonself-identical" (SP68). A concept alters, then, in being applied to a particular, which renders the concept no longer simply a general rule or definition. Beyond this kind of alteration, the application of a concept to a particular can in fact necessitate altering the rule of the concept itself. Derrida's criticism of the traditional concept of the concept hinges on this. Thinking the Concept Otherwise: Deleuze and Expression PETER COOK

Conceptualization itself kills truth!

Event?

It is the product of your conceptualization!

Singularities'!

Wave function itself collapses in singularity when the concepts are different!

Why?

Concepts are different and concomitant resultant orientationality of the event is much moiré phantasmagorical!

True words!

Deleuze, also, acknowledges this traditional conception of concepts as universal descriptions. In prefacing the English translation of *Dialogues*, 7 he says: In so-called rationalist philosophies, the abstract is given the task of explaining, and it is the abstract that is realized in the concrete. One starts with abstractions such as the One, the Whole, the Subject... Even if this means undergoing a terrible crisis each time that **one sees rational unity or totality turning into their opposites**, or the subject generating monstrosities (Dvii).(as quoted in Peter Cook's essay *Thinking the Concept Otherwise: Deleuze and Expression* PETER COOK)

Yes! Rational unity or totalities turn in to opposites!

What happens?

Abstract is not abstract!

It becomes distract!

Event ramifies in to some other which you understand as perception!

What in the ultimate is your idea of life?

Self preservation, self adaptation, self transcendence and self dissolution!

You have defined self as witness consciousness and dynamic in case of evolutionary case!

Yes!

Introduction:

Note: Models for the following abstracts are given in the next paper in the series, such procrastination as such, attributable and ascribable to the restrictions' on space.

Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect Phys. Rev. B 61, 10267 – Published 15 April 2000 N. Read and Dmitry Green

- (1) Authors analyze pairing of fermions in two dimensions for fully gapped cases with broken parity (P) and time reversal (T), especially cases in which the gap function is an orbital angular momentum (l) eigenstate, in particular $l=1$ (p wave, spinless, or spin triplet) and $l=2$ (d wave, spin singlet). For $l \neq 0$, this fall into two phases, weak and strong pairing, which may be distinguished topologically?
- (2) In the cases with conserved spin, we derive explicitly the Hall conductivity for spin as the corresponding topological invariant.
- (3) For the spinless p-wave case, the weak-pairing phase has a pair wave function that is asymptotically the same as that in the Moore-Read (Pfaffian) Quantum Hall State, and we argue that its other properties (edge states, quasiholes, and toroidal ground states) are also the same, indicating that nonabelian statistics is a generic property of such a paired phase.

- (4) The strong-pairing phase is an abelian state, and the transition between the two phases involves a bulk Majorana fermion, the mass of which changes sign at the transition.
- (5) For the d-wave case, we argue that the Haldane-Rezayi state is not the generic behavior of a phase but describes the asymptotics at the critical point between weak and strong pairing, and has gapless fermion excitations in the bulk.
- (6) In this case the weak-pairing phase is an abelian phase, which has been considered previously.
- (7) In the p-wave case with an unbroken U(1) symmetry, which can be applied to the double layer quantum Hall problem, the weak-pairing phase has the properties of the 331 state, and with nonzero tunneling there is a transition to the Moore-Read phase.
- (8) The effects of disorder on noninteracting quasiparticles are considered.
- (9) The gapped phases survive, but there is an intermediate thermally conducting phase in the spinless p-wave case, in which the quasiparticles are extended. DOI: <http://dx.doi.org/10.1103/PhysRevB.61.10267>
 Received 30 June 1999 Published in the issue dated 15 April 2000 © 2000 The American Physical Society

International Journal of Modern Physics B Condensed Matter Physics; Statistical Physics; Atomic, Molecular and Optical Physics Volume 06, Issue 01, January 10, 1992 SHOU CHENG ZHANG, Int. J. Mod. Phys B, 06, 25 (1992) DOI: 10.1142/S0217979292000037

- (10) This paper gives a systematic review of a field theoretical approach to the fractional quantum Hall effect (FQHE) that has been developed in the past few years.
- (11) Authors first illustrate some simple physical ideas to motivate such an approach and then present a systematic derivation of the Chern–Simons–Landau–Ginzburg (CSLG) action for the FQHE, starting from the microscopic Hamiltonian
- (12) It is demonstrated that all the phenomenological aspects of the FQHE can be derived from the mean field solution and the small fluctuations of the CSLG action.
- (13) Although this formalism is logically independent of Laughlin's wave function approach, their physical consequences are equivalent
- (14) The CSLG theory demonstrates a deep connection between the phenomena of superfluidity and the FQHE, and can provide a simple and direct formalism to address many new macroscopic phenomena of the FQHE.

Letter abstract Nature Physics 5, 258 - 261 (2009) Published online: 15 March 2009 | doi:10.1038/nphys1227 Signature of magnetic monopole and Dirac string dynamics in spin ice L. D. C. Jaubert¹ & P. C. W. Holdsworth¹

- (15) Magnetic monopoles have eluded experimental detection since their prediction nearly a century ago by Dirac..
- (16) It was recently shown that classical analogues of these enigmatic particles can occur as excitations out of the topological ground state of a model magnetic system, dipolar spin ice.
- (17) These quasi-particle excitations do not require a modification of Maxwell's equations, but they do interact through Coulomb's law and are of magnetic origin.
- (18) Here, we present an experimentally measurable signature of monopole dynamics.
- (19) In particular, we show that previous magnetic relaxation measurements in the spin-ice material Dy₂Ti₂O₇ (ref. 3) can be interpreted entirely in terms of the diffusive motion of monopoles in the grand canonical ensemble, constrained by a network of 'Dirac strings' filling the quasi-particle vacuum.
- (20) In a magnetic field, the topology of the network prevents charge flow in the steady state. Nevertheless, we demonstrate the existence of a monopole density gradient near the surface of an open system.

Boris Pioline 2006 Class. Quantum Grav 23 S981 doi:10.1088/0264-9381/23/21/S05 Lectures on black holes, topological strings and quantum attractors Boris Pioline

- (21) In these lecture notes, authors review some recent developments on the relation between the macroscopic entropy of four-dimensional BPS black holes and the microscopic counting of states, beyond the thermodynamical, large charge limit.
- (22) After a brief overview of charged black holes in supergravity and string theory, authors give an extensive introduction to special and very special geometry, attractor flows and topological string theory, including holomorphic anomalies.
- (23) Authors then expose the Ooguri–Strominger–Vafa (OSV) conjecture which relates microscopic degeneracies to the topological string amplitude and review precision tests of this formula on 'small' black holes.
- (24) Finally, motivated by a holographic interpretation of the OSV conjecture, we discuss the radial quantization of BPS black holes (i.e. quantum attractors) and present a recent conjecture relating exact black hole degeneracies to Fourier coefficients of certain automorphic forms.

Nature 464, 199-208 (11 March 2010) | doi: 10.1038/nature08917; Published online 10 March 2010 Spin liquids in frustrated magnets Leon Balents

- (25) Frustrated magnets are materials in which localized magnetic moments, or spins, interact through competing exchange interactions that cannot be simultaneously satisfied, giving rise to a large degeneracy of the system ground state.
- (26) Under certain conditions, this can lead to the formation of fluid-like states of matter, so-called spin liquids, in which the constituent spins are highly correlated but still fluctuate strongly down to a temperature of absolute zero.
- (27) The fluctuations of the spins in a spin liquid can be classical or quantum and show remarkable collective phenomena such as emergent gauge fields and fractional particle excitations.
- (28) This exotic behaviour is now being uncovered in the laboratory, providing insight into the properties of spin liquids and challenges to the theoretical description of these materials.

Superstrings and Topological Strings at Large N Cumrun Vafa

- (29) Authors embed the large N Chern-Simons/topological string duality in ordinary superstrings.
- (30) This corresponds to a large N duality between generalized gauge systems with $N=1$ supersymmetry in 4 dimensions and superstrings propagating on non-compact Calabi-Yau manifolds with certain fluxes turned on.
- (31) They also show that in a particular limit of the $N=1$ gauge theory system, certain superpotential terms in the $N=1$ system (including deformations if spacetime is non-commutative) are captured to all orders in $1/N$ by the amplitudes of non-critical bosonic strings propagating on a circle with self-dual radius.
- (32) They also consider D-brane/anti-D-brane system wrapped over vanishing cycles of compact Calabi-Yau manifolds and argue that at large N they induce a shift in the background to a topologically distinct Calabi-Yau, which we identify as the ground state system of the Brane/anti-Brane system. Journal reference: J.Math.Phys. 42 (2001) 2798-2817 Report number: HUTP-00/A035 arXiv: hep-th/0008142

Skyrmions and the crossover from the integer to fractional quantum Hall effect at small Zeeman energies Phys. Rev. B 47, 16419 – Published 15 June 1993 S. L. Sondhi, A. Karlhede, S. A. Kivelson, and E. H. Rezayi

- (33) **S. L. Sondhi, A. Karlhede, S. A. Kivelson, and E. H. Rezayi** study the two-dimensional electron gas in a high magnetic field at filling factor $\nu=1$ for an arbitrary ratio of the Zeeman energy $g\mu_B B$ to the typical interaction energy.
- (34) **They** find that the **system always has a gap**, even when the one-particle gap vanishes, i.e., when $g=0$. When g is sufficiently large, the quasiparticles are perturbatively related to those in the noninteracting limit; we compute their energies to second order in the Coulomb interaction.

- (35) For g smaller than a critical value g_c the quasiparticles change character; in the limit of $g \rightarrow 0$, they are skyrmions—spatially unbounded objects with infinite spin.
- (36) In GaAs heterojunctions, the gap is unambiguously predominantly due to correlation effects; indeed, we tentatively conclude that g is always smaller than g_c , so the relevant quasiparticles are the skyrmions.
- (37) The generalization to other odd-integer filling factors, and to $\nu=1/3$ and $1/5$, is discussed. DOI: <http://dx.doi.org/10.1103/PhysRevB.47.16419> © 1993 The American Physical Society

News and Views Nature 421, 796-797 (20 February 2003) | doi: 10.1038/421796a Gianni Blatter Scientist Position in Human Molecular Genetics Quantum computing: The qubit duet

- (38) Small, but consistent, steps are being taken towards the realization of a quantum computer.
- (39) The demonstration of the coupling of two quantum bits in a solid-state device moves us closer to that goal.
- (40) Basic steps towards the creation of a quantum computer have been taken, with the demonstrations of elementary data storage and manipulation using photons and atoms¹ or trapped ions^{2, 3} as the quantum bits, or 'qubits'.
- (41) Solid-state qubits (made, for example, from tiny samples of superconducting material) are an attractive alternative, however, as they could be more easily built into working devices, profiting from the highly developed methods of nanotechnology.

Published Online August 7 2003 Science 5 September 2003: Vol. 301 no. 5638 pp. 1348-135 DOI: 10.1126/science.1087128 Dissipationless Quantum Spin Current at Room Temperature Shuichi Murakami^{1,*}, Naoto Nagaosa^{1, 2, 3}, Shou-Cheng Zhang⁴

- (42) Although microscopic laws of physics are invariant under the reversal of the arrow of time, the transport of energy and information in most devices is an irreversible process.
- (43) It is this irreversibility that leads to intrinsic dissipations in electronic devices and limits the possibility of quantum computation.
- (44) Authors theoretically predict that the electric field can induce a substantial amount of dissipationless quantum spin current at room temperature, in hole-doped semiconductors such as Si, Ge, and GaAs. On the basis of a generalization of the quantum Hall effect, the predicted effect leads to efficient spin injection without the need for metallic ferromagnets.
- (45) Principles found here could enable quantum spintronic devices with integrated information processing and storage units, operating with low power consumption and performing reversible quantum computation.

T Thiemann 1998 Class Quantum Grav 15 1207 doi:10.1088/0264-9381/15/5/010 Quantum spin dynamics (QSD): III. Quantum constraint algebra and physical scalar product in quantum general relativity

- (46) This paper deals with several technical issues of non-perturbative four-dimensional Lorentzian canonical quantum gravity in the continuum that arose in connection with the recently constructed Wheeler-DeWitt quantum constraint operator.
- (47) The Wheeler-DeWitt constraint of quantum general relativity mixes the diffeomorphism superselection sectors for diffeomorphism-invariant theories of connections that were previously discussed in the literature.
- (48) From it one can construct diffeomorphism-invariant operators which do not necessarily commute with the Hamiltonian constraint but which still mix those sectors and which, at the diffeomorphism-invariant level, encode physical information.
- (49) Thus, if one adopts, as before in the literature, the strategy to solve the diffeomorphism constraint before the Hamiltonian constraints then those sectors become spurious.
- (50) The inner product for diffeomorphism-invariant states can be fixed by requiring that diffeomorphism group averaging is a partial isometry.

- (51) The established non-anomalous constraint algebra is clarified by computing commutators of duals of constraint operators.
- (52) The full classical constraint algebra is faithfully implemented on the diffeomorphism-invariant Hilbert space in an appropriate sense.
- (53) The Hilbert space of diffeomorphism-invariant states can be made separable if a natural new superselection principle is satisfied. We propose a natural physical scalar product for quantum general relativity by extending the group-average approach to the case of non-self-adjoint constraint operators like the Wheeler-DeWitt constraint. Equipped with this inner product, the construction of physical observables is straightforward.

Published Online September 1 2011 Science 7 October 2011: Vol. 334 no. 6052 pp. 57-61 DOI: 10.1126/science.1208001 Universal Digital Quantum Simulation with Trapped Ions B. P. Lanyon et al

- (54) A digital quantum simulator is an envisioned quantum device that can be programmed to efficiently simulate any other local system.
- (55) Authors demonstrate and investigate the digital approach to quantum simulation in a system of trapped ions.
- (56) With sequences of up to 100 gates and 6 qubits, the full time dynamics of a range of spin systems are digitally simulated.
- (57) Interactions beyond those naturally present in our simulator are accurately reproduced, and quantitative bounds are provided for the overall simulation quality.
- (58) Results demonstrate the key principles of digital quantum simulation and provide evidence that the level of control required for a full-scale device is within reach.

6, 382 - 388 (2010) Published online: 14 March 2010 | doi: 10.1038/nphys1614 Subject Categories: Atomic and molecular physics | Quantum physics | Condensed-matter physics A Rydberg quantum simulator Hendrik Weimer¹, Markus Müller², Igor Lesanovsky^{2,3}, Peter Zoller² & Hans Peter Büchler¹ Nature Physics

- (59) A universal quantum simulator is a controlled quantum device that reproduces the dynamics of any other many-particle quantum system with short-range interactions.
- (60) This dynamics can refer to both coherent Hamiltonian and dissipative open-system evolution.
- (61) Here we propose that laser-excited Rydberg atoms in large-spacing optical or magnetic lattices provide an efficient implementation of a universal quantum simulator for spin models involving n-body interactions, including such of higher order
- (62) This would allow the simulation of Hamiltonians of exotic spin models involving n-particle constraints, such as the Kitaev toric code, colour code and lattice gauge theories with spin-liquid phases.
- (63) In addition, our approach provides the ingredients for dissipative preparation of entangled states based on engineering n-particle reservoir couplings.
- (64) The basic building blocks of our architecture are efficient and high-fidelity n-qubit entangling gates using auxiliary Rydberg atoms, including a possible dissipative time step through optical pumping.
- (65) This enables mimicking the time evolution of the system by a sequence of fast, parallel and high-fidelity n-particle coherent and dissipative Rydberg gates.

Journal of Statistical Physics July 2003, Volume 112, Issue 1-2, pp 219-275 Pattern Selection: Determined by Symmetry and Modifiable by Distant Effects Mitchell J. Feigenbaum

- (66) Authors consider Saffman–Taylor channel flow without surface tension on a high-pressure driven interface, but modify the usual infinite-fluid in infinite-channel configuration.
- (67) Here we include the treatment of efflux by considering a finite connected body of fluid in an arbitrarily long channel, with its second free interface the efflux of this configuration.

- (68) They show that there is a uniquely determined translating solution for the driven interface, which is exactly the $1/2$ width S–T solution, following from correct symmetry for a finite channel flow.
- (69) Authors establish that there exist no perturbations about this solution corresponding to a finger propagating with any other width: Selection is locally unique and isolated.
- (70) The stability of this solution is anomalous, in that all freely impenetrable perturbations are stabilities, while unstable modes request power proportional to their strength from the external agencies that drive the flow, and so, in principle, are experimentally controllable.
- (71) This is very different from the behavior of the usual infinite fluid. We conjecture that surface tension on the efflux interface modifies channel-width λ according to $1-2\lambda=\sigma/v$ (i.e., $(2\pi)^2 B$ of the literature) with v the velocity of the high-pressure tip, but σ the surface tension of the efflux.
- (72) That is, λ is decreased below $1/2$ by the effect of smoothing the distant efflux.
- (73) The perturbation theory created here to deal with transport between two free boundaries is novel and dependent upon a symmetry implied by the equations of motion.

Journal of Statistical Physics June 2001, Volume 103, Issue 5-6, pp 973-1007 Dynamics of Finger Formation in Laplacian Growth without Surface Tension Mitchell J. Feigenbaum, Itamar Procaccia, Benny Davidovich

- (74) Authors study the dynamics of “finger” formation in Laplacian growth without surface tension in channel geometry (the **Saffman–Taylor problem**).
- (75) They present a pedagogical derivation of the dynamics of the conformal map from a strip in the complex plane to the physical channel. In doing so we pay attention to the boundary conditions (no flux rather than periodic) and derive a field equation of motion for the conformal map.
- (76) Authors first consider an explicit analytic class of conformal maps that form a basis for solutions in infinitely long channels, characterized by meromorphic derivatives.
- (77) The great bulk of these solutions can lose conformality due to finite time singularities.
- (78) By considerations of the nature of the analyticity of these solutions, they show that those solutions which are free of such singularities inevitably result in a single asymptotic “finger” whose width is determined by initial conditions.
- (79) This is in contradiction with the experimental results that indicate selection of a finger of width $1/2$.
- (80) In the last part of this paper we show that such a solution might be determined by the boundary conditions of a finite body of fluid, e.g. finiteness can lead to pattern selection.

Conformal dynamics of fractal growth patterns without randomness Phys. Rev. E 62, 1706 – Published 1 August 2000 Benny Davidovitch, M. J. Feigenbaum, H. G. E. Hentschel, and Itamar Procaccia

- (81) Many models of fractal growth patterns (such as diffusion limited aggregation and dielectric breakdown models) combine complex geometry with randomness; this double difficulty is a stumbling block to their elucidation.
- (82) In this paper we introduce a wide class of fractal growth models with highly complex geometry but without any randomness in their growth rules.
- (83) The models are defined in terms of deterministic itineraries of iterated conformal maps, generating the function $\Phi(n)(\omega)$ which maps the exterior of the unit circle to the exterior of an n -particle growing aggregate.
- (84) The complexity of the evolving interfaces is fully contained in the deterministic dynamics of the conformal map $\Phi(n)(\omega)$.
- (85) They focus attention on a class of growth models in which the itinerary is quasiperiodic. Such itineraries can be approached via a series of rational approximants.
- (86) The analytic power gained is used to introduce a scaling theory of the fractal growth patterns and to identify the exponent that determines the fractal dimension. DOI:

<http://dx.doi.org/10.1103/PhysRevE.62.1706> Published in the issue dated August 2000 © 2000 The American Physical Society

Sandpiles, avalanches, and the statistical mechanics of nonequilibrium stationary states Phys. Rev. E 47, 3099 – Published 1 May 1993 Ashvin B. Chhabra, Mitchell J. Feigenbaum, Leo P. Kadanoff, Amy J. Kolan, and Itamar Procaccia

- (87) The scaling properties of three nontrivial one-dimensional avalanche models are analyzed.
- (88) The first two of them are the local limited model with one open, one closed, and with periodic boundary conditions, respectively.
- (89) A theory for the scaling properties of these models based on the existence of two fundamental length scales, which diverge in the thermodynamic limit, is developed.
- (90) The third model studied is a trapless version of the nonperiodic local limited model.
- (91) They find that it is scale invariant. Our theoretical predictions are compared with extensive computer simulations in all three cases. DOI: <http://dx.doi.org/10.1103/PhysRevE.47.3099>

S Barbarino et al 2014 Smart Mater Struct 23 063001 doi:10.1088/0964-1726/23/6/063001 A review on shape memory alloys with applications to morphing aircraft FEATURED ARTICLE REVIEW ARTICLE

- (92) Shape memory alloys (SMAs) are a unique class of metallic materials with the ability to recover their original shape at certain characteristic temperatures (shape memory effect), even under high applied loads and large inelastic deformations, or to undergo large strains without plastic deformation or failure (super-elasticity).
- (93) In this review, authors describe the main features of SMAs, their constitutive models and their properties.
- (94) They also review the fatigue behavior of SMAs and some methods adopted to remove or reduce its undesirable effects.
- (95) SMAs have been used in a wide variety of applications in different fields.
- (96) In this review, we focus on the use of shape memory alloys in the context of morphing aircraft, with particular emphasis on variable twist and camber, and also on actuation bandwidth and reduction of power consumption.
- (97) These applications prove particularly challenging because novel configurations are adopted to maximize integration and effectiveness of SMAs, which play the role of an actuator (using the shape memory effect), often combined with structural, load-carrying capabilities.
- (98) Iterative and multi-disciplinary modeling is therefore necessary due to the fluid–structure interaction combined with the nonlinear behavior of SMAs.

Jong-Ha Chung et al 2007 Smart Mater Struct 16 N1 doi:10.1088/0964-1726/16/1/N01 Implementation strategy for the dual transformation region in the Brinson SMA constitutive model

- (99) The transformation kinetics formulation is the principal factor underlying the constitutive model of shape memory alloys (SMAs).
- (100) Therefore, the transformation kinetics formulation, which is applicable to any status of stress and temperature, is essential for predicting the material behavior of SMAs.
- (101) In this work we show that the transformation kinetics of the original Brinson model, which is the most widely used one-dimensional model, has shortcomings in the case where temperature decreases at low temperature ($T < M_s$).
- (102) In addition, we propose a modified transformation kinetics formula that can be used for dual transformation conditions.

- (103) The martensite transformation kinetics is modified so that the transformation from austenite into temperature-induced martensite, due to the decrease in temperature, is coupled with a transformation from austenite or temperature-induced martensite into stress-induced martensite, due to the increase in stress.
- (104) Through this modification, the suggested formulation can properly describe the behavior of martensite fractions in the dual transformation region.

Lisa Dyson et al JHEP08 (2002)045 doi:10.1088/1126-6708/2002/08/045 Is There Really a de Sitter/CFT Duality Lisa Dyson¹, James Lindesay¹ and Leonard Susskind¹

- (105) In this paper a de Sitter Space version of Black Hole Complementarity is formulated which states that an observer in de Sitter Space describes the surrounding space as a sealed finite temperature cavity bounded by a horizon which allows no loss of information.
- (106) Authors then discuss the implications of this for the existence of boundary correlators in the hypothesized dS/Cft correspondence.
- (107) They find that dS complementarity precludes the existence of the appropriate limits.
- (108) Authors find that the limits exist only in approximations in which the entropy of the de Sitter Space is infinite.
- (109) The reason that the correlators exist in quantum field theory in the de Sitter Space background is traced to the fact that horizon entropy is infinite in QFT.

Toda Theories, Matrix Models, Topological Strings, and N=2 Gauge Systems Robbert Dijkgraaf, Cumrun Vafa

- (110) Authors consider the topological string partition function, including the Nekrasov deformation, for type IIB geometries with an A_{n-1} singularity over a Riemann surface.
- (111) These models realize the $N=2$ $SU(n)$ superconformal gauge systems recently studied by Gaiotto and collaborators.
- (112) Employing large N dualities we show why the partition function of topological strings in these backgrounds is captured by the chiral blocks of A_{n-1} Toda systems and derive the dictionary recently proposed by Alday, Gaiotto and Tachikawa.
- (113) For the case of genus zero Riemann surfaces, we show how these systems can also be realized by Penner-like matrix models with logarithmic potentials.
- (114) The Seiberg-Witten curve can be understood as the spectral curve of these matrix models which arises holographically at large N . In this context the Nekrasov deformation maps to the beta-ensemble of generalized matrix models, that in turn maps to the Toda system with general background charge.
- (115) They also point out the notion of a double holography for this system, when both n and N are large.

Physics Reports Volume 94, Issue 6, March 1983, Pages 313–404 Quantum integrable systems related to lie algebras M.A. Olshanetsky, A.M. Perelomov doi:10.1016/0370-1573(83)90018-2

- (116) Some quantum integrable finite-dimensional systems related to Lie algebras are considered.
- (117) This review continues the previous review of the same authors [83] devoted to the classical aspects of these systems.
- (118) The dynamics of some of these systems is closely related to free motion in symmetric spaces.
- (119) Using this connection with the theory of symmetric spaces some results such as the forms of spectra, wave functions-matrices, quantum integrals of motion are derived.
- (120) In specific cases the considered systems describe the one-dimensional n -body systems interacting pairwise via potentials $g_2 v(q)$ of the following 5 types: $v_I(q) = q^{-2}$, $v_{II}(q) = \sinh^{-2} q$, $v_{III}(q) = \sin^{-2} q$, $v_{IV}(q) = \mathcal{P}(q)$, $v_V(q) = q^{-2} + \omega^2 q^2$.

- (121) Here $\wp(q)$ is the Weierstrass function, so that the first three cases are merely subcases of the fourth. The system characterized by the Toda nearest-neighbour potential $\exp(q|q| + 1)$ is moreover considered.
- (122) This review presents from a general and universal point of view results obtained mainly over the past fifteen years.
- (123) Besides, it contains some new results both of physical and mathematical interest. Copyright © 1983 Published by Elsevier B.V.

Shamik Banerjee et al 2013 Class. Quantum Grav. 30 104001 doi:10.1088/0264-9381/30/10/104001 Smoothed transitions in higher spin AdS gravity

- (124) Authors consider CFTs conjectured to be dual to higher spin theories of gravity in AdS3 and AdS4.
- (125) Two-dimensional CFTs with \mathcal{W}_N symmetry are considered in the $\lambda = 0$ ($k \rightarrow \infty$) limit where they are conjectured to be described by continuous orbifolds.
- (126) The torus partition function is computed, using reasonable assumptions, and equals that of a free-field theory.
- (127) They find no phase transition at temperatures of order 1; the usual Hawking–Page phase transition is removed by the highly degenerate light states associated with conical defect states in the bulk.
- (128) Three-dimensional Chern–Simons matter CFTs with vector-like matter are considered on T^3 , where the dynamics is described by an effective theory for the eigenvalues of the holonomies.
- (129) Likewise, we find no evidence for a Hawking–Page phase transition at a large level k .

M Akyol and G Papadopoulos 2014 Class. Quantum Grav 31 123001 doi:10.1088/0264-9381/31/12/123001 Spinorial geometry, horizons and superconformal symmetry in six dimensions

- (130) The spinorial geometry method of solving Killing spinor equations is reviewed as it applies to six-dimensional $(1, 0)$ supergravity.
- (131) In particular, it is explained how the method is used to identify both the fractions of supersymmetry preserved by and the geometry of all supersymmetric backgrounds.
- (132) Then two applications are described to systems that exhibit superconformal symmetry. The first is the proof that some six-dimensional black hole horizons are locally isometric to $\text{AdS}_3 \times \Sigma_3$, where Σ_3 is diffeomorphic to S^3 .
- (133) The second one is a description of all supersymmetric solutions of six-dimensional $(1, 0)$ superconformal theories and in particular of their brane solitons.

Naoyuki Kawahara et al JHEP10 (2007)097 doi:10.1088/1126-6708/2007/10/097 Phase structure of matrix quantum mechanics at finite temperature

- (134) Authors study matrix quantum mechanics at finite temperature by Monte Carlo simulation.
- (135) The model is obtained by dimensionally reducing 10d $U(N)$ pure Yang-Mills theory to 1d. Following Aharony et al., one can view the same model as describing the high temperature regime of $(1+1)$ d $U(N)$ super Yang-Mills theory on a circle.
- (136) In this interpretation an analog of the deconfinement transition was conjectured to be a continuation of the black-hole/black-string transition in the dual gravity theory.
- (137) Our detailed analysis in the critical regime up to $N = 32$ suggests the existence of the non-uniform phase, in which the eigenvalue distribution of the holonomy matrix is non-uniform but gapless.
- (138) The transition to the gapped phase is of second order.

- (139) The internal energy is constant (giving the ground state energy) in the uniform phase, and rises quadratically in the non-uniform phase, which implies that the transition between these two phases is of third order.

Advances in Space Research Available online 28 November 2014 Internal wave coupling processes in Earth's atmosphere Erdal Yiğita, Alexander S. Medvedev, c, doi:10.1016/j.asr.2014.11.020

- (140) This paper presents a contemporary review of vertical coupling in the atmosphere and ionosphere system induced by internal waves of lower atmospheric origin.
- (141) Atmospheric waves are primarily generated by meteorological processes, possess a broad range of spatial and temporal scales, and can propagate to the upper atmosphere.
- (142) A brief summary of internal wave theory is given, focusing on gravity waves, solar tides, and planetary Rossby and Kelvin waves.
- (143) Observations of wave signatures in the upper atmosphere, their relationship with the direct propagation of waves into the upper atmosphere, dynamical and thermal impacts as well as concepts, approaches, and numerical modeling techniques are outlined.
- (144) Recent progress in studies of sudden stratospheric warming and upper atmospheric variability are discussed in the context of wave-induced vertical coupling between the lower and upper atmosphere.

Journal of High Energy Physics January 2010, 2010:114 Date: 27 Jan 2010 Holographic quantum liquids in 1+1 dimensions Ling-Yan Hung, Aninda Sinha

- (145) In this paper we initiate the study of holographic quantum liquids in 1+1 dimensions. Since the Landau Fermi liquid theory breaks down in 1+1 dimensions, it is of interest to see what holographic methods have to say about similar models.
- (146) For theories with a gapless branch, the Luttinger conjecture states that there is an effective description of the physics in terms of a Luttinger liquid which is specified by two parameters.
- (147) The theory we consider is the defect CFT arising due to a probe D3 brane in the AdS Schwarzschild planar black hole background.
- (148) They turn on a fundamental string density on the worldvolume.
- (149) Unlike higher dimensional defects, a persistent dissipationless zero sound mode is found.
- (150) The thermodynamic aspects of these models are considered carefully and certain subtleties with boundary terms are explained which are unique to 1+1 dimensions.
- (151) Spectral functions of bosonic and fermionic fluctuations are also considered and quasinormal modes are analysed.
- (152) A prescription is given to compute spectral functions when there are mixing due to the worldvolume gauge field.
- (153) Comment is on the Luttinger conjecture in the light of our findings. ArXiv ePrint: 0909.3526

M J Feigenbaum 1988 Nonlinearity 1 577 doi:10.1088/0951-7715/1/4/005 Presentation functions and scaling function theory for circle maps

- (154) Considering first return maps, a most natural renormalisation group fixed point is determined.
- (155) From it a simple presentation function is constructed, immediately leading to the thermodynamics of critical rotation.
- (156) The rotation number is encoded in the topological action of the presentation function and the algebraic singularity of critically in that function's derivatives at its fixed points.
- (157) Any such presentation function determines a circle map dynamics of that rotation number and index of criticality.
- (158) These functions are naturally parametrised by a trajectory scaling function.

- (159) The requirement that the dynamics be smooth leads to a prescription for the calculation of the scaling function and hence the dynamics.
- (160) The theory is highly constrained and suffers in finite-order approximation from the extra constraint of commutativity, which however can be overcome.

Topological Strings from Quantum Mechanics Alba Grassi, Yasuyuki Hatsuda, Marcos Marino

- (161) Authors propose a general correspondence which associates a non-perturbative quantum-mechanical operator to a toric Calabi-Yau manifold, and conjecture an explicit formula for its spectral determinant in terms of an M-theoretic version of the topological string free energy.
- (162) As a consequence, authors derive an exact quantization condition for the operator spectrum, in terms of the vanishing of a generalized theta function.
- (163) The perturbative part of this quantization condition is given by the Nekrasov-Shatashvili limit of the refined topological string, but there are non-perturbative corrections determined by the conventional topological string.
- (164) They analyze in detail the cases of local P2, local P1xP1 and local F1. In all these cases, the predictions for the spectrum agree with the existing numerical results.
- (165) Authors also show explicitly that our conjectured spectral determinant leads to the correct spectral traces of the corresponding operators, which are closely related to topological string theory at orbifold points.
- (166) Physically, our results provide a Fermi gas picture of topological strings on toric Calabi-Yau manifolds, which is fully non-perturbative and background independent.
- (167) They also suggest the existence of an underlying theory of M2 branes behind this formulation.
- (168) Mathematically results lead to precise, surprising conjectures relating the spectral theory of functional difference operators to enumerative geometry. ArXiv: 1410.3382 [hep-th]

Baishali Chakraborty et al 2013 J. Phys.: Conf. Ser. 442 012017 doi:10.1088/1742-6596/442/1/012017 Topology, cosmic strings and quantum dynamics – a case study with graphene

- (169) Authors explore the possibility to study the quantum dynamics of Dirac fermions in presence of a cosmic string by introducing a conical topological defect in gapped graphene in the presence of a Coulomb charge.
- (170) When the Coulomb charge exceeds a certain critical strength, quantum instability sets in.
- (171) Below the critical regime and for certain values of the system parameters, the allowed boundary conditions in gapped graphene cone can be classified in terms of a single real quantity.
- (172) Observables such as local density of states, scattering phase shifts and the bound state spectra are dependent on the value of this real parameter, which has to be determined empirically.
- (173) For a supercritical Coulomb charge, we analyze the system with a regularized potential as well as with a zigzag boundary condition and find the effect of the sample topology on the observable features of the system.

Nuclear Physics B Volume 306, Issue 4, 5 September 1988, Pages 890–907 Axion-induced topology change in quantum gravity and string theory Steven B. Giddings doi:10.1016/0550-3213(88)90446-4

- (174) Authors consider a system comprised of an axion (described by rank-three antisymmetric tensor field strength) coupled to gravity. Instantons are found which describe the nucleation of a Planck-sized **baby Robertson-Walker universe**.
- (175) Information loss to the baby universe can lead to an effective loss of quantum coherence.
- (176) An estimate of the magnitude of this effect on particle propagation is made in the semi-classical approximation.

- (177) This magnitude depends on the parameters of the theory (which includes a cutoff since the theory is non-renormalizable) and on the quantum state of the many-universe system.
- (178) In contrast to the naive expectation that Planck-scale dynamics should lead to very small effects at low energies, the effects of these instantons can be large.
- (179) The case of string theory is considered in some detail, and it is found that a massless dilaton can suppress the tunneling.

Nuclear Physics B Volume 536, Issues 1–2, 21 December 1998, Pages 407–434 Non-perturbative Lorentzian quantum gravity, causality and topology change J. Ambjørn, R. Loll doi:10.1016/S0550-3213(98)00692-0

- (180) Authors formulate a non-perturbative lattice model of two-dimensional Lorentzian quantum gravity by performing the path integral over geometries with a causal structure.
- (181) The model can be solved exactly at the discretized level.
- (182) Its continuum limit coincides with the theory obtained by quantizing 2d continuum gravity in proper-time gauge, but it disagrees with 2d gravity defined via matrix models or Liouville theory.
- (183) By allowing topology change of the compact spatial slices (i.e. baby universe creation), one obtains agreement with the matrix models and Liouville theory.

Communications in Mathematical Physics January 2006, Volume 261, Issue 2, pp 451-516 Topological Strings and Integrable Hierarchies Mina Aganagic et al

- (184) Authors consider the topological B-model on local Calabi-Yau geometries.
- (185) They show how one can solve for the amplitudes by using $-$ -algebra symmetries which encode the symmetries of holomorphic diffeomorphisms of the Calabi-Yau.
- (186) In the highly effective fermionic/brane formulation this leads to a free fermion description of the amplitudes.
- (187) Furthermore we argue that topological strings on Calabi-Yau geometries provide a unifying picture connecting non-critical (super) strings, integrable hierarchies, and various matrix models.
- (188) In particular we show how the ordinary matrix model, the double scaling limit of matrix models, and Kontsevich-like matrix model are all related and arise from studying branes in specific local Calabi-Yau three-folds.
- (189) They also show how an A-model topological string on P^1 and local toric threefolds (and in particular the topological vertex) can be realized and solved as B-model topological string amplitudes on a Calabi-Yau manifold. Communicated by N.A. Nekrasov

Toda Theories, Matrix Models, Topological Strings, and $N=2$ Gauge Systems Robbert Dijkgraaf, Cumrun Vafa

- (190) Authors consider the topological string partition function, including the Nekrasov deformation, for type IIB geometries with an A_{n-1} singularity over a Riemann surface.
- (191) These models realize the $N=2$ $SU(n)$ superconformal gauge systems recently studied by Gaiotto and collaborators. Employing large N dualities we show why the partition function of topological strings in these backgrounds is captured by the chiral blocks of A_{n-1}
- (192) Toda systems and derive the dictionary recently proposed by Alday, Gaiotto and Tachikawa.
- (193) For the case of genus zero Riemann surfaces, we show how these systems can also be realized by Penner-like matrix models with logarithmic potentials.
- (194) The Seiberg-Witten curve can be understood as the spectral curve of these matrix models which arises holographically at large N .
- (195) In this context the Nekrasov deformation maps to the beta-ensemble of generalized matrix models, that in turn maps to the Toda system with general background charge.

- (196) They also point out the notion of a double holography for this system, when both n and N are large. ArXiv: 0909.2453 [hep-th]

Journal of High Energy Physics Quantum geometry of refined topological strings Mina Aganagic et al

- (197) Authors consider branes in refined topological strings.
 (198) They argue that their wavefunctions satisfy a Schrödinger equation depending on multiple times and prove this in the case where the topological string has a dual matrix model description.
 (199) Furthermore, in the limit where one of the equivariant rotations approaches zero, the brane partition function satisfies a time-independent Schrödinger equation.
 (200) Authors use this observation, as well as the back reaction of the brane on the closed string geometry, to offer an explanation of the connection between integrable systems and $N=2$ gauge systems in four dimensions observed by Nekrasov and Shatashvili.

Journal of Statistical Physics August 1988, Volume 52, Issue 3-4, pp 527-569 Presentation functions, fixed points, and a theory of scaling function dynamics Mitchell J. Feigenbaum

- (201) Presentation functions provide the time-ordered points of the forward dynamics of a system as successive inverse images.
 (202) They generally determine objects constructed on trees, regular or otherwise and immediately determine a functional form of the transfer matrix of these systems.
 (203) Presentation functions for regular binary trees determine the associated forward dynamics to be that of a period doubling fixed point.
 (204) They are generally parametrized by the trajectory scaling function of the dynamics in a natural way.
 (205) The requirement that the forward dynamics be smooth with a critical point determines a complete set of equations whose solution is the scaling function.
 (206) These equations are compatible with a dynamics in the space of scalings which is conjectured, with numerical and intuitive support, to possess its solution as a unique, globally attracting fixed point.
 (207) It is argued that such dynamics is to be sought as a program for the solution of chaotic dynamics. In the course of the exposition new information pertaining to universal mode locking is presented.

Quantum simulation of frustrated Ising spins with trapped ions K. Kim, M.-S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G.-D. Lin, L.-M. Duan & C. Monroe Nature 465, 590–593 (03 June 2010) doi: 10.1038/nature09071

- (208) A network is frustrated when competing interactions between nodes prevent each bond from being satisfied.
 (209) This compromise is central to the behaviour of many complex systems, from social¹ and neural² networks to protein folding³ and magnetism^{4, 5}.
 (210) Frustrated networks have highly degenerate ground states, with excess entropy and disorder even at zero temperature.
 (211) In the case of quantum networks, frustration can lead to massively entangled ground states, underpinning exotic materials such as quantum spin liquids and spin glasses^{6, 7, 8, 9}.
 (212) Here we realize a quantum simulation of frustrated Ising spins in a system of three trapped atomic ions^{10, 11, and 12}, whose interactions are precisely controlled using optical forces¹³.
 (213) They study the ground state of this system as it adiabatically evolves from a transverse polarized state, and observe that frustration induces extra degeneracy
 (214) They also measure the entanglement in the system, finding a link between frustration and ground-state entanglement.

- (215) This experimental system can be scaled to simulate larger numbers of spins, the ground states of which (for frustrated interactions) cannot be simulated on a classical computer.

International Journal of Modern Physics B Condensed Matter Physics; Statistical Physics; Atomic, Molecular and Optical Physics Volume 04, Issue 02, February 1990 Add to Favorites Download to Citation Manager Citation Alert X. G. WEN, Int. J. Mod. Phys. B, 04, 239 (1990) DOI: 10.1142/S0217979290000139 TOPOLOGICAL ORDERS IN RIGID STATES

- (216) Authors study a new kind of ordering — topological order — in rigid states (the states with no local gapless excitations).
- (217) They concentrate on characterization of the different topological orders.
- (218) As an example we discuss in detail chiral spin states of $2 + 1$ dimensional spin systems.
- (219) Chiral spin states are described by the topological Chern-Simons theories in the continuum limit.
- (220) They show that the topological orders can be characterized by a non-Abelian gauge structure over the moduli space which parametrizes a family of the model Hamiltonians supporting topologically ordered ground states. In $2 + 1$ dimensions, the non-Abelian gauge structure determines possible fractional statistics of the quasi-particle excitations over the topologically ordered ground states.
- (221) The dynamics of the low lying global excitations is shown to be independent of random spatial dependent perturbations.
- (222) The ground state degeneracy and the non-Abelian gauge structures discussed in this paper are very robust, even against those perturbations that break translation symmetry.
- (223) They also discuss the symmetry properties of the degenerate ground states of chiral spin states.
- (224) Authors find that some degenerate ground states of chiral spin states on torus carry non-trivial quantum numbers of the 90° rotation.

Interacting Anyons in Topological Quantum Liquids: The Golden Chain Phys. Rev. Lett. 98, 160409 – Published 20 April 2007 Adrian Feiguin, Simon Trebst, Andreas W. W. Ludwig, Matthias Troyer, Alexei Kitaev, Zhenghan Wang, and Michael H. Freedman

- (225) Authors discuss generalizations of quantum spin Hamiltonians using anyonic degrees of freedom.
- (226) The simplest model for interacting anyons energetically favors neighboring anyons to fuse into the trivial (“identity”) channel, similar to the quantum Heisenberg model favoring neighboring spins to form spin singlets.
- (227) Numerical simulations of a chain of Fibonacci anyons show that the model is critical with a dynamical critical exponent $z=1$, and described by a two-dimensional (2D) conformal field theory with central charge $c=7/10$.
- (228) An exact mapping of the anyonic chain onto the 2D tricritical Ising model is given using the restricted-solid-on-solid representation of the Temperley-Lieb algebra.
- (229) The gaplessness of the chain is shown to have topological origin. DOI: <http://dx.doi.org/10.1103/PhysRevLett.98.160409>

Loop Quantum Gravity Carlo Rovelli

- (230) The problem of finding the quantum theory of the gravitational field, and thus understanding what is quantum spacetime, **is still open**.
- (231) One of the most active of the current approaches is loop quantum gravity.
- (232) Loop quantum gravity is a mathematically well-defined, non-perturbative and background independent quantization of general relativity, with its conventional matter couplings.
- (233) The research in loop quantum gravity forms today a vast area, ranging from mathematical foundations to physical applications.

- (234) Among the most significative results obtained are: (i) the computation of the physical spectra of geometrical quantities such as area and volume; which yields quantitative predictions on Planck-scale physics.
- (235) (ii) A derivation of the Bekenstein-Hawking black hole entropy formula.
- (236) (iii) An intriguing physical picture of the microstructure of quantum physical space, characterized by a polymer-like Planck scale discreteness.
- (237) This discreteness emerges naturally from the quantum theory and provides a mathematically well-defined realization of Wheeler's intuition of a spacetime ``foam".
- (238) Long standing open problems within the approach (lack of a scalar product, over completeness of the loop basis, implementation of reality conditions) have been fully solved.
- (239) The weak part of the approach is the treatment of the dynamics: at present there exist several proposals, which are intensely debated.
- (240) Here, **Carlo Rovelli** provides a general overview of ideas, techniques, results and open problems of this candidate theory of quantum gravity, and a guide to the relevant literature. Journal reference: LivingRev.Rel.1:1, 1998 Cite as: arXiv:gr-qc/9710008

Vertex Operator Solutions of 2d dimensionally Reduced Gravity Denis Bernard, Nicolas Regnault Commun.Math.Phys 210 (2000) 177-201, solv-int/9902017

- (241) **Denis Bernard, Nicolas Regnault** applies algebraic and vertex operator techniques to solve two dimensional reduced vacuum Einstein's equations.
 - (242) This leads to explicit expressions for the coefficients of metrics solutions of the vacuum equations as ratios of determinants.
 - (243) No quadratures are left. These formulas rely on the identification of dual pairs of vertex operators corresponding to dual metrics related by the Kramer-Neugebauer symmetry.
 - (244) **Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin** develop a systematic method for renormalizing the AdS/CFT prescription for computing correlation functions.
 - (245) This involves regularizing the bulk on-shell supergravity action in a covariant way, computing all divergences, adding counterterms to cancel them and then removing the regulator.
 - (246) Authors explicitly work out the case of pure gravity up to six dimensions and of gravity coupled to scalars.
 - (247) The method can also be viewed and visualized as providing a holographic reconstruction of the bulk spacetime metric and of bulk fields on this spacetime, out of conformal field theory data.
 - (248) Knowing which sources are turned on is sufficient in order to obtain an asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order so that all infrared divergences of the on-shell action are obtained.
 - (249) To continue the holographic reconstruction of the bulk fields one needs new CFT data: the expectation value of the dual operator.
 - (250) In particular, in order to obtain the bulk metric one needs to know the expectation value of stress-energy tensor of the boundary theory.
 - (251) They provide completely explicit formulae for the holographic stress-energy tensors up to six dimensions.
 - (252) Authors show that both the gravitational and matter conformal anomalies of the boundary theory are correctly reproduced.
 - (253) They also obtain the conformal transformation properties of the boundary stress-energy tensors.
- Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin.**

- (254) The theory of holographic space-time (HST) generalizes both string theory and quantum field theory.
- (255) It provides a geometric rationale for supersymmetry (SUSY) and a formalism in which super-Poincare invariance follows from Poincare invariance. HST unifies particles and black holes, realizing both as excitations of non-commutative geometrical variables on a holographic screen.
- (256) Compact extra dimensions are interpreted as finite dimensional unitary representations of super-algebras, and have no moduli.
- (257) Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of super-Poincare invariant limits in which the number of holographic degrees of freedom goes to infinity.
- (258) Finite radius de Sitter (dS) spaces have no moduli, and break SUSY with a gravitino mass scaling like $\Lambda^{1/4}$. **T. Banks** presents a holographic theory of inflation and fluctuations.
- (259) The **inflaton field** is an emergent concept, describing the geometry of an underlying HST model, rather than "a field associated with a microscopic string theory".
- (260) **T. Banks** argues that the phrase in quotes is meaningless in the HST formalism. Cite as: arXiv: 1109.2435 [hep-th] **Holographic Space-Time: The Takeaway T. Banks.**
- (261) **T. Banks, W. Fischler** use the formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain a firewall, i.e. an in-falling observer encounters highly excited states at a time much shorter than the light crossing time of the Schwarzschild radius.
- (262) This conclusion is much less dramatic in HST than in the hypothetical models of quantum gravity used in [1].
- (263) In HST there is no dramatic change in particle physics inside the horizon until a time of order the Schwarzschild radius. Report number: UTTG-15-12; TCC-015-12; RUNHETC-2012-17; SCIPP 12/11 arXiv: 1208.4757 [hep-th] **Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler.**
- (264) **Tom Banks** reviews the holographic theory of space-time and its applications to cosmology.
- (265) Much of this has appeared before, but this discussion is more unified and concise.
- (266) He also includes some material on work in progress, whose aim is to understand compactification in terms of finite-dimensional super-algebras. **Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era**
- (267) .The spinorial geometry method of solving Killing spinor equations is reviewed as it applies to six-dimensional (1,0) supergravity.
- (268) In particular, it is explained how the method is used to identify both the fractions of supersymmetry preserved by and the geometry of all supersymmetric backgrounds.
- (269) Then two applications are described to systems that exhibit superconformal symmetry. The first is the proof that some six-dimensional black hole horizons are locally isometric to $AdS_3 \times \Sigma_3$, where Σ_3 is diffeomorphic to S^3 .
- (270) The second one is a description of all supersymmetric solutions of six-dimensional (1,0) superconformal theories and in particular of their brane solitons.
- (271) **M Akyol and G Papadopoulos 2014 Class. Quantum Grav 31 123001 doi:10.1088/0264-9381/31/12/123001 Spinorial geometry, horizons and superconformal symmetry in six dimensions.**
- (272) Field of squeezed states for gravitational-wave (GW) detector enhancement is rapidly maturing. In this review paper, **S S Y Chua¹, B J J Slagmolen, D A Shaddock and D E McClelland** provide an analysis of the field circa 2013.
- (273) They begin by outlining the concept and description of quantum squeezed states.
- (274) This is followed by an overview of how quantum squeezed states can improve GW detection, and the requirements on squeezed states to achieve such enhancement.
- (275) Next, an overview of current technology for producing squeezed states, using atoms, optomechanical methods and nonlinear crystals, is provided.

- (276) Authors finally highlight the milestone squeezing implementation experiments at the GEO600 and LIGO GW detectors. **S S Y Chua et al 2014 Class. Quantum Grav 31 183001 doi:10.1088/0264-9381/31/18/183001 Quantum squeezed light in gravitational-wave detectors**

@@
@@
@@

SECTION ONE

Asymptotic Symmetry And Three-Dimensional Higher Spin Ads Gravity

INTRODUCTION—VARIABLES USED

Journal of High Energy Physics December 2010, 2010:7 Date: 01 Dec 2010 Nonlinear W_∞ as asymptotic symmetry of three-dimensional higher spin AdS gravity Marc Henneaux, Soo-Jong Rey

- (1) **Marc Henneaux, Soo-Jong Rey** investigate the asymptotic symmetry algebra of (e) (2+1)-dimensional higher spin, anti-de Sitter gravity.
- (2) They use the formulation of the theory as a Chern-Simons gauge theory based on (e) the higher spin algebra $hs(1, 1)$.
- (3) Expanding the gauge connection around (e&eb) asymptotically anti-de Sitter spacetime, they specify consistent boundary conditions on (eb) the higher spin gauge fields.
- (4) Authors then study residual gauge transformation (e&eb) the corresponding surface terms and (e&eb) their Poisson bracket algebra.
- (5) They find that the asymptotic symmetry algebra is (=) nonlinearly realized W_∞ algebra with classical central charges.
- (6) They discuss implications of results to (e&eb) quantum gravity and to various situations in string theory. ArXiv ePrint: 1008.4579

Annals of Physics Volume 172, Issue 2, December 1986, Pages 304–347 Black holes in higher dimensional space-times R.C Myers, M.J Perry doi: 10.1016/0003-4916(86)90186-7

- (7) Black hole solutions to Einstein's equations are examined in (eb) asymptotically flat $N + 1$ dimensional space-times.
- (8) First generalizations of Schwarzschild and Reissner-Nordström solutions are examined in a discussion of (e) static black holes in $N + 1$ dimension.

NOTATION

Module One

Marc Henneaux, Soo-Jong Rey investigate the

asymptotic symmetry algebra of (e) (2+1)-dimensional higher spin, anti-de Sitter gravity

G_{13} : Category one of (2+1)-dimensional higher spin, anti-de Sitter gravity

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of asymptotic symmetry algebra

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

They use the

formulation of the theory as a Chern-Simons gauge theory based on (e) the higher spin algebra $hs(1, 1)$

G_{16} : Category one of higher spin algebra $hs(1, 1)$

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of formulation of the theory as a Chern-Simons gauge theory

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Expanding the

gauge connection around (e&eb) asymptotically anti-de Sitter spacetime, they specify consistent boundary conditions on (eb) the higher spin gauge fields

G_{20} : Category one of gauge connection; asymptotically anti-de Sitter spacetime, they specify consistent boundary conditions on (eb) the higher spin gauge fields

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of asymptotically anti-de Sitter spacetime, they specify consistent boundary conditions on (eb) the higher spin gauge fields ;gauge connection

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

gauge connection around asymptotically anti-de Sitter spacetime, they specify consistent boundary conditions on (eb) the higher spin gauge fields

G_{24} : Category one of gauge connection around asymptotically anti-de Sitter spacetime, they specify consistent boundary conditions

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of higher spin gauge fields

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Authors then study

residual gauge transformation (e&eb) the corresponding surface terms and (e&eb) their Poisson bracket algebra

G_{28} : Category one of residual gauge transformation; corresponding surface terms and (e&eb) their Poisson bracket algebra

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of corresponding surface terms and (e&eb) their Poisson bracket algebra ;residual gauge transformation

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

residual gauge transformation of the corresponding surface terms and (e&eb) their Poisson bracket algebra

G_{32} : Category one of residual gauge transformation of the corresponding surface terms; Poisson bracket algebra

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Poisson bracket algebra; residual gauge transformation of the corresponding surface terms

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

They find that the

asymptotic symmetry algebra is (=) nonlinearly realized W_{∞} algebra with classical central charges

G_{36} : Category one of asymptotic symmetry algebra

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of nonlinearly realized W_∞ algebra with classical central charges

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

They discuss implications of

Results of **Nonlinear W_∞ as asymptotic symmetry of three-dimensional higher spin** to (e&eb) quantum gravity and to various situations in string theory. ArXiv ePrint: 1008.4579

G_{40} : Category one of Results of **Nonlinear W_∞ as asymptotic symmetry of three-dimensional higher spin**; quantum gravity and to various situations in string theory

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of quantum gravity and to various situations in string theory; Results of **Nonlinear W_∞ as asymptotic symmetry of three-dimensional higher spin**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Black hole solutions to Einstein's equations are examined in (eb) asymptotically flat $N + 1$ dimensional space-times

G_{44} : Category one of Black hole solutions to Einstein's equations

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of asymptotically flat $N + 1$ dimensional space-times

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$

$$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$$

$$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$$

are Accentuation coefficients

$$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$$

$$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$$

$$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$$

$$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$$

$$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$$

$$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$$

$$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
 $-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
 $-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\begin{aligned}\frac{dG_{36}}{dt} &= (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} & 42 \\ &+ (a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}\end{aligned}$$

Module Numbered Eight

The differential system of this model is now

$$\begin{aligned}\frac{dG_{40}}{dt} &= (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} & 44 \\ \frac{dG_{42}}{dt} &= (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} & 54 \\ &+ (a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor} \\ &- (b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}\end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for

category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for

category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth

detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a''_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a''_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a''_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19}, t) & -(b''_{13})^{(1,1)}(G, t) & -(b''_{20})^{(3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ & -(b''_{36})^{(7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19}, t) & -(b''_{14})^{(1,1)}(G, t) & -(b''_{21})^{(3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ & -(b''_{37})^{(7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19}, t) & -(b''_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ & -(b''_{38})^{(7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21}, t) & +(a''_{16})^{(2,2,2)}(T_{17}, t) & +(a''_{13})^{(1,1,1)}(T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ & +(a''_{36})^{(7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21}, t) & +(a''_{17})^{(2,2,2)}(T_{17}, t) & +(a''_{14})^{(1,1,1)}(T_{14}, t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ & +(a''_{37})^{(7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{22}$$

$+(a'_{20})^{(3)}(T_{21}, t)$, $+(a'_{21})^{(3)}(T_{21}, t)$, $+(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & - (b''_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{36})^{(7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & - (b''_{21})^{(3)}(G_{23}, t) & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{37})^{(7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{21}$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} & - (b''_{22})^{(3)}(G_{23}, t) & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{38})^{(7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{22}$$

$-(b'_{20})^{(3)}(G_{23}, t)$, $-(b'_{21})^{(3)}(G_{23}, t)$, $-(b'_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7)}(G_{39}, t) \text{ are seventh detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8)}(G_{43}, t) \text{ are eight detrition coefficients for}$$

category 1, 2 and 3

$$-(b''_{46})^{(9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for}$$

category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$$(a''_{24})^{(4)}(T_{25}, t), (a''_{25})^{(4)}(T_{25}, t), (a''_{26})^{(4)}(T_{25}, t) \text{ are first augmentation coefficients}$$

category 1, 2 3

$$+(a''_{28})^{(5,5)}(T_{29}, t), +(a''_{29})^{(5,5)}(T_{29}, t), +(a''_{30})^{(5,5)}(T_{29}, t) \text{ are second augmentation}$$

coefficient for category 1, 2 and 3

$$+(a''_{32})^{(6,6)}(T_{33}, t), +(a''_{33})^{(6,6)}(T_{33}, t), +(a''_{34})^{(6,6)}(T_{33}, t) \text{ are third augmentation}$$

coefficient for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1)}(T_{14}, t)$$

are fourth augmentation coefficients for category 1, 2 and 3

$$+(a''_{16})^{(2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2)}(T_{17}, t)$$

are fifth augmentation coefficients for category 1, 2 and 3

$$+(a''_{20})^{(3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3)}(T_{21}, t)$$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for}$$

category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)}(G_{27}, t) & -(b'_{28})^{(5,5)}(G_{31}, t) & -(b'_{32})^{(6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1)}(G, t) & -(b'_{16})^{(2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b'_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)}(G_{27}, t) & -(b'_{29})^{(5,5)}(G_{31}, t) & -(b'_{33})^{(6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1)}(G, t) & -(b'_{17})^{(2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b'_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)}(G_{27}, t) & -(b'_{30})^{(5,5)}(G_{31}, t) & -(b'_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$ are first augmentation coefficients for category 1, 2 and 3

And $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$ are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$ are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$ are sixth augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)}$ are eighth augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} \boxed{(b'_{28})^{(5)}} & \boxed{-(b''_{28})^{(5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} & \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)} & \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} \boxed{(b'_{29})^{(5)}} & \boxed{-(b''_{29})^{(5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} & \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)} & \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} \boxed{(b'_{30})^{(5)}} & \boxed{-(b''_{30})^{(5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)} & \end{array} \right] T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$ are sixth detrition coefficients

for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{32} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} & + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$[-(b''_{32})^{(6)}(G_{35}, t)], [-(b''_{33})^{(6)}(G_{35}, t)], [-(b''_{34})^{(6)}(G_{35}, t)]$ are first detrition coefficients
for category 1, 2 and 3

$[-(b''_{28})^{(5,5,5)}(G_{31}, t)], [-(b''_{29})^{(5,5,5)}(G_{31}, t)], [-(b''_{30})^{(5,5,5)}(G_{31}, t)]$ are second detrition coefficients
for category 1, 2 and 3

$[-(b''_{24})^{(4,4,4)}(G_{27}, t)], [-(b''_{25})^{(4,4,4)}(G_{27}, t)], [-(b''_{26})^{(4,4,4)}(G_{27}, t)]$ are third detrition coefficients
for category 1, 2 and 3

$[-(b''_{13})^{(1,1,1,1,1,1)}(G, t)], [-(b''_{14})^{(1,1,1,1,1,1)}(G, t)], [-(b''_{15})^{(1,1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients
for category 1, 2, and 3

$[-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)], [-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)], [-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition
coefficients for category 1, 2, and 3

$[-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)], [-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)], [-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition
coefficients for category 1, 2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)], [-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)], [-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition
coefficients for category 1, 2, and 3

$[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)], [-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)], [-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$
are eighth detrition coefficients for category 1, 2, and 3

$[-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)], [-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)], [-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & - (b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & - (b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for

category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition

coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition

coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition

coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition

coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition

coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition

coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third

augmentation coefficient for category 1, 2 and 3

$$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) \text{ are seventh}$$

augmentation coefficient for 1,2,3

$$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t), -(b''_{37})^{(7)}(G_{39}, t), -(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t), -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) \text{ are sixth detrition coefficients}$$

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition

coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition

coefficients for category 1, 2 and 3

$$\begin{aligned} & \frac{dG_{44}}{dt} \\ &= (a_{44})^{(9)}G_{45} \\ & - \left[\begin{array}{l} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} & \frac{dG_{45}}{dt} \\ &= (a_{45})^{(9)}G_{44} \\ & - \left[\begin{array}{l} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) + (a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} & \frac{dG_{46}}{dt} \\ &= (a_{46})^{(9)}G_{45} \\ & - \left[\begin{array}{l} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\begin{aligned} \frac{dT_{44}}{dt} &= \\ (b_{44})^{(9)}T_{45} &- \left[\begin{array}{ccc} (b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t) & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\ \frac{dT_{45}}{dt} &= \\ (b_{45})^{(9)}T_{44} &- \left[\begin{array}{ccc} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\ \frac{dT_{46}}{dt} &= \\ (b_{46})^{(9)}T_{45} &- \left[\begin{array}{ccc} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15} \end{aligned}$$

Where $-(b''_{44})^{(9)}(G_{47}, t)$, $-(b''_{45})^{(9)}(G_{47}, t)$, $-(b''_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1$$

111

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22$$

112

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)}$$

113

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition:

114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$:

115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$\begin{aligned} (a''_i)^{(4)}(T_{25}, t) &\leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)} \\ (b''_i)^{(4)}((G_{27}), t) &\leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)} \\ \lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) &= (p_i)^{(4)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) &= (r_i)^{(4)} \end{aligned} \quad 118$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$\begin{aligned} |(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| &\leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)}t} \\ |(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| &< (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)}t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\begin{aligned} \frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] &< 1 \\ \frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] &< 1 \end{aligned}$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$\begin{aligned} (a''_i)^{(5)}(T_{29}, t) &\leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)} \\ (b''_i)^{(5)}((G_{31}), t) &\leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)} \\ \lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) &= (p_i)^{(5)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) &= (r_i)^{(5)} \end{aligned} \quad 123$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$\begin{aligned} |(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| &\leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t} \\ |(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| &< (\hat{k}_{28})^{(5)} \|(G_{31}) - (G_{31})'\| e^{-(\hat{M}_{28})^{(5)} t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(A) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

$$(B) \quad \text{The functions } (a''_i)^{(7)}, (b''_i)^{(7)} \text{ are positive continuous increasing and bounded.}$$

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(C) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(D) \quad \lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ are positive constants and $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T'_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

$$(E) \quad (\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, \text{ are positive constants}$$

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

$$(F) \quad \text{There exists two constants } (\hat{P}_{36})^{(7)} \text{ and } (\hat{Q}_{36})^{(7)} \text{ which together with } (\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} \text{ and } (\hat{B}_{36})^{(7)} \text{ the constants}$$

$(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$
 $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42,$

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(a) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} | (a'_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a'_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a'_{13})^{(1)}$ and $(b'_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(1)}$ and $(b'_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e \left[- \int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)} \right] \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\bar{M}_{13})^{(1)})_1, ((\bar{M}_{13})^{(1)})_2$ and $((\bar{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < (\bar{M}_{13})^{(1)} \text{ it follows } \frac{dG_{14}}{dt} \leq ((\bar{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)} (G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)} (G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[(\widehat{P}_{16})^{(2)} + ((\widehat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[((\widehat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{16})^{(2)} \right] \leq (\widehat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$

197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\bar{A}_{16})^{(2)} + (\bar{P}_{16})^{(2)} (\bar{k}_{16})^{(2)}) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}); (G_{19})^{(2)}, (T_{19})^{(2)} \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\bar{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ .

199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

200

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{ (a'_i)^{(2)} - (a''_i)^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{16})^{(2)})_1, ((\bar{M}_{16})^{(2)})_2$ and $((\bar{M}_{16})^{(2)})_3 :$

201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\bar{M}_{16})^{(2)})$ it follows $\frac{dG_{17}}{dt} \leq ((\bar{M}_{16})^{(2)})_1 - (a'_{17})^{(2)} G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(2)} ((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b'_i)^{(2)} ((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b''_{18})^{(2)} ((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[(\widehat{P}_{20})^{(3)} + ((\widehat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[((\widehat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{20})^{(3)} \right] \leq (\widehat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \}$$

Indeed if we denote

212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : ((\widetilde{G}_{23}), (\widetilde{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}; (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

216

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \} ds_{(20)}]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(3)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{20})^{(3)})_1, ((\bar{M}_{20})^{(3)})_2$ and $((\bar{M}_{20})^{(3)})_3 :$

217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b'_i)^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b''_{22})^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[(\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \}$$

Indeed if we denote

$$\textbf{Definition of } (\widetilde{G_{27}}, \widetilde{T_{27}}) : (\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_{24}^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)} \right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right); (G_{27})^{(2)}, (T_{27})^{(2)}\right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{24})^{(4)})_1, ((\bar{M}_{24})^{(4)})_2$ and $((\bar{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b'_i)^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[(\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric

$$d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) = \\ \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_{28}^{(2)}| \leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ \int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{-(\bar{M}_{28})^{(5)}s_{(28)}} + \\ (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} + \\ G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{1}{(\bar{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\bar{A}_{28})^{(5)} + (\bar{P}_{28})^{(5)} (\bar{k}_{28})^{(5)}) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right); (G_{31})^{(2)}, (T_{31})^{(2)}\right) \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$ and $(\bar{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \} ds_{(28)}} \geq 0 \\ T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{28})^{(5)})_1, ((\bar{M}_{28})^{(5)})_2$ and $((\bar{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)}G_{29}$ and by integrating

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) = \\ \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$|\tilde{G}_{32}^{(1)} - \tilde{G}_{32}^{(2)}| \leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ \int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{-(\bar{M}_{32})^{(6)}s_{(32)}} + \\ (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} + \\ G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{1}{(\bar{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\bar{A}_{32})^{(6)} + (\bar{P}_{32})^{(6)} (\bar{k}_{32})^{(6)}) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}$ and $(\bar{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0 \\ T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{32})^{(6)})_1, ((\bar{M}_{32})^{(6)})_2$ and $((\bar{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying

Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric

258

$$d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right), \left((G_{39})^{(2)}, (T_{39})^{(2)}\right)\right) = \sup\left\{\max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}\right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}, \widetilde{T_{39}}) : (\widetilde{G_{39}}, \widetilde{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} |(a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}\right) d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right); (G_{39})^{(2)}, (T_{39})^{(2)}\right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$:

262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$G_{36} < ((\widehat{M}_{36})^{(7)})$ it follows $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266

$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

268

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\hat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{40})^{(8)} \right] \leq (\hat{Q}_{40})^{(8)}$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d \left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widehat{G_{43}}, \widehat{T_{43}})$: $(\widehat{G_{43}}, \widehat{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} \} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} &((a_{40})^{(8)} + (a'_{40})^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}) d \left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)}) \right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1, ((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$$G_{40} < ((\widehat{M}_{40})^{(8)}) \text{ it follows } \frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)}G_{41} \text{ and by integrating}$$

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b_{42}'')^{(8)}((G_{43})(t), t(t), t) = (b_{42}')^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[\left((\hat{Q}_{44})^{(9)} + T_j^0 \right) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \}$$

Indeed if we denote

Definition of $(\bar{G}_{47}, \bar{T}_{47}) : (\bar{G}_{47}, \bar{T}_{47}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a_{44}')^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a_{44}'')^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a_{44}'')^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a_{44}'')^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a_{44}')^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a_{44}'')^{(9)}$ and $(b_{44}'')^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a_i')^{(9)} - (a_i'')^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \} ds_{(44)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1, ((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$G_{44} < ((\widehat{M}_{44})^{(9)})_1$ it follows $\frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b'_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
 $(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, if $(v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}$, if $(v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}$,

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}$, if $(\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously

284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, if $(u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}$, if $(u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}$, if $(\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \leq$$

286

$$\frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$$

287

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

288

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq$$

289

$$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^0e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying
 $-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$ 294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$ 305

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$ 306

and $(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(\nu_1)^{(3)} > 0, (\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(\nu^{(3)})^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{\nu}_1)^{(3)} > 0, (\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(\nu^{(3)})^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}, \text{ if } (\nu_0)^{(3)} < (\nu_1)^{(3)}$$

$$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}, \text{ if } (\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)},$$

$$\text{and } \boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}, \text{ if } (\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(s_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(s_1)^{(3)}t}$$

323

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right.$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(\nu_1)^{(5)} > 0, (\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{\nu}_1)^{(5)}, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{\nu}_1)^{(5)} > 0, (\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)} (u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$
Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and $(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \right.$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}}$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:-

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)} (m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)} (\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) - (b''_{41})^{(8)}(G_{43}, t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((s_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(s_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t} \right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$
Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(s_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_2)^{(9)}} G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(s_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((s_1)^{(9)} - (p_{44})^{(9)} - (s_2)^{(9)})} \left[e^{((s_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(s_2)^{(9)}t} \right] + G_{46}^0 e^{-(s_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((s_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(s_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-

$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad (\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{c})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}) t]}}{1 + (\bar{c})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}) t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)} G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)} (T_{17}, t) \right) - (a_{17}'')^{(2)} (T_{17}, t) v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

Definition of $v^{(2)}$:-

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

388

It follows

389

$$- \left((a_{17})^{(2)} (v^{(2)})^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)} (v^{(2)})^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof: From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- 399

$$v^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (C)^{(3)}(\bar{\nu}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)})t]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain

404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$- \left((a_{25})^{(4)}(\nu^{(4)})^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} \right) \leq \frac{d\nu^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(\nu^{(4)})^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{[-(a_{25})^{(4)}((\nu_1)^{(4)} - (\nu_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((\nu_1)^{(4)} - (\nu_0)^{(4)})t]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)} , \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)} , \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$-\left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)}\right) \leq \frac{dv^{(5)}}{dt} \leq -\left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \left(v_0\right)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (\bar{C})^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad (\bar{C})^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad (\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (\bar{C})^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \left(v_0\right)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of $(u_0)^{(5)}$.**

Proof : From global equations we obtain

412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:-

$$\nu^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left((a_{33})^{(6)}(\nu^{(6)})^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} \right) \leq \frac{d\nu^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(\nu^{(6)})^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get

413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case,

414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_2)^{(6)})t]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case**.

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a_{36}')^{(7)} - (a_{37}')^{(7)} + (a_{36}'')^{(7)} (T_{37}, t) \right) - (a_{37}'')^{(7)} (T_{37}, t) v^{(7)} - (a_{37})^{(7)} v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$- \left((a_{37})^{(7)} (v^{(7)})^2 + (\sigma_2)^{(7)} v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)} (v^{(7)})^2 + (\sigma_1)^{(7)} v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_0)^{(7)}) t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_0)^{(7)}) t]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)} , \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)} , \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition

$(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof: From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

For $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner, we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case .**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)} (a_{14}')^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)} (a_{14}')^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} + (a_{13})^{(1)} (p_{13})^{(1)} + (a_{14}')^{(1)} (p_{14})^{(1)} + (p_{13})^{(1)} (p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{28})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{32})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{36})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{40})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied, then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution, which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution, which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution, which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0$	475
$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0$	476
$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0$	477
$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0$	478
$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(a) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that

there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42})$, G_{40}, G_{42} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and

512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

G_{25}^* given by $\varphi(G_{27}) = 0, T_{25}^*$ given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]} \\ T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]} \\ T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]} \\ T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]} \\ T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]} \quad 523$$

Finally we obtain the unique solution of 89 to 99 523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19})^* = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a_{16}')^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a_{17}')^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a_{18}')^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b_{16}')^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b_{17}')^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b_{18}')^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23})^* = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a_{20}')^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a_{21}')^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a_{22}')^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b_{20}')^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b_{21}')^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22}(s_{(21)(j)}T_{21}^*G_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b_{22}')^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(22)(j)}T_{22}^*G_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a_{24}')^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a_{25}')^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a_{26}')^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b_{24}')^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b_{25}')^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b_{26}')^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a_{28}')^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a_{29}')^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a_{30}')^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b_{28}')^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)} T_{29}^* G_j) \quad 561$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)} T_{30}^* G_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 564

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}} (T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j} ((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^* T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^* T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^* T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)} T_{32}^* G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)} T_{33}^* G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)} T_{34}^* G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}} (T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial (b_i'')^{(7)}}{\partial G_j} ((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^* \mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^* \mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^* \mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^* \mathbb{G}_j \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^* \mathbb{G}_j \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^* \mathbb{G}_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^* \mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^* \mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^* \mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(40)(j)})T_{40}^* \mathbb{G}_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)})T_{41}^* \mathbb{G}_j \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)})T_{42}^* \mathbb{G}_j \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a_{44}')^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad 586 \text{ B}$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a_{45}')^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad 586 \text{ C}$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a_{46}')^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad 586 \text{ D}$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b_{44}')^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)}T_{44}^*\mathbb{G}_j) \quad 586 \text{ E}$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b_{45}')^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)}T_{45}^*\mathbb{G}_j) \quad 586 \text{ F}$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b_{46}')^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)}T_{46}^*\mathbb{G}_j) \quad 586 \text{ G}$$

The characteristic equation of this system is

587

$$\begin{aligned} & ((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}) \\ & \left[((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right] \\ & \left(((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^* \right) \\ & + \left(((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^* \right) \\ & \left(((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^* \right) \\ & \left(((\lambda)^{(1)})^2 + ((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & \left(((\lambda)^{(1)})^2 + ((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & + \left(((\lambda)^{(1)})^2 + ((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\ & + ((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\ & \left. \left(((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \right\} = 0 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \right) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ (\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \} \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \} \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & +
 \end{aligned}$$

$$\begin{aligned}
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & + \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ (\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)} \} \\
 & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 & \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 & \left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 & + \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & \left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & + \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWO

String Theory, Black Holes And Asymptotic Symmetries

INTRODUCTION—VARIABLES USED

Annals of Physics Volume 172, Issue 2, December 1986, Pages 304–347 Black holes in higher dimensional space-times R.C Myers, M.J Perry doi: 10.1016/0003-4916(86)90186-7

- (1) Then a new family of solutions is found which describe (eb) spinning black holes in (eb) higher dimensional space-times.
- (2) In many respects these new solutions are (=) similar to the familiar Kerr and Schwarzschild metrics which are recovered for (e) $N = 3$.
- (3) One exceptional case though is that for $N \geq 5$, black holes with a fixed mass may have (e) arbitrarily large angular momentum.

Journal of High Energy Physics November 2010, 2010:7 Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields A. Campoleoni, S. Fredenhagen, S. Pfenninger, S. Theisen

- (4) **A. Campoleoni, S. Fredenhagen, S. Pfenninger, S. Theisen** discuss the emergence of (e) W -algebras as (=) asymptotic symmetries of higher-spin gauge theories coupled to (e&eb) three-dimensional Einstein gravity with (e&eb) a negative cosmological constant.
- (5) **They** focus on models involving a finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to gravity.
- (6) Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to gravity is described by (e) a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W_3 -algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant.
- (7) Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to gravity is described by (e) a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W_3 -algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

String theory and black holes Phys. Rev. D 44, 314 – Published 15 July 1991 Edward Witten

- (8) An exact conformal field theory describing a black hole in (eb) two-dimensional space-time is found as (=) an $SL(2, \mathbb{R})/U(1)$ gauged Wess-Zumino-Witten model.
- (9) For $k=9/4$, the conformal field theory can be regarded as (=) a classical solution of the same system that is probed in the $c=1$ matrix model.

NOTATION

Module One

First generalizations of Schwarzschild and Reissner-Nordstrøm solutions are examined in a discussion of (e) static black holes in $N + 1$ dimension.

G_{13} : Category one of First generalizations of Schwarzschild and Reissner-Nordstrøm solutions; static black holes in $N + 1$ dimension

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of static black holes in $N + 1$ dimension; First generalizations of Schwarzschild and Reissner-Nordstrøm solutions

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Then a

new family of solutions is found which describe (eb) spinning black holes in (eb) higher dimensional space-times

G_{16} : Category one of new family of solutions

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of spinning black holes in (eb) higher dimensional space-times

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

spinning black holes in (eb) higher dimensional space-times

G_{20} : Category one of spinning black holes

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of higher dimensional space-times

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

In many respects these

new solutions are (=) similar to the familiar Kerr and Schwarzschild metrics which are recovered for (e) $N =$

G_{24} : Category one of new solutions

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of similar to the familiar Kerr and Schwarzschild metrics which are recovered for (e) $N = 3$

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

One exceptional case though is that

black holes with a fixed mass for $N \geq 5$ may have (e) arbitrarily large angular momentum

G_{28} : Category one of arbitrarily large angular momentum

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of black holes with a fixed mass for $N \geq 5$

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Campoleoni, S. Fredenhagen, S. Pfenninger, S. Theisen discuss the

emergence of W-algebras as (=) asymptotic symmetries of higher-spin gauge theories coupled to (e&eb) three-dimensional Einstein gravity with (e&eb) a negative cosmological constant

G_{32} : Category one of emergence of W-algebras

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of asymptotic symmetries of higher-spin gauge theories coupled to (e&eb) three-dimensional Einstein gravity with (e&eb) a negative cosmological constant

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

emergence of W-algebras as asymptotic symmetries of higher-spin gauge theories coupled to (e&eb) three-dimensional Einstein gravity with (e&eb) a negative cosmological constant

G_{36} : Category one of emergence of W-algebras as asymptotic symmetries of higher-spin gauge theories; three-dimensional Einstein gravity with (e&eb) a negative cosmological constant

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of three-dimensional Einstein gravity with (e&eb) a negative cosmological constant ;emergence of W-algebras as asymptotic symmetries of higher-spin gauge theories

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

emergence of W-algebras as asymptotic symmetries of higher-spin gauge theories coupled to three-dimensional Einstein gravity with (e&eb) a negative cosmological constant

G_{40} : Category one of emergence of W-algebras as asymptotic symmetries of higher-spin gauge theories coupled to three-dimensional Einstein gravity; **negative** cosmological constant

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of negative cosmological constant; emergence of W-algebras as asymptotic symmetries of higher-spin gauge theories coupled to three-dimensional Einstein gravity

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

They focus on models involving a

finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to gravity.

G_{44} : Category one of finite number of bosonic higher-spin fields; coupling of a spin-3 field to gravity.

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of coupling of a spin-3 field to gravity.; finite number of bosonic higher-spin fields

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \end{aligned}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}), t) =$ First detritions factor

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a'_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a'_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a'_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t), +(a''_{21})^{(3)}(T_{21}, t), +(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{c} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients

for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for

category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients

for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition

coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition

coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation

coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation

coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \begin{bmatrix} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \begin{bmatrix} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

96

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition:

99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31} - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

128

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(G) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(H) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(I) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(J) \quad \lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(K) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(L) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(b) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right] \quad 181$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})$ it follows $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b''_{22})^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

$$\textbf{Definition of } (\widetilde{G_{31}}, \widetilde{T_{31}}): (\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a'_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} , \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\textbf{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-
$$v^{(2)} = \frac{G_{16}}{G_{17}}$$
 388

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$ $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)} , \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)} , \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)} (v_2)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)} (T_{45}, t) \right) - (a_{45}'')^{(9)} (T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(b) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]}, \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}, \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]}, \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]}, \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]}, \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]}, \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}) T_{13}^* \mathbb{G}_j \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}) T_{14}^* \mathbb{G}_j \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}) T_{15}^* \mathbb{G}_j \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ \left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \right. \\
 &\left. \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 &+ \left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \right. \\
 &\left. \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THREE

SL (3) × SL (3) Chern-Simons Theory And Its Asymptotic Symmetry Algebra

INTRODUCTION—VARIABLES USED

Journal of High Energy Physics November 2010, 2010:7 Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields A. Campoleoni, S. Fredenhagen, S. Pfenninger, S. Theisen

- (1) Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to gravity is described by (e) a SL (3) × SL (3) Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant.
- (2) Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to gravity is described by (e) a SL (3) × SL (3) Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

String theory and black holes Phys. Rev. D 44, 314 – Published 15 July 1991 Edward Witten

- (3) An exact conformal field theory describing a black hole in (eb) two-dimensional space-time is found as (=) an SL (2, openR)/U (1) gauged Wess-Zumino-Witten model.
- (4) For k=9/4, the conformal field theory can be regarded as (=) a classical solution of the same system that is probed in the c=1 matrix model.

NOTATION

Module One

Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by (e) the coupling of a spin-3 field to (e&eb) gravity is described by (e) a SL (3) × SL (3) Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{13} : Category one of coupling of a spin-3 field to (e&eb) gravity is described by (e) a SL (3) × SL (3) Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Models involving a finite number of bosonic higher-spin fields, and especially on the example

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to (e&eb) gravity is described by (e) a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{16} : Category one of Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field; gravity is described by (e) a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of gravity is described by (e) a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant; Models involving a finite number of bosonic higher-spin fields, and especially on the example coupling of a spin-3 field

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by (e) a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{20} : Category one of $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by (=) two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{24} : Category one of Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by two copies of the classical W3-algebra with (e& eb) central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{28} : Category one of Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by two copies of the classical W3-algebra; central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant ;Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by two copies of the classical W3-algebra

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by two copies of the classical W3-algebra with central charge the one computed by Brown and Henneaux in (eb) pure gravity with negative cosmological constant

G_{32} : Category one of Models involving a finite number of bosonic higher-spin fields, and especially on the example provided by the coupling of a spin-3 field to gravity is described by a $SL(3) \times SL(3)$ Chern-Simons theory and its asymptotic symmetry algebra is given by two copies of the classical W3-algebra with central charge the one computed

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of pure gravity with negative cosmological constant

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

An

exact conformal field theory describing a black hole in (eb) two-dimensional space-time is found as (=) an $SL(2, \text{openR})/U(1)$ gauged Wess-Zumino-Witten model

G_{36} : Category one of exact conformal field theory

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of black hole in (eb) two-dimensional space-time is found as (=) an $SL(2, \text{openR})/U(1)$ gauged Wess-Zumino-Witten model

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

exact conformal field theory describing a black hole in two-dimensional space-time is found as (=) an $SL(2, \text{openR})/U(1)$ gauged Wess-Zumino-Witten model

G_{40} : Category one of exact conformal field theory describing a black hole in two-dimensional space-time

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of SL (2, openR)/U (1) gauged Wess-Zumino-Witten model

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

conformal field theory for $k=9/4$ can be regarded as (=) a classical solution of the same system that is probed in the $c=1$ matrix model

G_{44} : Category one of conformal field theory for $k=9/4$

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of classical solution of the same system that is probed in the $c=1$ matrix model (systemic differentiation or take all the three stratifications as equivalent).

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\begin{aligned}\frac{dT_{40}}{dt} &= (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} & 54 \\ + (a''_{44})^{(9)}(T_{45}, t) &= \text{First augmentation factor} \\ - (b''_{44})^{(9)}((G_{47}), t) &= \text{First detrition factor}\end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$, $(a''_{21})^{(3,3)}(T_{21}, t)$, $(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7)}(T_{37}, t)$, $(a''_{37})^{(7,7)}(T_{37}, t)$, $(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8)}(T_{41}, t)$, $(a''_{41})^{(8,8)}(T_{41}, t)$, $(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad -(b'_{16})^{(2,2)}(G_{19}, t) \quad -(b'_{20})^{(3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4)}(G_{27}, t) \quad -(b'_{28})^{(5,5,5,5)}(G_{31}, t) \quad -(b'_{32})^{(6,6,6,6)}(G_{35}, t) \\ -(b'_{36})^{(7,7)}(G_{39}, t) \quad -(b'_{40})^{(8,8)}(G_{43}, t) \quad -(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{c} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad -(b'_{17})^{(2,2)}(G_{19}, t) \quad -(b'_{21})^{(3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4)}(G_{27}, t) \quad -(b'_{29})^{(5,5,5,5)}(G_{31}, t) \quad -(b'_{33})^{(6,6,6,6)}(G_{35}, t) \\ -(b'_{37})^{(7,7)}(G_{39}, t) \quad -(b'_{41})^{(8,8)}(G_{43}, t) \quad -(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad -(b'_{18})^{(2,2)}(G_{19}, t) \quad -(b'_{22})^{(3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4)}(G_{27}, t) \quad -(b'_{30})^{(5,5,5,5)}(G_{31}, t) \quad -(b'_{34})^{(6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7)}(G_{39}, t) \quad -(b'_{42})^{(8,8)}(G_{43}, t) \quad -(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17}, t)] \quad +(a'_{13})^{(1,1)}(T_{14}, t) \quad +(a'_{20})^{(3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4)}(T_{25}, t) \quad +(a'_{28})^{(5,5,5,5,5)}(T_{29}, t) \quad +(a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a'_{36})^{(7,7,7)}(T_{37}, t) \quad +(a'_{40})^{(8,8,8)}(T_{41}, t) \quad +(a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}[+(a''_{17})^{(2)}(T_{17}, t)] \quad +(a'_{14})^{(1,1)}(T_{14}, t) \quad +(a'_{21})^{(3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4)}(T_{25}, t) \quad +(a'_{29})^{(5,5,5,5,5)}(T_{29}, t) \quad +(a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a'_{37})^{(7,7,7)}(T_{37}, t) \quad +(a'_{41})^{(8,8,8)}(T_{41}, t) \quad +(a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b'_{20})^{(3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b'_{21})^{(3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{37})^{(7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b'_{22})^{(3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{38})^{(7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients

for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients

for category 1,2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for

category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for

category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category

1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)}(T_{21}, t)$, $+(a'_{21})^{(3)}(T_{21}, t)$, $+(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients

for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients

for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation

coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation

coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients

for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} [- (b''_{20})^{(3)}(G_{23}, t)] & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} [- (b''_{21})^{(3)}(G_{23}, t)] & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} [- (b''_{22})^{(3)}(G_{23}, t)] & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$[- (b''_{20})^{(3)}(G_{23}, t)]$, $[- (b''_{21})^{(3)}(G_{23}, t)]$, $[- (b''_{22})^{(3)}(G_{23}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[- (b'_{16})^{(2,2,2)}(G_{19}, t)]$, $[- (b'_{17})^{(2,2,2)}(G_{19}, t)]$, $[- (b'_{18})^{(2,2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[- (b'_{13})^{(1,1,1)}(G, t)]$, $[- (b'_{14})^{(1,1,1)}(G, t)]$, $[- (b'_{15})^{(1,1,1)}(G, t)]$ are third detrition coefficients for category 1, 2 and 3

$[- (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)]$, $[- (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)]$, $[- (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[- (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)]$, $[- (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)]$, $[- (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[- (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)]$, $[- (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)]$, $[- (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[- (b''_{36})^{(7,7,7,7,7)}(G_{39}, t)]$, $[- (b''_{37})^{(7,7,7,7,7)}(G_{39}, t)]$, $[- (b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[- (b''_{40})^{(8,8,8,8)}(G_{43}, t)]$, $[- (b''_{41})^{(8,8,8,8)}(G_{43}, t)]$, $[- (b''_{42})^{(8,8,8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[- (b''_{46})^{(9,9,9)}(G_{47}, t)]$, $[- (b''_{45})^{(9,9,9)}(G_{47}, t)]$, $[- (b''_{44})^{(9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} [+ (a''_{24})^{(4)}(T_{25}, t)] & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} [+ (a''_{25})^{(4)}(T_{25}, t)] & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a'_{30})^{(5,5)}(T_{29}, t) & + (a'_{34})^{(6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t), (a'_{25})^{(4)}(T_{25}, t), (a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a'_{28})^{(5,5)}(T_{29}, t), +(a'_{29})^{(5,5)}(T_{29}, t), +(a'_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6)}(T_{33}, t), +(a'_{33})^{(6,6)}(T_{33}, t), +(a'_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1)}(T_{14}, t), +(a'_{14})^{(1,1,1,1)}(T_{14}, t), +(a'_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2)}(T_{17}, t), +(a'_{17})^{(2,2,2,2)}(T_{17}, t), +(a'_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3)}(T_{21}, t), +(a'_{21})^{(3,3,3,3)}(T_{21}, t), +(a'_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a'_{46})^{(9,9,9,9)}(T_{45}, t), +(a'_{45})^{(9,9,9,9)}(T_{45}, t), +(a'_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a'_{24})^{(4,4)}(T_{25}, t) & + (a'_{32})^{(6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a'_{25})^{(4,4)}(T_{25}, t) & + (a'_{33})^{(6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a'_{26})^{(4,4)}(T_{25}, t) & + (a'_{34})^{(6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a'_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a'_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a'_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}[-(b''_{28})^{(5)}(G_{31}, t)] & -(b''_{24})^{(4,4)}(G_{27}, t) & -(b''_{32})^{(6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}[-(b''_{29})^{(5)}(G_{31}, t)] & -(b''_{25})^{(4,4)}(G_{27}, t) & -(b''_{33})^{(6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}[-(b''_{30})^{(5)}(G_{31}, t)] & -(b''_{26})^{(4,4)}(G_{27}, t) & -(b''_{34})^{(6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)}[+(a''_{32})^{(6)}(T_{33}, t)] & +(a''_{28})^{(5,5,5)}(T_{29}, t) & +(a''_{24})^{(4,4,4)}(T_{25}, t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)}[+(a''_{33})^{(6)}(T_{33}, t)] & +(a''_{29})^{(5,5,5)}(T_{29}, t) & +(a''_{25})^{(4,4,4)}(T_{25}, t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients

for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients

for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients

for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition

coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{36})^{(7)}(T_{37}, t)$, $(a''_{37})^{(7)}(T_{37}, t)$, $(a''_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) & + (a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)} + (a_{41}'')^{(8)}(T_{41}, t) & + (a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)} + (a_{42}'')^{(8)}(T_{41}, t) & + (a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41}, t)$, $+(a_{41}'')^{(8)}(T_{41}, t)$, $+(a_{42}'')^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

96

$$= (a_{44})^{(9)}G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}[+(a''_{45})^{(9)}(T_{45}, t)] & +(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & +(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}[+(a''_{46})^{(9)}(T_{37}, t)] & +(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & +(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}[-(b''_{44})^{(9)}(G_{47}, t)] & -(b'_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t) & -(b'_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) \quad - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) \quad - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47}, t)$, $-(b''_{45})^{(9)}(G_{47}, t)$, $-(b''_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b'_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a'_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b'_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition:

99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$:

106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition:

114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21}' - T_{21}| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23}' - G_{23}| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$:

115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26$$

117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

118

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition:

119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$:

120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30$$

122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

123

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

128

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33}' - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(M) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(N) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(O) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(P)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ are positive constants and $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37}' - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(Q) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(R) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions

147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(c) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$| (G_{19})^{(1)} - (G_{19})^{(2)} | e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e \left[- \int_0^t \{ (a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right] \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) &d((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on
 (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t}$$

376

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq$$

377

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $v^{(2)} = \frac{G_{16}}{G_{17}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (\bar{v}_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(c) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42}(s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FOUR

Band Offsets In Znse-Znxsxse1-X Strained-Layer Superlattices

INTRODUCTION—VARIABLES USED

String theory and black holes Phys. Rev. D 44, 314 – Published 15 July 1991 Edward Witten

- (1) The conformal field theory governing (e&eb) the space-time is regular at (eb) the Riemannian singularity, but it appears that generic perturbations blow up there.
- (2) It is argued that the **end point** of the Hawking black-hole evaporation is (=) the standard space-time of the c=1 matrix model, which should be regarded as (=) an analog of the extreme Reissner-Nordström black hole of (e) four-dimensional general relativity.

- (3) The $c=1$ model is thus a model of the quantum mechanics of matter interacting with (e&eb) a black hole. DOI: <http://dx.doi.org/10.1103/PhysRevD.44.314>

Optical characterization and band offsets in ZnSe-ZnSxSe1-x strained-layer superlattices Phys. Rev. B 38, 1417 – Published 15 July 1988; Erratum Phys. Rev. B 43, 1830 (1991)Khalid Shahzad, Diego J. Olego, and Chris G. Van de Walle

- (4) Photoluminescence and excitation experiments were carried out to study effects of the strain and carrier confinement in (eb) ZnSe-ZnSxSe1-x strained-layer superlattices (SLS's) grown by (e) molecular-beam epitaxy on GaAs substrates.
- (5) For the case where the total thickness of the SLS is very small (1000 Å), the structure grows (eb) pseudomorphically to the buffer layer.
- (6) The ZnSe well layers are not strained and all the blue shift in the optical spectra is attributed to (e) the carrier confinement effects only.
- (7) At the other extreme, for the case of a SLS with very large total thickness ($\sim 4 \mu\text{m}$), they show that it can be treated as (=) free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers under (e&eb) biaxial tension.
- (8) In the cases of intermediate total thicknesses, they show that SLS's are not fully relaxed to their equilibrium states by (e) measuring the strains directly in the ZnSe well layers.
- (9) Empirical values for the band offsets are obtained from (e) the analysis of the optical response as (=) a function of the sample parameters.

NOTATION

Module One

conformal field theory governing (e&eb) the space-time is regular at (eb) the Riemannian singularity, but it appears that generic perturbations blow up there

G_{13} : Category one of conformal field theory; space-time is regular at (eb) the Riemannian singularity, but it appears that generic perturbations blow up there

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of space-time is regular at (eb) the Riemannian singularity, but it appears that generic perturbations blow up there; conformal field theory

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

conformal field theory governing the space-time is regular at the Riemannian singularity, but it appears that generic perturbations blow up there

G_{16} : Category one of conformal field theory governing the space-time

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Riemannian singularity, but it appears that generic perturbations blow up there

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

It is argued that the

end point of the Hawking black-hole evaporation is (=) the standard space-time of the $c=1$ matrix model, which should be regarded as (=) an analog of the extreme Reissner-Nordström black hole of (e) four-dimensional general relativity

G_{20} : Category one of **end point** of the Hawking black-hole evaporation

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of standard space-time of the $c=1$ matrix model, which should be regarded as (=) an analog of the extreme Reissner-Nordström black hole of (e) four-dimensional general relativity

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

end point of the Hawking black-hole evaporation is the standard space-time of the $c=1$ matrix model, which should be regarded as (=) an analog of the extreme Reissner-Nordström black hole of (e) four-dimensional general relativity

G_{24} : Category one of **end point** of the Hawking black-hole evaporation is the standard space-time of the $c=1$ matrix model

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of analog of the extreme Reissner-Nordström black hole of (e) four-dimensional general relativity

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

end point of the Hawking black-hole evaporation is the standard space-time of the $c=1$ matrix model, which should be regarded as an analog of the extreme Reissner-Nordström black hole of (e) four-dimensional general relativity

G_{28} : Category one of four-dimensional general relativity

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of **end point** of the Hawking black-hole evaporation is the standard space-time of the $c=1$ matrix model, which should be regarded as an analog of the extreme Reissner-Nordström black hole

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

$c=1$ model is thus a model of the quantum mechanics of matter interacting with (e&eb) a black hole. DOI:
<http://dx.doi.org/10.1103/PhysRevD.44.314>

G_{32} : Category one of $c=1$ model is thus a model of the quantum mechanics of matter; black holes

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of black holes; $c=1$ model is thus a model of the quantum mechanics of matter

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Photoluminescence and excitation experiments were carried out to study effects of the strain and carrier confinement in (eb) ZnSe-ZnSxSe1-x strained-layer superlattices (SLS's) grown by (e) molecular-beam epitaxy on GaAs substrates

G_{36} : Category one of Photoluminescence and excitation experiments were carried out to study effects of the strain and carrier confinement

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of ZnSe-ZnSxSe1-x strained-layer superlattices (SLS's) grown by (e) molecular-beam epitaxy on GaAs substrates

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Photoluminescence and excitation experiments were carried out to study effects of the strain and carrier confinement in ZnSe-ZnSxSe1-x strained-layer superlattices (SLS's) grown by (e) molecular-beam epitaxy

on GaAs substrates

G_{40} : Category one of molecular-beam epitaxy on GaAs substrates

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Photoluminescence and excitation experiments were carried out to study effects of the strain and carrier confinement in ZnSe-ZnS_xSe_{1-x} strained-layer superlattices (SLS's)

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Structure (for the case where the total thickness of the SLS is very small (1000 Å)) grows (eb) pseudomorphically to the buffer layer

G_{44} : Category one of Structure (for the case where the total thickness of the SLS is very small (1000 Å))

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of pseudomorphically to the buffer layer

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20\end{aligned}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & - (b''_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b''_{22})^{(3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{38})^{(7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) + (a''_{24})^{(4,4)}(T_{25}, t) + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T_{29}'| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} \|(G_{31}) - (G_{31})'\| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(S) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(T) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(U) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(V) \quad \lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(W) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(X) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(d) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) &d((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a'_{13})^{(1)}(T_{14}, t) + (a'_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b'_{13})^{(1)}(G, t) - (b'_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-

$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(d) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b_i'')^{(5)}}{\partial G_j}(G_{31}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIVE

Quantum Spin Dynamics (QSD)

INTRODUCTION—VARIABLES USED

Optical characterization and band offsets in ZnSe-ZnSxSe1-x strained-layer superlattices Phys. Rev. B 38, 1417 – Published 15 July 1988; Erratum Phys. Rev. B 43, 1830 (1991) Khalid Shahzad, Diego J. Olego, and Chris G. Van de Walle

- (1) The ZnSe well layers are not strained and all the blue shift in the optical spectra is attributed to (e) the carrier confinement effects only.
- (2) At the other extreme, for the case of a SLS with very large total thickness ($\sim 4 \mu\text{m}$), they show that

it can be treated as (=) free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers under (e&eb) biaxial tension.

- (3) In the cases of intermediate total thicknesses, they show that SLS's are not fully relaxed to their equilibrium states by (e) measuring the strains directly in the ZnSe well layers.
- (4) Empirical values for the band offsets are obtained from (e) the analysis of the optical response as (=) a function of the sample parameters.

T Thiemann 1998 Class Quantum Grav.15 875 doi:10.1088/0264-9381/15/4/012 Quantum spin dynamics (QSD): II. The kernel of the Wheeler - DeWitt constraint operator

- (5) Authors determine the complete and rigorous kernel of the Wheeler - DeWitt constraint operator for (e) four-dimensional, Lorentzian, non-perturbative, canonical vacuum quantum gravity in the continuum.
- (6) They construct complete and rigorous kernel of the Wheeler - DeWitt constraint operator for (e) the non-symmetric version of the operator constructed previously in this series.
- (7) They also construct a symmetric, regulated constraint operator. For the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of (e) the Wick transform from (e) the Euclidean to the Lorentzian regime they prove (eb) existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.
- (8) Both constraint operators evaluated at unit lapse as well as the generator of the Wick transform can be shown to have (e) regulator-independent and symmetric duals on (eb) the diffeomorphism-invariant Hilbert space.
- (9) Finally, comment is on the status of the Wick rotation transform in the light of the present results that give an intuitive description of (e) the action of the Hamiltonian constraint.

NOTATION

Module One

ZnSe well layers are not strained and all the blue shift in the optical spectra is attributed to (e) the carrier confinement effects only

G_{13} : Category one of carrier confinement effects only

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of ZnSe well layers are not strained and all the blue shift in the optical spectra

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

At the other extreme, for the case of a SLS with very large total thickness ($\sim 4 \mu\text{m}$), they show that it can be treated as (=) free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers under (e&eb) biaxial tension

G_{16} : Category one of SLS with very large total thickness

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers under (e&eb) biaxial tension

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers under (e&eb) biaxial tension

G_{20} : Category one of free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers; biaxial tension

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of biaxial tension ;free-standing with ZnSe layers under biaxial compression and ZnSxSe1-x layers

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

they show that

SLS's in the cases of intermediate total thicknesses are not fully relaxed to their equilibrium states by (e) measuring the strains directly in the ZnSe well layers

G_{24} : Category one of measuring the strains directly in the ZnSe well layers

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of SLS's in the cases of intermediate total thicknesses are not fully relaxed to their equilibrium states

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Empirical values for the band offsets are obtained from (e) the analysis of the optical response as (=) a function of the sample parameters

G_{28} : Category one of analysis of the optical response as (=) a function of the sample parameters

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Empirical values for the band offsets

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Empirical values for the band offsets are obtained from the analysis of the optical response as (=) a function of the sample parameters

G_{32} : Category one of Empirical values for the band offsets are obtained from the analysis of the optical response

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of function of the sample parameters

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Authors determine the

complete and rigorous kernel of the Wheeler - DeWitt constraint operator for (e) four-dimensional, Lorentzian, non-perturbative, canonical vacuum quantum gravity in the continuum

G_{36} : Category one of four-dimensional, Lorentzian, non-perturbative, canonical vacuum quantum gravity in the continuum

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of complete and rigorous kernel of the Wheeler - DeWitt constraint operator

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

They construct

complete and rigorous kernel of the Wheeler - DeWitt constraint operator for (e) the non-symmetric version of the operator

constructed previously in this series

G_{40} : Category one of non-symmetric version of the operator

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of complete and rigorous kernel of the Wheeler - DeWitt constraint operator

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

They also construct a

symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of (e) the Wick transform from (e) the Euclidean to the Lorentzian regime they prove (eb) existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.

G_{44} : Category one of Wick transform from (e) the Euclidean to the Lorentzian regime they prove (eb) existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &- (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18\end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}), t) =$ First detritions factor

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ First augmentation factor

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for

1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7)}(G_{39}, t)}$ are seventh detrition coefficients for

category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth

detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a''_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a''_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a''_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{l} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) - (b''_{13})^{(1,1)}(G, t) - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1)}(G, t)} & \boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1)}(G, t)} & \boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{46})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a'_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a'_{21})^{(3)}(T_{21}, t)} & \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} \boxed{+(a'_{22})^{(3)}(T_{21}, t)} & \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \quad 69$$

$\boxed{+(a'_{20})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3)}(T_{21}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)}$ are second augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$ are third augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$ are eight augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{20})^{(3)}(G_{23}, t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} \boxed{(b'_{21})^{(3)}(G_{23}, t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{22})^{(3)}(G_{23}, t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25}, t) & +(a''_{28})^{(5,5)}(T_{29}, t) & +(a''_{32})^{(6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25}, t) & +(a''_{29})^{(5,5)}(T_{29}, t) & +(a''_{33})^{(6,6)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25}, t) & +(a''_{30})^{(5,5)}(T_{29}, t) & +(a''_{34})^{(6,6)}(T_{33}, t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14}, t) & +(a''_{18})^{(2,2,2,2)}(T_{17}, t) & +(a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35}) - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(Y) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

$$(Z) \quad \text{The functions } (a_i'')^{(7)}, (b_i'')^{(7)} \text{ are positive continuous increasing and bounded.}$$

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(AA) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(BB)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(CC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(DD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\tilde{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\tilde{M}_{28})^{(5)}} \left[((\tilde{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\tilde{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\tilde{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\tilde{P}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\tilde{P}_{32})^{(6)}}{(\tilde{M}_{32})^{(6)}} \left(e^{(\tilde{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\tilde{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\tilde{M}_{32})^{(6)}} \left[((\tilde{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\tilde{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\tilde{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(e) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\tilde{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\tilde{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\tilde{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[((\tilde{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\tilde{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\tilde{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\tilde{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\tilde{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\tilde{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[((\tilde{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\tilde{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\tilde{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on
 (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $v^{(2)} = \frac{G_{16}}{G_{17}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case.**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(e) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* \mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* \mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^* \mathbb{G}_j \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^* \mathbb{G}_j \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^* \mathbb{G}_j \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* \mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^* \mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^* \mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^* \mathbb{G}_j \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^* \mathbb{G}_j \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^* \mathbb{G}_j \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIX

Gauge Field Theory Coherent States (GCS)

INTRODUCTION—VARIABLES USED

Optical characterization and band offsets in ZnSe-ZnSxSe1-x strained-layer superlattices Phys. Rev. B 38, 1417 – Published 15 July 1988; Erratum Phys. Rev. B 43, 1830 (1991) Khalid Shahzad, Diego J. Olego, and Chris G. Van de Walle

- (1) Theoretical calculations of the band offsets, based on the model solid approach, were also performed and are found to agree with (=) the experimental observations to within 0.05 eV.
- (2) They indicate that all possible ZnSe-ZnSxSe1-x interfaces will exhibit (eb) a very small

conduction-band offset. DOI: <http://dx.doi.org/10.1103/PhysRevB.38.1417> Received 19 October 1987Published in the issue dated 15 July 1988 © 1988 The American Physical Society

T Thiemann and O Winkler 2001 Class Quantum Gra 18 4997 doi:10.1088/0264-9381/18/23/302 Gauge field theory coherent states (GCS): IV. Infinite tensor product and thermodynamical limit

- (3) In the canonical approach to Lorentzian quantum general relativity in four spacetime dimensions an important step forward has been made by Ashtekar, Isham and Lewandowski some eight years ago through the introduction of a Hilbert space structure, which was later proved to be a faithful representation of (e) the canonical commutation and adjointness relations of (e) the quantum field algebra of (e) diffeomorphism invariant gauge field theories by Ashtekar, Lewandowski, Marolf, Mourão and Thiemann.
- (4) This Hilbert space, together with its generalization due to Baez and Sawin, is appropriate for (e) semi-classical quantum general relativity if (e) the spacetime is spatially compact.
- (5) In the spatially non-compact case, however, an extension of the Hilbert space is needed in order to (e) approximate metrics that are (=) macroscopically nowhere degenerate.
- (6) For this purpose, in this paper authors apply the theory of the infinite tensor product (ITP) of Hilbert Spaces, developed by von Neumann more than sixty years ago, to (e&eb) quantum general relativity.
- (7) The cardinality of the number of tensor product factors can take the value of (e) any possible Cantor aleph, making this mathematical theory well suited to (e) our problem in which a Hilbert space is (=) attached to each edge of an arbitrarily complicated, generally infinite graph.
- (8) The new framework opens access to a new arsenal of techniques, appropriate to describe fascinating physics such as quantum topology change, (e&eb) semi-classical quantum gravity, (e&eb) effective low-energy physics etc from the universal point of view of the ITP. In particular, the study of photons and gravitons propagating on (eb) fluctuating quantum spacetimes should now be in reach.

NOTATION

Module One

T Thiemann 1998 Class Quantum Grav.15 875 doi:10.1088/0264-9381/15/4/012 Quantum spin dynamics (QSD): II. The kernel of the Wheeler - DeWitt constraint operator

symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform from (e) the Euclidean to the Lorentzian regime they prove (eb) existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.

G_{13} : Category one of Euclidean to the Lorentzian regime they prove (eb) existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform from the Euclidean to the Lorentzian regime they prove (eb) existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.

G_{16} : Category one of symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform from the Euclidean to the Lorentzian regime

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform from the Euclidean to the Lorentzian regime they prove existence of self-adjoint extensions and based on these propose (eb) a method of proof of self-adjoint extensions for the regulated Lorentzian operator.

G_{20} : Category one of symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform from the Euclidean to the Lorentzian regime they prove existence of self-adjoint extensions; method of proof of self-adjoint extensions for the regulated Lorentzian operator.

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of method of proof of self-adjoint extensions for the regulated Lorentzian operator.; symmetric, regulated constraint operator for the regulated Euclidean Wheeler - DeWitt operator as well as for the regulated generator of the Wick transform from the Euclidean to the Lorentzian regime they prove existence of self-adjoint extensions

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Theoretical calculations of the band offsets, based on the model solid approach, were also performed and are found to agree with (=) the experimental observations to within 0.05 eV

G_{24} : Category one of Theoretical calculations of the band offsets, based on the model solid approach

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of experimental observations to within 0.05 eV

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

They indicate that

all possible ZnSe-ZnSxSe1-x interfaces will exhibit (eb) a very small conduction-band offset.

DOI: <http://dx.doi.org/10.1103/PhysRevB.38.1417> Received 19 October 1987Published in the issue dated
 15 July 1988 © 1988 The American Physical Society

G_{28} : Category one of all possible ZnSe-ZnSxSe1-x interfaces

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of very small conduction-band offset.

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

In the canonical approach to Lorentzian quantum general relativity in four spacetime dimensions an important step forward has been made by Ashtekar, Isham and Lewandowski some eight years ago through the introduction of a

Hilbert space structure, which was later proved to be a faithful representation of (e) the canonical commutation and adjointness relations of (e) the quantum field algebra of (e) diffeomorphism invariant gauge field theories

by Ashtekar, Lewandowski, Marolf, Mourão and Thiemann

G_{32} : Category one of canonical commutation and adjointness relations of (e) the quantum field algebra of (e) diffeomorphism invariant gauge field theories

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Hilbert space structure

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Hilbert space structure, which was later proved to be a faithful representation of the canonical commutation and adjointness relations of (e) the quantum field algebra of (e) diffeomorphism invariant gauge field theories

G_{36} : Category one of quantum field algebra of (e) diffeomorphism invariant gauge field theories

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Hilbert space structure, which was later proved to be a faithful representation of the canonical commutation and adjointness relations

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Hilbert space structure, which was later proved to be a faithful representation of the canonical commutation and adjointness relations of the quantum field algebra of (e) diffeomorphism invariant gauge field theories

G_{40} : Category one of diffeomorphism invariant gauge field theories

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Hilbert space structure, which was later proved to be a faithful representation of the canonical commutation and adjointness relations of the quantum field algebra

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

This

Hilbert space, together with its generalization due to Baez and Sawin, is appropriate for (e) semi-classical quantum general relativity if (e) the spacetime is spatially compact

G_{44} : Category one of semi-classical quantum general relativity if (e) the spacetime is spatially compact

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Hilbert space, together with its generalization due to Baez and Sawin, is appropriate

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \end{aligned}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor}$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a'_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a'_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a'_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{l} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{l} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{l} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{l} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{l} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients

for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for

category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients

for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition

coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition

coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \begin{bmatrix} \boxed{+(a'_{40})^{(8)}(T_{41}, t)} & \boxed{+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \begin{bmatrix} \boxed{+(a'_{41})^{(8)}(T_{41}, t)} & \boxed{+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} \quad 96$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44}$$

$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t) & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47}, t)$, $-(b''_{45})^{(9)}(G_{47}, t)$, $-(b''_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$$-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(EE) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(FF) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(GG) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(HH)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(II) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(JJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$, Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(f) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right] \quad 181$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)} \right) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a'_{13})^{(1)}$ and $(b'_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}\right), \left((G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})$ it follows $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \quad \text{If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(M_{24})^{(4)}} \left[(\hat{P}_{24})^{(4)} + ((\hat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\hat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(M_{24})^{(4)}} \left[((\hat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)} \right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), ((G_{27})^{(2)}, (T_{27})^{(2)}) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}\right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{-(\hat{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} |(a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a'_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} ((b_{38}'')^{(7)}((G_{39})(t), t)) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

$$\text{roots of the equations } (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t) \right) - (a_{25}'')^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(f) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)})$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)})$$

$$\begin{aligned}
 & + \left((\lambda)^{(1)} \right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)} \right) (\lambda)^{(1)} (q_{15})^{(1)} G_{15} \\
 & + \left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)} \right) \left((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^* \right) \\
 & \left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)} \right) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)} \right) \{ (\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)} \} \\
 & \left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + \left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)} \right) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \right) \\
 & \left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(\left((\lambda)^{(2)} \right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \right) \\
 & \left(\left((\lambda)^{(2)} \right)^2 + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \right) \\
 & + \left(\left((\lambda)^{(2)} \right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \right) (q_{18})^{(2)} G_{18} \\
 & + \left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) \left((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \\
 & \left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \{ (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \} \\
 & \left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(\left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(\left((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(\left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(\left((\lambda)^{(3)} \right)^2 + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \right) \\
 & \left(\left((\lambda)^{(3)} \right)^2 + \left((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVEN

Canonical Commutation And Adjointness Relations Of The Quantum Field Algebra Of Diffeomorphism Invariant Gauge Field Theories By Ashtekar, Lewandowski, Marolf, Mourão And Thiemann.

INTRODUCTION—VARIABLES USED

T Thiemann and O Winkler 2001 Class Quantum Gra 18 4997 doi:10.1088/0264-9381/18/23/302 Gauge field theory coherent states (GCS): IV. Infinite tensor product and thermodynamical limit

- (1) This Hilbert space, together with its generalization due to Baez and Sawin, is appropriate for (e) semi-classical quantum general relativity if (e) the spacetime is spatially compact.
- (2) In the spatially non-compact case, however, an extension of the Hilbert space is needed in order to (e) approximate metrics that are (=) macroscopically nowhere degenerate.
- (3) For this purpose, in this paper authors apply the theory of the infinite tensor product (ITP) of Hilbert Spaces, developed by von Neumann more than sixty years ago, to (e&eb) quantum general relativity.
- (4) The cardinality of the number of tensor product factors can take the value of (e) any possible Cantor aleph, making this mathematical theory well suited to (e) our problem in which a Hilbert space is (=) attached to each edge of an arbitrarily complicated, generally infinite graph.
- (5) The new framework opens access to a new arsenal of techniques, appropriate to describe fascinating physics such as quantum topology change, (e&eb) semi-classical quantum gravity, (e&eb) effective low-energy physics etc from the universal point of view of the ITP. In particular, the study of photons and gravitons propagating on (eb) fluctuating quantum spacetimes should now be in reach.

NOTATION

Module One

This

Hilbert space, together with its generalization due to Baez and Sawin, is appropriate for semi-classical quantum general relativity if (e) the spacetime is spatially compact

G_{13} : Category one of spacetime is spatially compact

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Hilbert space, together with its generalization due to Baez and Sawin, is appropriate for semi-classical quantum general relativity

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

In the

spatially non-compact case calls for an extension of the Hilbert space is needed in order to (e) approximate metrics that are (=) macroscopically nowhere degenerate

G_{16} : Category one of approximate metrics that are (=) macroscopically nowhere degenerate

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of spatially non-compact case calls for an extension of the Hilbert space is needed

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

spatially non-compact case calls for an extension of the Hilbert space is needed in order to approximate metrics that are (=) macroscopically nowhere degenerate

G_{20} : Category one of spatially non-compact case calls for an extension of the Hilbert space is needed in order to approximate metrics

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of macroscopically nowhere degenerate

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

For this purpose, in this paper authors apply the

theory of the infinite tensor product (ITP) of Hilbert Spaces, developed by von Neumann more than sixty years ago, to (e&eb) quantum general relativity

G_{24} : Category one of theory of the infinite tensor product (ITP) of Hilbert Spaces; quantum general relativity

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of quantum general relativity; theory of the infinite tensor product (ITP) of Hilbert Spaces

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The

cardinality of the number of tensor product factors can take the value of (e) any possible Cantor aleph, making this mathematical theory well suited to (e) our problem in which a Hilbert space is (=) attached to each edge of an arbitrarily complicated, generally infinite graph

G_{28} : Category one of any possible Cantor aleph, making mathematical theory well suited to (e) extant problem in which a Hilbert space is (=) attached to each edge of an arbitrarily complicated, generally infinite graph

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of cardinality of the number of tensor product factors can take the value

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

cardinality of the number of tensor product factors can take the value of any possible Cantor aleph, making this mathematical theory well suited to (e) extant problem in which a Hilbert space is (=) attached to each edge of an arbitrarily complicated, generally infinite graph

G_{32} : Category one of extant problem in which a Hilbert space is (=) attached to each edge of an arbitrarily complicated, generally infinite graph

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of cardinality of the number of tensor product factors can take the value of any possible Cantor aleph, making this mathematical theory well suited

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

cardinality of the number of tensor product factors can take the value of any possible Cantor aleph, making this mathematical theory well suited to extant problem in which a Hilbert space is attached to each edge of an arbitrarily complicated, generally infinite graph

G_{36} : Category one of cardinality of the number of tensor product factors can take the value of any possible Cantor aleph, making this mathematical theory well suited to extant problem in which a Hilbert space; each edge of an arbitrarily complicated, generally infinite graph

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of each edge of an arbitrarily complicated, generally infinite graph; cardinality of the number of tensor product factors can take the value of any possible Cantor aleph, making this mathematical theory well suited to extant problem in which a Hilbert space

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

The new framework opens access to a new arsenal of techniques, appropriate to describe fascinating physics such as quantum topology change, semi-classical quantum gravity, effective low-energy physics etc from the universal point of view of the ITP.

In particular, the study of photons and gravitons propagating on fluctuating quantum spacetimes should now be in reach.

G_{40} : Category one of quantum topology change; semi-classical quantum gravity

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of semi-classical quantum gravity ;quantum topology change

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

In particular, the study of

photons and gravitons propagating on fluctuating quantum spacetimes
 should now be in reach.

G_{44} : Category one of photons and gravitons; fluctuating quantum spacetimes

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of fluctuating quantum spacetimes; photons and gravitons

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \end{aligned}$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$

$-(b''_{44})^{(9)}(G_{47}, t) = \text{First detrition factor}$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} \boxed{+(a''_{13})^{(1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} \boxed{+(a''_{14})^{(1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} \boxed{+(a''_{15})^{(1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{15} \quad 57$$

Where $\boxed{(a'_{13})^{(1)}(T_{14}, t)}$, $\boxed{(a'_{14})^{(1)}(T_{14}, t)}$, $\boxed{(a'_{15})^{(1)}(T_{14}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)} \boxed{-(b''_{13})^{(1)}(G, t)} \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{c} (b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] & -(b''_{18})^{(2,2)}(G_{19}, t) & -(b''_{22})^{(3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7)}(G_{39}, t) & -(b''_{42})^{(8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $[-(b'_{13})^{(1)}(G, t)]$, $[-(b'_{14})^{(1)}(G, t)]$, $[-(b'_{15})^{(1)}(G, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b'_{16})^{(2,2)}(G_{19}, t)]$, $[-(b'_{17})^{(2,2)}(G_{19}, t)]$, $[-(b'_{18})^{(2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b'_{20})^{(3,3)}(G_{23}, t)]$, $[-(b'_{21})^{(3,3)}(G_{23}, t)]$, $[-(b'_{22})^{(3,3)}(G_{23}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b'_{24})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b'_{25})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b'_{26})^{(4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b'_{28})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b'_{29})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b'_{30})^{(5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b'_{32})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b'_{33})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b'_{34})^{(6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b'_{37})^{(7,7)}(G_{39}, t)]$, $[-(b'_{36})^{(7,7)}(G_{39}, t)]$, $[-(b'_{38})^{(7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b'_{40})^{(8,8)}(G_{43}, t)]$, $[-(b'_{41})^{(8,8)}(G_{43}, t)]$, $[-(b'_{42})^{(8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17}, t)] & +(a''_{13})^{(1,1)}(T_{14}, t) & +(a''_{20})^{(3,3,3)}(T_{21}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{36})^{(7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)}[+(a''_{17})^{(2)}(T_{17}, t)] & +(a''_{14})^{(1,1)}(T_{14}, t) & +(a''_{21})^{(3,3,3)}(T_{21}, t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{37})^{(7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}[+(a''_{18})^{(2)}(T_{17}, t)] & +(a''_{15})^{(1,1)}(T_{14}, t) & +(a''_{22})^{(3,3,3)}(T_{21}, t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{38})^{(7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $[(a'_{16})^{(2)}(T_{17}, t)]$, $[(a'_{17})^{(2)}(T_{17}, t)]$, $[(a'_{18})^{(2)}(T_{17}, t)]$ are first augmentation coefficients for category 1, 2 and 3

$[(a'_{13})^{(1,1)}(T_{14}, t)]$, $[(a'_{14})^{(1,1)}(T_{14}, t)]$, $[(a'_{15})^{(1,1)}(T_{14}, t)]$ are second augmentation coefficient for category 1, 2 and 3

$[(a'_{20})^{(3,3,3)}(T_{21}, t)]$, $[(a'_{21})^{(3,3,3)}(T_{21}, t)]$, $[(a'_{22})^{(3,3,3)}(T_{21}, t)]$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)}(G_{19}, t) & -(b'_{13})^{(1,1)}(G, t) & -(b'_{20})^{(3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b'_{36})^{(7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)}(G_{19}, t) & -(b'_{14})^{(1,1)}(G, t) & -(b'_{21})^{(3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b'_{37})^{(7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)}(G_{19}, t) & -(b'_{15})^{(1,1)}(G, t) & -(b'_{22})^{(3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a''_{16})^{(2,2,2)}(T_{17}, t) + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a''_{17})^{(2,2,2)}(T_{17}, t) + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a''_{18})^{(2,2,2)}(T_{17}, t) + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{l} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{16})^{(2,2,2)}(G_{19}, t) - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{l} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) - (b''_{17})^{(2,2,2)}(G_{19}, t) - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$[-(b'_{20})^{(3)}(G_{23}, t)], [-(b'_{21})^{(3)}(G_{23}, t)], [-(b'_{22})^{(3)}(G_{23}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b'_{16})^{(2,2,2)}(G_{19}, t)], [-(b'_{17})^{(2,2,2)}(G_{19}, t)], [-(b'_{18})^{(2,2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b'_{13})^{(1,1,1)}(G, t)], [-(b'_{14})^{(1,1,1)}(G, t)], [-(b'_{15})^{(1,1,1)}(G, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b'_{36})^{(7,7,7,7)}(G_{39}, t)], [-(b'_{37})^{(7,7,7,7)}(G_{39}, t)], [-(b'_{38})^{(7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b'_{40})^{(8,8,8,8)}(G_{43}, t)], [-(b'_{41})^{(8,8,8,8)}(G_{43}, t)], [-(b'_{42})^{(8,8,8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b'_{46})^{(9,9,9)}(G_{47}, t)], [-(b'_{45})^{(9,9,9)}(G_{47}, t)], [-(b'_{44})^{(9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)}[+(a''_{24})^{(4)}(T_{25}, t)] & +(a'_{28})^{(5,5)}(T_{29}, t) & +(a'_{32})^{(6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)}[+(a''_{25})^{(4)}(T_{25}, t)] & +(a'_{29})^{(5,5)}(T_{29}, t) & +(a'_{33})^{(6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)}[+(a''_{26})^{(4)}(T_{25}, t)] & +(a'_{30})^{(5,5)}(T_{29}, t) & +(a'_{34})^{(6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$[(a'_{24})^{(4)}(T_{25}, t)], [(a'_{25})^{(4)}(T_{25}, t)], [(a'_{26})^{(4)}(T_{25}, t)]$ are first augmentation coefficients category 1, 2 3

$[(a'_{28})^{(5,5)}(T_{29}, t)], [(a'_{29})^{(5,5)}(T_{29}, t)], [(a'_{30})^{(5,5)}(T_{29}, t)]$ are second augmentation coefficient for category 1, 2 and 3

$[(a'_{32})^{(6,6)}(T_{33}, t)], [(a'_{33})^{(6,6)}(T_{33}, t)], [(a'_{34})^{(6,6)}(T_{33}, t)]$ are third augmentation coefficient for category 1, 2 and 3

$[(a'_{13})^{(1,1,1,1)}(T_{14}, t)], [(a'_{14})^{(1,1,1,1)}(T_{14}, t)], [(a'_{15})^{(1,1,1,1)}(T_{14}, t)]$

are fourth augmentation coefficients for category 1, 2 and 3

$$+(a''_{16})^{(2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2)}(T_{17}, t)$$

are fifth augmentation coefficients for category 1, 2 and 3

$$+(a''_{20})^{(3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3)}(T_{21}, t)$$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a''_{25})^{(4,4)}(T_{25}, t) & + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1, 2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1, 2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1, 2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1, 2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \text{ sixth augmentation}$$

coefficients

$$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \text{ ninth augmentation}$$

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41}, t)} & \boxed{+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41}, t)} & \boxed{+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}$, $(b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}$, $(\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}$, $(\hat{B}_{13})^{(1)}$, $(p_i)^{(1)}$, $(r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}$, $(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}$, $(\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(KK) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(LL) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(MM) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(NN)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(OO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(PP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(g) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right] \quad 181$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$|\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} +$$

$$\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} \leq$$

$$\frac{1}{(\bar{M}_{20})^{(3)}} \left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)} \right) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{-(\hat{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)}t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as 249
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\bar{G}_{40}^{(1)} - \bar{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &((a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
 $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
 $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
 $(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

409

410

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines $(\nu_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{[-(a_{41})^{(8)}((\nu_1)^{(8)} - (\nu_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((\nu_1)^{(8)} - (\nu_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get

422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (\bar{C})^{(8)} (v_2)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)} (T_{45}, t) \right) - (a_{45}'')^{(9)} (T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(g) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $\mathcal{C}^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $\mathcal{C}^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ \left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \right. \\
 &\left. \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 &+ \left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \right. \\
 &\left. \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left((\lambda)^{(9)} \right)^2 + \left((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 &+ \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 &\left(\left((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)} \right) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION EIGHT

Renormalization In The Ads/CFT Correspondence

INTRODUCTION—VARIABLES USED

Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin

- (1) **Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin** develop a systematic method for renormalizing the AdS/CFT prescription for (e) computing correlation functions.
- (2) This involves regularizing the bulk on-shell supergravity action in (eb) a covariant way, computing all divergences, adding counterterms to cancel (e&eb) them and then removing the regulator.
- (3) They explicitly work out the case of pure gravity up to six dimensions and (e&eb) of gravity coupled to scalars.
- (4) The method can also be viewed and visualized as providing a holographic reconstruction of (e) the bulk spacetime metric and of bulk fields on this spacetime, out of (e) conformal field theory data.
- (5) Knowing which sources are turned on is sufficient in order to obtain (eb) an asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order so that all infrared divergences of (e) the on-shell action are obtained.
- (6) To continue the holographic reconstruction of the bulk fields one needs (e) new CFT data: the **expectation value of the dual operator**
- (7) In particular, in order to obtain the bulk metric one needs (e) to know the expectation value of stress-energy tensor of the boundary theory.
- (8) **Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin** provide completely explicit formulae for (e) the holographic stress-energy tensors up to six dimensions.
- (9) They show that both the gravitational and matter conformal anomalies of (e) the boundary theory are correctly reproduced.

NOTATION

Module One

Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin develop a systematic method

renormalizing the AdS/CFT prescription for (e) computing correlation functions

G_{13} : Category one of computing correlation functions

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of renormalizing the AdS/CFT prescription

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

This involves

regularizing the bulk on-shell supergravity action in (eb) a covariant way, computing all divergences, adding counterterms to cancel (e&eb) them and then removing the regulator

G_{16} : Category one of regularizing the bulk on-shell supergravity action

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of covariant way, computing all divergences, adding counterterms to cancel (e&eb) them and then removing the regulator

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

regularizing the bulk on-shell supergravity action in a covariant way, computing all divergences, adding counterterms to cancel (e&eb) them and then removing the regulator

G_{20} : Category one of regularizing the bulk on-shell supergravity action in a covariant way, computing all divergences, adding counterterms; them (concomitant terms for divergences) and then removing the regulator

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of them (concomitant terms for divergences) and then removing the regulator; regularizing the bulk on-shell supergravity action in a covariant way, computing all divergences, adding counterterms

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

They explicitly work out the

case of pure gravity up to six dimensions and (e&eb) of gravity coupled to scalars

G_{24} : Category one of case of pure gravity up to six dimensions; gravity coupled to scalars

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of gravity coupled to scalars; case of pure gravity up to six dimensions

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The method can also be viewed and visualized as providing a

holographic reconstruction of (e) the bulk spacetime metric and of bulk fields on this spacetime, out of (e) conformal field theory data

G_{28} : Category one of bulk spacetime metric and of bulk fields on this spacetime, out of (e) conformal field theory data

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of holographic reconstruction

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

holographic reconstruction of the bulk spacetime metric and of bulk fields on this spacetime, out of (e) conformal field theory data

G_{32} : Category one of conformal field theory data

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of holographic reconstruction of the bulk spacetime metric and of bulk fields on this spacetime

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Knowing which sources are turned on is sufficient in order to obtain (eb) an asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order so that all infrared divergences of (e) the on-shell action are obtained

G_{36} : Category one of sources turned on information

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order so that all infrared divergences of (e) the on-shell action are obtained

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

sources turned on information provides asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order so that all infrared divergences of the on-shell action are obtained

G_{40} : Category one of sources turned on information provides asymptotic expansion of the bulk metric and of bulk fields near the boundary to high enough order

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of all infrared divergences of the on-shell action are obtained

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

To continue the holographic reconstruction of the bulk fields one needs (e) new CFT data: the **expectation value of the dual operator**

G_{44} : Category one of CFT data: the **expectation value of the dual operator**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of continuation of the holographic reconstruction of the bulk fields

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$

$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$

$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{|c|c|c|c|} \hline (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14}, t) & +(a''_{16})^{(2,2)}(T_{17}, t) & +(a''_{20})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{|c|c|c|c|} \hline (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14}, t) & +(a''_{17})^{(2,2)}(T_{17}, t) & +(a''_{21})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a'_{18})^{(2,2)}(T_{17}, t) + (a'_{22})^{(3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7)}(T_{37}, t) + (a'_{42})^{(8,8)}(T_{41}, t) + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for

category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for

category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient

for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient

for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for

1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{c} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b'_{13})^{(1)}(G, t)$, $-(b'_{14})^{(1)}(G, t)$, $-(b'_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1,

2 and 3

$-(b'_{16})^{(2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for

category 1, 2 and 3

$-(b'_{20})^{(3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for

category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients

for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for

category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} \boxed{-(b''_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b''_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category

1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition

coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients

for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients

for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for

category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for

category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category

1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a''_{21})^{(3)}(T_{21}, t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{22}$$

$+(a'_{20})^{(3)}(T_{21}, t)$, $+(a'_{21})^{(3)}(T_{21}, t)$, $+(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & - (b''_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{36})^{(7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & - (b''_{21})^{(3)}(G_{23}, t) & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{37})^{(7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{21}$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} & - (b''_{22})^{(3)}(G_{23}, t) & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{38})^{(7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{22}$$

$-(b'_{20})^{(3)}(G_{23}, t)$, $-(b'_{21})^{(3)}(G_{23}, t)$, $-(b'_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7)}(G_{39}, t) \text{ are seventh detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8)}(G_{43}, t) \text{ are eight detrition coefficients for}$$

category 1, 2 and 3

$$-(b''_{46})^{(9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for}$$

category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{c} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$$(a''_{24})^{(4)}(T_{25}, t), (a''_{25})^{(4)}(T_{25}, t), (a''_{26})^{(4)}(T_{25}, t) \text{ are first augmentation coefficients}$$

category 1, 2 3

$$+(a''_{28})^{(5,5)}(T_{29}, t), +(a''_{29})^{(5,5)}(T_{29}, t), +(a''_{30})^{(5,5)}(T_{29}, t) \text{ are second augmentation}$$

coefficient for category 1, 2 and 3

$$+(a''_{32})^{(6,6)}(T_{33}, t), +(a''_{33})^{(6,6)}(T_{33}, t), +(a''_{34})^{(6,6)}(T_{33}, t) \text{ are third augmentation}$$

coefficient for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1)}(T_{14}, t)$$

are fourth augmentation coefficients for category 1, 2 and 3

$$+(a''_{16})^{(2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2)}(T_{17}, t)$$

are fifth augmentation coefficients for category 1, 2 and 3

$$+(a''_{20})^{(3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3)}(T_{21}, t)$$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for}$$

category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)}(G_{27}, t) & -(b'_{28})^{(5,5)}(G_{31}, t) & -(b'_{32})^{(6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1)}(G, t) & -(b'_{16})^{(2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b'_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)}(G_{27}, t) & -(b'_{29})^{(5,5)}(G_{31}, t) & -(b'_{33})^{(6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1)}(G, t) & -(b'_{17})^{(2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b'_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)}(G_{27}, t) & -(b'_{30})^{(5,5)}(G_{31}, t) & -(b'_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$ are first augmentation coefficients for category 1, 2 and 3

And $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$ are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$ are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$ are sixth augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)}$ are eighth augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} \boxed{(b'_{28})^{(5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} \boxed{(b'_{29})^{(5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} \boxed{(b'_{30})^{(5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$ are sixth detrition coefficients

for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{32} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} & + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35}, t) & -(b''_{28})^{(5,5,5)}(G_{31}, t) & -(b''_{24})^{(4,4,4)}(G_{27}, t) \\ -(b''_{13})^{(1,1,1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35}, t) & -(b''_{29})^{(5,5,5)}(G_{31}, t) & -(b''_{25})^{(4,4,4)}(G_{27}, t) \\ -(b''_{14})^{(1,1,1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b''_{30})^{(5,5,5)}(G_{31}, t) & -(b''_{26})^{(4,4,4)}(G_{27}, t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & - (b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & - (b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for

category 1, 2 and 3

$$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} \text{ are fifth detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \text{ are sixth detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)}$$

are seventh detrition coefficients for category 1, 2 and 3

$$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} \text{ are eighth detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \text{ are ninth detrition}$$

coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} \boxed{(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} \boxed{(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt}$$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} \boxed{(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8)}(T_{41}, t)}$ are first augmentation coefficients for

category 1, 2 and 3

$$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)} \text{ are second}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) \text{ are seventh}$$

augmentation coefficient for 1,2,3

$$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t), -(b''_{37})^{(7)}(G_{39}, t), -(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2 and 3

$$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t), -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) \text{ are sixth detrition coefficients}$$

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition

coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition

coefficients for category 1, 2 and 3

$$\begin{aligned} & \frac{dG_{44}}{dt} \\ &= (a_{44})^{(9)}G_{45} \\ & - \left[\begin{array}{l} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} & \frac{dG_{45}}{dt} \\ &= (a_{45})^{(9)}G_{44} \\ & - \left[\begin{array}{l} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} & \frac{dG_{46}}{dt} \\ &= (a_{46})^{(9)}G_{45} \\ & - \left[\begin{array}{l} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\begin{aligned} \frac{dT_{44}}{dt} &= \\ (b_{44})^{(9)}T_{45} - & \left[\begin{array}{ccc} (b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t) & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\ \frac{dT_{45}}{dt} &= \\ (b_{45})^{(9)}T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\ \frac{dT_{46}}{dt} &= \\ (b_{46})^{(9)}T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15} \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1$$

111

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22$$

112

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)}$$

113

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition:

114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$:

115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$\begin{aligned} (a''_i)^{(4)}(T_{25}, t) &\leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)} \\ (b''_i)^{(4)}((G_{27}), t) &\leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)} \\ \lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) &= (p_i)^{(4)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) &= (r_i)^{(4)} \end{aligned} \quad 118$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$\begin{aligned} |(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| &\leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)}t} \\ |(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| &< (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)}t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\begin{aligned} \frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] &< 1 \\ \frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] &< 1 \end{aligned}$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$\begin{aligned} (a''_i)^{(5)}(T_{29}, t) &\leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)} \\ (b''_i)^{(5)}((G_{31}), t) &\leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)} \\ \lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) &= (p_i)^{(5)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) &= (r_i)^{(5)} \end{aligned} \quad 123$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$\begin{aligned} |(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| &\leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t} \\ |(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| &< (\hat{k}_{28})^{(5)} \|(G_{31}) - (G_{31})'\| e^{-(\hat{M}_{28})^{(5)} t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(QQ) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(RR) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(SS) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(TT)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ are positive constants and $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T'_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(UU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(VV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ the constants

$(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a'')^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b'')^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a'')^{(8)}, (b'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a'')^{(8)}(T'_{41}, t) - (a'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b'')^{(8)}((G_{43})', t) - (b'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a'')^{(8)}(T'_{41}, t)$ and $(a'')^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$
 $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42,$

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0} \right)} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(h) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)}(\bar{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e \left[- \int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)} \right] \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\bar{M}_{13})^{(1)})_1, ((\bar{M}_{13})^{(1)})_2$ and $((\bar{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < (\bar{M}_{13})^{(1)} \text{ it follows } \frac{dG_{14}}{dt} \leq ((\bar{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)} (G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)} (G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[(\widehat{P}_{16})^{(2)} + ((\widehat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[((\widehat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{16})^{(2)} \right] \leq (\widehat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$

197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\bar{A}_{16})^{(2)} + (\bar{P}_{16})^{(2)} (\bar{k}_{16})^{(2)}) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}); (G_{19})^{(2)}, (T_{19})^{(2)} \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

198

Remark 6: The fact that we supposed $(a'_{16})^{(2)}$ and $(b'_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\bar{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ .

199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(2)}$ and $(b'_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

200

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{ (a'_i)^{(2)} - (a''_i)^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{16})^{(2)})_1, ((\bar{M}_{16})^{(2)})_2$ and $((\bar{M}_{16})^{(2)})_3$:

201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\bar{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq ((\bar{M}_{16})^{(2)})_1 - (a'_{17})^{(2)} G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(2)} ((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b'_i)^{(2)} ((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b''_{18})^{(2)} ((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[(\widehat{P}_{20})^{(3)} + ((\widehat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[((\widehat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{20})^{(3)} \right] \leq (\widehat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \}$$

Indeed if we denote

212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}) d\left((G_{23})^{(1)}, (T_{23})^{(1)}; (G_{23})^{(2)}, (T_{23})^{(2)}\right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

216

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \} ds_{(20)}]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(3)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{20})^{(3)})_1, ((\bar{M}_{20})^{(3)})_2$ and $((\bar{M}_{20})^{(3)})_3 :$

217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b'_i)^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b''_{22})^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[(\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying 224
Equations into itself

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \}$$

Indeed if we denote

$$\textbf{Definition of } (\widetilde{G_{27}}, \widetilde{T_{27}}) : (\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_{24}^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)} \right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right); (G_{27})^{(2)}, (T_{27})^{(2)}\right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{24})^{(4)})_1, ((\bar{M}_{24})^{(4)})_2$ and $((\bar{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b'_i)^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[(\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric

$$d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) = \\ \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_{28}^{(2)}| \leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ \int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{-(\bar{M}_{28})^{(5)}s_{(28)}} + \\ (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} + \\ G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{1}{(\bar{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\bar{A}_{28})^{(5)} + (\bar{P}_{28})^{(5)} (\bar{k}_{28})^{(5)}) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$ and $(\bar{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \} ds_{(28)}} \geq 0 \\ T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{28})^{(5)})_1, ((\bar{M}_{28})^{(5)})_2$ and $((\bar{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)}G_{29}$ and by integrating

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) = \\ \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$|\tilde{G}_{32}^{(1)} - \tilde{G}_{32}^{(2)}| \leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ \int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{-(\bar{M}_{32})^{(6)}s_{(32)}} + \\ (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} + \\ G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{1}{(\bar{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\bar{A}_{32})^{(6)} + (\bar{P}_{32})^{(6)} (\bar{k}_{32})^{(6)}) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right); (G_{35})^{(2)}, (T_{35})^{(2)}\right) \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}$ and $(\bar{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0 \\ T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{32})^{(6)})_1, ((\bar{M}_{32})^{(6)})_2$ and $((\bar{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying

Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric

258

$$d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right), \left((G_{39})^{(2)}, (T_{39})^{(2)}\right)\right) = \sup_{t \in \mathbb{R}_+} \left\{ \max_i |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_i |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}, \widetilde{T_{39}}) : (\widetilde{G_{39}}, \widetilde{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} |(a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)} \right) d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right); (G_{39})^{(2)}, (T_{39})^{(2)}\right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$:

262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$G_{36} < ((\widehat{M}_{36})^{(7)})$ it follows $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266

$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

268

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\hat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{40})^{(8)} \right] \leq (\hat{Q}_{40})^{(8)}$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d \left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widehat{G_{43}}, \widehat{T_{43}})$: $(\widehat{G_{43}}, \widehat{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} \} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} &((a_{40})^{(8)} + (a'_{40})^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}) d \left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)}) \right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1, ((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$$G_{40} < (\widehat{M}_{40})^{(8)} \text{ it follows } \frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)}G_{41} \text{ and by integrating}$$

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b_{42}'')^{(8)}((G_{43})(t), t(t), t) = (b_{42}')^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[\left((\hat{Q}_{44})^{(9)} + T_j^0 \right) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \}$$

Indeed if we denote

Definition of $(\bar{G}_{47}, \bar{T}_{47}) : (\bar{G}_{47}, \bar{T}_{47}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a_{44}')^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a_{44}'')^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a_{44}'')^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a_{44}'')^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a_{44}')^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a_{44}'')^{(9)}$ and $(b_{44}'')^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a_i')^{(9)} - (a_i'')^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \} ds_{(44)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1, ((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$G_{44} < ((\widehat{M}_{44})^{(9)})_1$ it follows $\frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b'_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \leq$$

286

$$\frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$$

287

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

288

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq$$

289

$$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^0e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying
 $-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$ 294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$ 305

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$ 306

and $(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(\nu_1)^{(3)} > 0, (\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(\nu^{(3)})^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{\nu}_1)^{(3)} > 0, (\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}(\nu^{(3)})^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}, \text{ if } (\nu_0)^{(3)} < (\nu_1)^{(3)}$$

$$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}, \text{ if } (\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)},$$

$$\text{and } \boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}, \text{ if } (\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(s_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(s_1)^{(3)}t}$$

323

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right.$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(\nu_1)^{(5)} > 0, (\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{\nu}_1)^{(5)}, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{\nu}_1)^{(5)} > 0, (\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$
Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and $(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \right.$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}}$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{33}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:-

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) - (b''_{41})^{(8)}(G_{43}, t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations
 $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
 and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((s_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(s_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t} \right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$
Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(s_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_2)^{(9)}} G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(s_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((s_1)^{(9)} - (p_{44})^{(9)} - (s_2)^{(9)})} \left[e^{((s_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(s_2)^{(9)}t} \right] + G_{46}^0 e^{-(s_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((s_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(s_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-

$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad (\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{c})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}) t]}}{1 + (\bar{c})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}) t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)} G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)} (T_{17}, t) \right) - (a_{17}'')^{(2)} (T_{17}, t) v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

Definition of $v^{(2)}$:-

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

388

It follows

389

$$- \left((a_{17})^{(2)} (v^{(2)})^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)} (v^{(2)})^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof: From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- 399

$$v^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (C)^{(3)}(\bar{\nu}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)})t]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain

404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$- \left((a_{25})^{(4)}(\nu^{(4)})^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} \right) \leq \frac{d\nu^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(\nu^{(4)})^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{[-(a_{25})^{(4)}((\nu_1)^{(4)} - (\nu_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((\nu_1)^{(4)} - (\nu_0)^{(4)})t]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then

$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$-\left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)}\right) \leq \frac{dv^{(5)}}{dt} \leq -\left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \left(v_0\right)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad (C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad (\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \left(v_0\right)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of $(u_0)^{(5)}$.**

Proof : From global equations we obtain

412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:-

$$\nu^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left((a_{33})^{(6)}(\nu^{(6)})^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} \right) \leq \frac{d\nu^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(\nu^{(6)})^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get

413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case,

414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((\nu_1)^{(6)} - (\nu_2)^{(6)})t]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)})t]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case**.

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a_{36}')^{(7)} - (a_{37}')^{(7)} + (a_{36}'')^{(7)} (T_{37}, t) \right) - (a_{37}'')^{(7)} (T_{37}, t) v^{(7)} - (a_{37})^{(7)} v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$- \left((a_{37})^{(7)} (v^{(7)})^2 + (\sigma_2)^{(7)} v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)} (v^{(7)})^2 + (\sigma_1)^{(7)} v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_0)^{(7)}) t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_0)^{(7)}) t]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)} , \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)} , \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition

$(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof: From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

For $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner, we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case .**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)} (a_{14}')^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)} (a_{14}')^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} + (a_{13})^{(1)} (p_{13})^{(1)} + (a_{14}')^{(1)} (p_{14})^{(1)} + (p_{13})^{(1)} (p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{28})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{32})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{36})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{40})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied, then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution, which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution, which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution, which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(h) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that

there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42})$, G_{40}, G_{42} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and

512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

G_{25}^* given by $\varphi(G_{27}) = 0, T_{25}^*$ given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]} \\ T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]} \\ T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]} \\ T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]} \\ T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]} \quad 523$$

Finally we obtain the unique solution of 89 to 99 523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19})^* = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a_{16}')^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a_{17}')^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a_{18}')^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b_{16}')^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b_{17}')^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b_{18}')^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23})^* = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a_{20}')^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a_{21}')^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a_{22}')^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b_{20}')^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b_{21}')^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22}(s_{(21)(j)}T_{21}^*G_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b_{22}')^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(22)(j)}T_{22}^*G_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

548

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

556

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)} T_{29}^* G_j) \quad 561$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)} T_{30}^* G_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 564

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}} (T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j} ((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^* T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^* T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^* T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)} T_{32}^* G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)} T_{33}^* G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)} T_{34}^* G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}} (T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial (b_i'')^{(7)}}{\partial G_j} ((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*\mathbb{G}_j \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*\mathbb{G}_j \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*\mathbb{G}_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}(G_{43}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*\mathbb{G}_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}(s_{(41)(j)})T_{41}^*\mathbb{G}_j \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}(s_{(42)(j)})T_{42}^*\mathbb{G}_j \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a_{44}')^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad 586 \text{ B}$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a_{45}')^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad 586 \text{ C}$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a_{46}')^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad 586 \text{ D}$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b_{44}')^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)}T_{44}^*\mathbb{G}_j) \quad 586 \text{ E}$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b_{45}')^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)}T_{45}^*\mathbb{G}_j) \quad 586 \text{ F}$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b_{46}')^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)}T_{46}^*\mathbb{G}_j) \quad 586 \text{ G}$$

The characteristic equation of this system is

587

$$\begin{aligned} & ((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}) \\ & \left[((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right] \\ & \left(((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^* \right) \\ & + \left(((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^* \right) \\ & \left(((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^* \right) \\ & \left(((\lambda)^{(1)})^2 + ((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & \left(((\lambda)^{(1)})^2 + ((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & + \left(((\lambda)^{(1)})^2 + ((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\ & + ((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\ & \left. \left(((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \right\} = 0 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \right) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & +
 \end{aligned}$$

$$\begin{aligned}
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & + \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ (\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)} \} \\
 & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 & \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 & \left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 & + \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & \left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & + \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION NINE

Ads/CFT Correspondence

INTRODUCTION—VARIABLES USED

Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin

- (1) In particular, in order to obtain the bulk metric one needs (e) to know the expectation value of stress-energy tensor of the boundary theory.
- (2) **Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin** provide completely explicit formulae for (e) the holographic stress-energy tensors up to six dimensions.
- (3) They show that both the gravitational and matter conformal anomalies of (e) the boundary theory are correctly reproduced.
- (4) They also obtain the conformal transformation properties of (e) the boundary stress-energy tensors.
Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin.
- (5) The theory of holographic space-time (HST) generalizes both string theory and quantum field theory. Theory of holographic space-time (HST) provides (eb) a geometric rationale for supersymmetry (SUSY) and formalism in (eb) which super-Poincare invariance follows from (e) Poincare invariance.
- (6) HST unifies particles and (e&eb) black holes, realizing both as (=) excitations of non-commutative geometrical variables on (eb) a holographic screen.
- (7) Compact extra dimensions are interpreted as (=) finite dimensional unitary representations of (e) super-algebras, and have (e) no moduli.
- (8) Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of (e) super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity.
- (9) Finite radius de Sitter (dS) spaces have (e) no moduli, and break (e) SUSY with a gravitino mass scaling like $\Lambda^{1/4}$

NOTATION

Module One

In particular, in order to obtain the bulk metric one needs (e) to know the expectation value of stress-energy tensor of the boundary theory

G_{13} : Category one of expectation value of stress-energy tensor of the boundary theory

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of bulk metric

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin provide completely explicit formulae for (e) the

holographic stress-energy tensors up to six dimensions

G_{16} : Category one of six dimensions; holographic stress-energy tensors

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of holographic stress-energy tensors; six dimensions

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

They show that both the gravitational and matter conformal anomalies of (e) the boundary theory are correctly reproduced
 G_{20} : Category one of boundary theory are correctly reproduced

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of gravitational and matter conformal anomalies

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

They also obtain the conformal transformation properties of (e) the boundary stress-energy tensors.

Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin

G_{24} : Category one of boundary stress-energy tensors

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of conformal transformation properties

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The
 theory of holographic space-time (HST) generalizes both string theory and quantum field theory.

Theory of holographic space-time (HST) provides (eb) a geometric rationale for supersymmetry (SUSY) and formalism in (eb) which super-Poincare invariance follows from (e) Poincare invariance
 G_{28} : Category one of theory of holographic space-time (HST); string theory and quantum field theory.

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of string theory and quantum field theory. ; theory of holographic space-time (HST)

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Theory of holographic space-time (HST) provides (eb) a geometric rationale for supersymmetry (SUSY) and formalism in (eb) which super-Poincare invariance follows from (e) Poincare invariance

G_{32} : Category one of Theory of holographic space-time (HST)

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of geometric rationale for supersymmetry (SUSY) and formalism in (eb) which super-Poincare invariance follows from (e) Poincare invariance

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Theory of holographic space-time (HST) provides a geometric rationale for supersymmetry (SUSY) and formalism in (eb) which super-Poincare invariance follows from (e) Poincare invariance

G_{36} : Category one of Theory of holographic space-time (HST) provides a geometric rationale for supersymmetry (SUSY) and formalism

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of super-Poincare invariance follows from (e) Poincare invariance

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Theory of holographic space-time (HST) provides a geometric rationale for supersymmetry (SUSY) and formalism in which super-Poincare invariance follows from (e) Poincare invariance

G_{40} : Category one of Poincare invariance

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Theory of holographic space-time (HST) provides a geometric rationale for supersymmetry (SUSY) and formalism in which super-Poincare invariance follows

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

HST unifies particles and (e&eb) black holes, realizing both as (=) excitations of non-commutative geometrical variables on (eb) a holographic screen

G_{44} : Category one of HST unifies particles; black holes, realizing both as (=) excitations of non-commutative geometrical variables on (eb) a holographic screen

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of black holes, realizing both as (=) excitations of non-commutative geometrical variables on (eb) a holographic screen; HST unifies particles

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &- (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} & 18 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}, t))]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}, t))]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}, t))]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}, t)) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}, t))]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}, t))]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}, t))]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}, t)) =$ First detritions factor

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}, t))]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}, t))]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}, t))]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ First augmentation factor

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 56$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 57$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for

1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{13})^{(1)}(G, t)} \quad \boxed{-(b'_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b'_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{14})^{(1)}(G, t)} \quad \boxed{-(b'_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b'_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{15})^{(1)}(G, t)} \quad \boxed{-(b'_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b'_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for

category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b'_{37})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b'_{36})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7)}(G_{39}, t)}$ are seventh detrition coefficients for

category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth

detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a''_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a''_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a''_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{l} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) - (b''_{13})^{(1,1)}(G, t) - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1)}(G, t)} & \boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1)}(G, t)} & \boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{46})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a'_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a'_{21})^{(3)}(T_{21}, t)} & \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} \boxed{+(a'_{22})^{(3)}(T_{21}, t)} & \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \quad 69$$

$\boxed{+(a'_{20})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3)}(T_{21}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)}$ are second augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$ are third augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} \boxed{(b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{36})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(WW) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(XX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(YY) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(ZZ)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(AAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(BBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(i) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d \left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}) \right) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{31}}), (\widehat{T_{31}})$: $(\widehat{G_{31}}, \widehat{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} &((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on
 (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(9)}$ and $(b''_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t})$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484 A
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(i) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}(s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}(s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}(s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + \left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \{ (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \} \\ & \left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + \left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) \left((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)} \right) \{ (\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)} \} \\ & \left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TEN

Holographic Reconstruction Of Spacetime

INTRODUCTION—VARIABLES USED

Communications in Mathematical Physics March 2001, Volume 217, Issue 3, pp 595-622 Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin.

- (1) HST unifies particles and (e&eb) black holes, realizing both as (=) excitations of non-commutative geometrical variables on (eb) a holographic screen.
- (2) Compact extra dimensions are interpreted as (=) finite dimensional unitary representations of (e) super-

algebras, and have (e) no moduli.

- (3) Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of (e) super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity.
- (4) Finite radius de Sitter (dS) spaces have (e) no moduli, and break (e) SUSY with a gravitino mass scaling like $\Lambda^{1/4}$

NOTATION

Module One

HST unifies particles and black holes, realizing both as (=) excitations of non-commutative geometrical variables on (eb) a holographic screen

G_{13} : Category one of HST unifies particles and black holes, realizing

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of excitations of non-commutative geometrical variables on (eb) a holographic screen

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

HST unifies particles and black holes, realizing both as excitations of non-commutative geometrical variables on (eb) a holographic screen

G_{16} : Category one of HST unifies particles and black holes, realizing both as excitations of non-commutative geometrical variables

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of holographic screen

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Compact extra dimensions are interpreted as (=) finite dimensional unitary representations of (e) super-algebras, and have (e) no moduli.

G_{20} : Category one of Compact extra dimensions

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of finite dimensional unitary representations of (e) super-algebras, and have (e) no moduli.

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Compact extra dimensions are interpreted as finite dimensional unitary representations of (e) super-algebras, and have (e) no moduli

G_{24} : Category one of super-algebras, and have (e) no moduli

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Compact extra dimensions are interpreted as finite dimensional unitary representations

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Compact extra dimensions are interpreted as finite dimensional unitary representations of super-algebras, and have (e) no moduli

G_{28} : Category one of no moduli

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Compact extra dimensions are interpreted as finite dimensional unitary representations of super-algebras

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of (e) super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity

G_{32} : Category one of Full field theoretic Fock spaces and continuous moduli

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of emergent phenomena of (e) super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of (e) super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity

G_{36} : Category one of super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Full field theoretic Fock spaces and continuous moduli are both emergent phenomena

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of super-Poincare invariant limits in which the number of holographic degrees of (e) freedom goes to infinity

G_{40} : Category one of freedom goes to infinity

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Full field theoretic Fock spaces and continuous moduli are both emergent phenomena of super-Poincare invariant limits in which the number of holographic degrees

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Finite radius de Sitter (dS) spaces have (e) no moduli, and break (e) SUSY with a gravitino mass scaling like $\Lambda^{1/4}$

G_{44} : Category one of no moduli, and break (e) SUSY with a gravitino mass scaling like $\Lambda^{1/4}$

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Finite radius de Sitter (dS) spaces

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}
 \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\
 \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\
 \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\
 \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\
 \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\
 \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\
 &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\
 &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor}
 \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}
 \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\
 \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\
 \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\
 \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\
 \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\
 \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\
 &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\
 &- (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}
 \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} \left[\begin{array}{c} + (a''_{13})^{(1)}(T_{14}, t) \left[\begin{array}{c} + (a''_{16})^{(2,2)}(T_{17}, t) \left[\begin{array}{c} + (a''_{20})^{(3,3)}(T_{21}, t) \left[\begin{array}{c} + (a''_{24})^{(4,4,4,4)}(T_{25}, t) \left[\begin{array}{c} + (a''_{28})^{(5,5,5,5)}(T_{29}, t) \left[\begin{array}{c} + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \left[\begin{array}{c} + (a''_{36})^{(7,7)}(T_{37}, t) \left[\begin{array}{c} + (a''_{40})^{(8,8)}(T_{41}, t) \left[\begin{array}{c} + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a'_{17})^{(2,2)}(T_{17}, t) + (a'_{21})^{(3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7)}(T_{37}, t) + (a'_{41})^{(8,8)}(T_{41}, t) + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a'_{18})^{(2,2)}(T_{17}, t) + (a'_{22})^{(3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7)}(T_{37}, t) + (a'_{42})^{(8,8)}(T_{41}, t) + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b'_{16})^{(2,2)}(G_{19}, t) - (b'_{20})^{(3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4)}(G_{27}, t) - (b'_{28})^{(5,5,5,5)}(G_{31}, t) - (b'_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7)}(G_{39}, t) - (b'_{40})^{(8,8)}(G_{43}, t) - (b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b'_{17})^{(2,2)}(G_{19}, t) - (b'_{21})^{(3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4)}(G_{27}, t) - (b'_{29})^{(5,5,5,5)}(G_{31}, t) - (b'_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{37})^{(7,7)}(G_{39}, t) - (b'_{41})^{(8,8)}(G_{43}, t) - (b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b'_{18})^{(2,2)}(G_{19}, t) - (b'_{22})^{(3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4)}(G_{27}, t) - (b'_{30})^{(5,5,5,5)}(G_{31}, t) - (b'_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{38})^{(7,7)}(G_{39}, t) - (b'_{42})^{(8,8)}(G_{43}, t) - (b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b'_{13})^{(1)}(G, t)$, $-(b'_{14})^{(1)}(G, t)$, $-(b'_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19}, t) & -(b'_{13})^{(1,1)}(G, t) & -(b''_{20})^{(3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{36})^{(7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19}, t) & -(b'_{14})^{(1,1)}(G, t) & -(b''_{21})^{(3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{37})^{(7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19}, t) & -(b'_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{38})^{(7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b'_{44})^{(9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9)}(G_{47}, t)$, $-(b'_{46})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21}, t) & +(a'_{16})^{(2,2,2)}(T_{17}, t) & +(a'_{13})^{(1,1,1)}(T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{36})^{(7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & + (a''_{21})^{(3)}(T_{21}, t) & + (a'_{17})^{(2,2,2)}(T_{17}, t) & + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{37})^{(7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)}(T_{21}, t), +(a'_{21})^{(3)}(T_{21}, t), +(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t), +(a'_{17})^{(2,2,2)}(T_{17}, t), +(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t), +(a'_{14})^{(1,1,1)}(T_{14}, t), +(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t), +(a'_{37})^{(7,7,7,7)}(T_{37}, t), +(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t), +(a'_{41})^{(8,8,8,8)}(T_{41}, t), +(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t), +(a'_{45})^{(9,9,9)}(T_{45}, t), +(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & - (b'_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{36})^{(7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & - (b'_{21})^{(3)}(G_{23}, t) & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{37})^{(7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} & - (b'_{22})^{(3)}(G_{23}, t) & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{38})^{(7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{22} \quad 72$$

$-(b'_{20})^{(3)}(G_{23}, t), -(b'_{21})^{(3)}(G_{23}, t), -(b'_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t), -(b'_{17})^{(2,2,2)}(G_{19}, t), -(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for

category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{c} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a''_{25})^{(4,4)}(T_{25}, t) & +(a''_{33})^{(6,6,6)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients

for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients

for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition

coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation

coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation

coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} \boxed{(b'_{32})^{(6)}(G_{35}, t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} \boxed{(b'_{33})^{(6)}(G_{35}, t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} \boxed{(b'_{34})^{(6)}(G_{35}, t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6)}(G_{35}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}, \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} \boxed{(a'_{36})^{(7)}(T_{37}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} \boxed{(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} \boxed{(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} \quad 96$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44}$$

$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31} - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(CCC) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(DDD) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(EEE) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(FFF)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(GGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(HHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = 169$$

$$\left(1 + (a_{16})^{(2)} t \right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t \right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(j) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right) \quad 181$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a'_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\textbf{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\bar{G}_{40}^{(1)} - \bar{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b_{42}'')^{(8)}((G_{43})(t), t(t), t) = (b_{42}')^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a'_{44})^{(9)}$ and $(b'_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(\mu_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)}) - (S_2)^{(2)}} \right) \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$$

317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$

318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions

319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)}) - (S_2)^{(6)}} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-
$$v^{(2)} = \frac{G_{16}}{G_{17}}$$
 388

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t) \right) - (a_{25}'')^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)} (v_2)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)} (T_{45}, t) \right) - (a_{45}'')^{(9)} (T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(j) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{45})^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b'_{47})^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ \left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \right. \\
 &\left. \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 &+ \left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \right. \\
 &\left. \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION ELEVEN

Holographic Space-Time: Big Bang To The De Sitter Era

INTRODUCTION—VARIABLES USED

Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era. Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler.

- (1) **T. Banks** presents a holographic theory of inflation and fluctuations. The **inflaton field** is (=) an emergent concept, describing (eb) the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory". **T. Banks** argues that the phrase in quotes is meaningless in the HST formalism. Cite as: arXiv: 1109.2435 [hep-th] **Holographic Space-Time: The Takeaway T. Banks.**
- (2) **T. Banks, W. Fischler** use the formalism of Holographic Space-time (HST) to investigate (e&eb) the claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius.
- (3) This conclusion is much less dramatic in HST than in (eb) the hypothetical models of quantum gravity used in [1].
- (4) In HST there is no dramatic change in particle physics inside the horizon until a time of (e) order the Schwarzschild radius. Report number: UTTG-15-12; TCC-015-12; RUNHETC-2012-17; SCIPP 12/11 arXiv: 1208.4757 [hep-th] **Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler.**

Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era.

- (5) **Tom Banks** reviews the holographic theory of space-time and its applications to (e&eb) cosmology.
- (6) Much of this has appeared before, but this discussion is more unified and concise. He also includes some material on work in progress, whose aim is to understand compactification in terms of (e&eb) finite-dimensional super-algebras. **Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era.**
- (7) The spinorial geometry method of (e) solving Killing spinor equations is reviewed as it applies to (e&eb) six-dimensional (1, 0) supergravity

NOTATION

Module One

Finite radius de Sitter (dS) spaces have no moduli, and break (e) SUSY with a gravitino mass scaling like $\Lambda^{1/4}$

G_{13} : Category one of SUSY with a gravitino mass scaling like $\Lambda^{1/4}$

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Finite radius de Sitter (dS) spaces have no moduli, and break

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

T. Banks presents a

Holographic theory of inflation and fluctuations.

Inflaton field is (=) an emergent concept, describing (eb) the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory". **T. Banks** argues that the phrase in quotes is meaningless in the HST formalism. Cite as: arXiv: 1109.2435 [hep-th] **Holographic Space-Time: The Takeaway T. Banks**

G_{16} : Category one of fluctuations.

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of holographic theory

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Inflaton field is (=) an emergent concept, describing (eb) the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory".

T. Banks argues that the phrase in quotes is meaningless in the HST formalism.

Cite as: arXiv: 1109.2435 [hep-th] **Holographic Space-Time: The Takeaway T. Banks**

G_{20} : Category one of **Inflaton field**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of emergent concept, describing (eb) the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory"

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Inflaton field is an emergent concept, describing (eb) the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory".

G_{24} : Category one of **Inflaton field** is an emergent concept

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory".

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Inflaton field is an emergent concept, describing the geometry of an underlying HST model, rather than "a field associated with (e&eb) a microscopic string theory"

G_{28} : Category one of **Inflaton field** is an emergent concept, describing the geometry of an underlying HST model, rather than "a field; microscopic string theory"

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of microscopic string theory; **Inflaton field** is an emergent concept, describing the geometry of an underlying HST model, rather than "a field"

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

T. Banks, W. Fischler use the

formalism of Holographic Space-time (HST) can be used to investigate (e&eb) the claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{32} : Category one of claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of formalism of Holographic Space-time (HST)

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{36} : Category one of firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes contain a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{40} : Category one of formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes contain a firewall, i.e. an in-falling observer; highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius ;formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes contain a firewall, i.e. an in-falling observer

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes contain a firewall, i.e. an in-falling observer encounters highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{44} : Category one of formalism of Holographic Space-time (HST) can be used to investigate the claim that old black holes contain a firewall, i.e. an in-falling observer encounters highly excited states at a time much shorter

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of light crossing time of the Schwarzschild radius

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)},$
 $(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \end{aligned}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{c} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $[-(b''_{13})^{(1)}(G, t)]$, $[-(b''_{14})^{(1)}(G, t)]$, $[-(b''_{15})^{(1)}(G, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{16})^{(2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{20})^{(3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3)}(G_{23}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b''_{28})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b''_{29})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b''_{30})^{(5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b''_{36})^{(7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b''_{40})^{(8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8)}(G_{43}, t)]$, $[-(b''_{42})^{(8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17}, t)] \quad [+(a''_{13})^{(1,1)}(T_{14}, t)] \quad [+(a''_{20})^{(3,3)}(T_{21}, t)] \\ [+(a''_{24})^{(4,4,4,4)}(T_{25}, t)] \quad [+(a''_{28})^{(5,5,5,5)}(T_{29}, t)] \quad [+(a''_{32})^{(6,6,6,6)}(T_{33}, t)] \\ [+(a''_{36})^{(7,7,7)}(T_{37}, t)] \quad [+(a''_{40})^{(8,8,8)}(T_{41}, t)] \quad [+(a''_{44})^{(9,9)}(T_{45}, t)] \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}[+(a''_{17})^{(2)}(T_{17}, t)] \quad [+(a''_{14})^{(1,1)}(T_{14}, t)] \quad [+(a''_{21})^{(3,3)}(T_{21}, t)] \\ [+(a''_{25})^{(4,4,4,4)}(T_{25}, t)] \quad [+(a''_{29})^{(5,5,5,5)}(T_{29}, t)] \quad [+(a''_{33})^{(6,6,6,6)}(T_{33}, t)] \\ [+(a''_{37})^{(7,7,7)}(T_{37}, t)] \quad [+(a''_{41})^{(8,8,8)}(T_{41}, t)] \quad [+(a''_{45})^{(9,9)}(T_{45}, t)] \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b'_{20})^{(3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b'_{21})^{(3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{37})^{(7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b'_{22})^{(3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{38})^{(7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients

for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients

for category 1,2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for

category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for

category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category

1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients

for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients

for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation

coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation

coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients

for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} \boxed{-(b''_{20})^{(3)}(G_{23}, t)} & \boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} \boxed{-(b''_{21})^{(3)}(G_{23}, t)} & \boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} \boxed{-(b''_{22})^{(3)}(G_{23}, t)} & \boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b'_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} \boxed{+(a''_{24})^{(4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{16})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{20})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} \boxed{+(a''_{25})^{(4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a'_{14})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{17})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{21})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t), (a'_{25})^{(4)}(T_{25}, t), (a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a'_{28})^{(5,5)}(T_{29}, t), +(a'_{29})^{(5,5)}(T_{29}, t), +(a'_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6)}(T_{33}, t), +(a'_{33})^{(6,6)}(T_{33}, t), +(a'_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1)}(T_{14}, t), +(a'_{14})^{(1,1,1,1)}(T_{14}, t), +(a'_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2)}(T_{17}, t), +(a'_{17})^{(2,2,2,2)}(T_{17}, t), +(a'_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3)}(T_{21}, t), +(a'_{21})^{(3,3,3,3)}(T_{21}, t), +(a'_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a'_{46})^{(9,9,9,9)}(T_{45}, t), +(a'_{45})^{(9,9,9,9)}(T_{45}, t), +(a'_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a'_{24})^{(4,4)}(T_{25}, t) & + (a'_{32})^{(6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a'_{25})^{(4,4)}(T_{25}, t) & + (a'_{33})^{(6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a'_{26})^{(4,4)}(T_{25}, t) & + (a'_{34})^{(6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a'_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a'_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a'_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}[-(b''_{28})^{(5)}(G_{31}, t)] & -(b''_{24})^{(4,4)}(G_{27}, t) & -(b''_{32})^{(6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}[-(b''_{29})^{(5)}(G_{31}, t)] & -(b''_{25})^{(4,4)}(G_{27}, t) & -(b''_{33})^{(6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}[-(b''_{30})^{(5)}(G_{31}, t)] & -(b''_{26})^{(4,4)}(G_{27}, t) & -(b''_{34})^{(6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)}[+(a''_{32})^{(6)}(T_{33}, t)] & +(a''_{28})^{(5,5,5)}(T_{29}, t) & +(a''_{24})^{(4,4,4)}(T_{25}, t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)}[+(a''_{33})^{(6)}(T_{33}, t)] & +(a''_{29})^{(5,5,5)}(T_{29}, t) & +(a''_{25})^{(4,4,4)}(T_{25}, t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients

for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients

for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{36})^{(7)}(T_{37}, t)$, $(a''_{37})^{(7)}(T_{37}, t)$, $(a''_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & - (b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) & + (a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)} + (a_{41}'')^{(8)}(T_{41}, t) & + (a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)} + (a_{42}'')^{(8)}(T_{41}, t) & + (a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41}, t)$, $+(a_{41}'')^{(8)}(T_{41}, t)$, $+(a_{42}'')^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

96

$$= (a_{44})^{(9)}G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)} - (b'_{44})^{(9)}(G_{47}, t) & - (b'_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) \quad - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) \quad - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47}, t)$, $-(b''_{45})^{(9)}(G_{47}, t)$, $-(b''_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b'_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a'_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b'_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition:

99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$:

106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} \|(G_{19})' - (G_{19})\| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition:

114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21}' - T_{21}| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23}' - G_{23}| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$:

115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26$$

117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

118

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition:

119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$:

120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30$$

122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

123

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

128

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33}' - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(III) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(JJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(KKK) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(LLL)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ are positive constants and $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37}' - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(MMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(NNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40, 41, 42$$

136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T'_{45}, t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47}) - (G_{47})'| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T'_{45}, t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions

147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions

152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(k) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{ (a'_i)^{(4)} - (a''_i)^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \} ds_{(24)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : (\widehat{G}_{31}), (\widehat{T}_{31}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as 249
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on
 (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right]$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)} (a_{14}')^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)} (a_{14}')^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} + (a_{13})^{(1)} (p_{13})^{(1)} + (a_{14}')^{(1)} (p_{14})^{(1)} + (p_{13})^{(1)} (p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)} (b_{14}')^{(1)} - (b_{13})^{(1)} (b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)} (b_{14}')^{(1)} - (b_{13})^{(1)} (b_{14})^{(1)} - (b_{13}')^{(1)} (r_{14})^{(1)} - (b_{14}')^{(1)} (r_{14})^{(1)} + (r_{13})^{(1)} (r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)} (a_{17}')^{(2)} - (a_{16})^{(2)} (a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(k) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}(G_{43}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWELVE

Holographic Space-Time Does Not Predict Firewalls

INTRODUCTION—VARIABLES USED

Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era. Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler.

- (1) **T. Banks, W. Fischler** use the formalism of Holographic Space-time (HST) to investigate (e&eb) the claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the

Schwarzschild radius. This conclusion is much less dramatic in HST than in (eb) the hypothetical models of quantum gravity used in [1].

- (2) In HST there is no dramatic change in particle physics inside the horizon until a time of (e) order the Schwarzschild radius. Report number: UTTG-15-12; TCC-015-12; RUNHETC-2012-17; SCIPP 12/11 arXiv: 1208.4757 [hep-th] **Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler.**

Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era.

- (3) **Tom Banks** reviews the holographic theory of space-time and its applications to (e&eb) cosmology.
- (4) Much of this has appeared before, but this discussion is more unified and concise. He also includes some material on work in progress, whose aim is to understand compactification in terms of (e&eb) finite-dimensional super-algebras. **Tom Banks 2009 J. Phys A: Math. Theor 42 304002 doi:10.1088/1751-8113/42/30/304002 Holographic space-time from the Big Bang to the de Sitter era.**
- (5) The spinorial geometry method of (e) solving Killing spinor equations is reviewed as it applies to (e&eb) six-dimensional (1, 0) supergravity

NOTATION

Module One

T. Banks, W. Fischler use the

formalism of Holographic Space-time (HST) to investigate (e&eb) the claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

. This conclusion is much less dramatic in HST than in (eb) the hypothetical models of quantum gravity used in [1].(Please see the original paper for references)

G_{13} : Category one of formalism of Holographic Space-time (HST); old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius; formalism of Holographic Space-time (HST)

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

formalism of Holographic Space-time (HST) to investigate (e&eb) the claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{16} : Category one of formalism of Holographic Space-time (HST); old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius ;formalism of Holographic Space-time (HST)

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain (e) a firewall, i.e. an in-falling observer encounters (e&eb) highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{20} : Category one of formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain a firewall, i.e. an in-falling observer; highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius ;formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain a firewall, i.e. an in-falling observer

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain a firewall, i.e. an in-falling observer encounters highly excited states at a time much shorter than (eb) the light crossing time of the Schwarzschild radius

G_{24} : Category one of formalism of Holographic Space-time (HST) to investigate the claim of [1] that old black holes contain a firewall, i.e. an in-falling observer encounters highly excited states at a time much shorter

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of light crossing time of the Schwarzschild radius

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

In HST there is no dramatic change in particle physics inside the horizon until a time of order the Schwarzschild radius.

Report number: UTTG-15-12; TCC-015-12; RUNHETC-2012-17; SCIPP 12/11 arXiv: 1208.4757 [hep-th]

Holographic Space-Time Does Not Predict Firewalls T. Banks, W. Fischler

G_{28} : Category one of time of order the Schwarzschild radius

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of particle physics in HST remains unchanged (or approximately so)

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

The

spinorial geometry method of solving Killing spinor equations is reviewed as it applies to (e&eb) six-dimensional (1, 0) supergravity

G_{32} : Category one of spinorial geometry method of solving Killing spinor equations; six-dimensional (1, 0) supergravity

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of six-dimensional (1, 0) supergravity ;spinorial geometry method of solving Killing spinor equations

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

T Thiemann 1998 Class Quantum Grav 15 839 doi:10.1088/0264-9381/15/4/011 Quantum spin dynamics (QSD)

- (1) An anomaly-free operator corresponding to (e&eb) the Wheeler - DeWitt constraint of Lorentzian, four-dimensional, canonical, non-perturbative vacuum gravity is constructed in (eb) the continuum.
- (2) This operator is entirely free (e) of factor-ordering singularities and can be defined in (eb) symmetric and non-symmetric form.
- (3) Authors work in the real connection representation and obtain (eb) a well defined quantum theory. The action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting (e&eb) the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states.

- (4) Authors argue that the spin-network representation is (=) the 'nonlinear Fock representation' of quantum gravity, thus justifying the term 'quantum spin dynamics (QSD)'. This paper is the first in a series of seven papers with the title 'quantum spin dynamics (QSD)'.

An anomaly-free operator corresponding to (e&eb) the Wheeler - DeWitt constraint of Lorentzian, four-dimensional, canonical, non-perturbative vacuum gravity is (=) constructed in the continuum

G_{36} : Category one of anomaly-free operator; Wheeler - DeWitt constraint of Lorentzian, four-dimensional, canonical, non-perturbative vacuum gravity is constructed in (eb) the continuum

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Wheeler - DeWitt constraint of Lorentzian, four-dimensional, canonical, non-perturbative vacuum gravity is constructed in (eb) the continuum ;anomaly-free operator

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

An anomaly-free operator corresponding to the Wheeler - DeWitt constraint of Lorentzian, four-dimensional, canonical, non-perturbative vacuum gravity is (=) constructed in the continuum

G_{40} : Category one of anomaly-free operator corresponding to the Wheeler - DeWitt constraint of Lorentzian, four-dimensional, canonical, non-perturbative vacuum gravity

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of constructed in the continuum

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

This

operator is entirely free (e) of factor-ordering singularities and can be defined in (eb) symmetric and non-symmetric form

G_{44} : Category one of factor-ordering singularities and can be defined in (eb) symmetric and non-symmetric form

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of operator

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \end{aligned}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}), t) =$ First detritions factor

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t), +(a''_{21})^{(3)}(T_{21}, t), +(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{l} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{l} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{l} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{l} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{l} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients

for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients

for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients

for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition

coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$\begin{aligned} & - \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86 \end{aligned}$$

$$\begin{aligned} \frac{dG_{34}}{dt} &= (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87 \end{aligned}$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation

coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation

coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)} G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41}, t)} & \boxed{+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41}, t)} & \boxed{+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(OOO) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(PPP) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(QQQ) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(RRR)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T'_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(SSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(TTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(l) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right] \quad 181$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)} \right) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is 188
analogous with the preceding one. An analogous property is true if G_{14} is bounded from below.

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a'_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} ((b'')^{(7)}((G_{39})(t), t)) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

Definition of $(\bar{G}_{43}), (\bar{T}_{43})$: $(\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 269$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 270$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$$

317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$

318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions

319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

$$\text{roots of the equations } (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)} , \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)} , \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)} (v_2)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)} (T_{45}, t) \right) - (a_{45}'')^{(9)} (T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(l) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTEEN

Factor-Ordering Singularities Free Operator

INTRODUCTION—VARIABLES USED

T Thiemann 1998 Class Quantum Grav 15 839 doi:10.1088/0264-9381/15/4/011 Quantum spin dynamics (QSD)

- (1) This operator is entirely free (e) of factor-ordering singularities and can be defined in (eb) symmetric and non-symmetric form.
- (2) Authors work in the real connection representation and obtain (eb) a well defined quantum theory. The action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting (e&eb) the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states.
- (3) Authors argue that the spin-network representation is (=) the 'nonlinear Fock representation' of quantum gravity, thus justifying the term 'quantum spin dynamics (QSD)'. This paper is the first in a series of seven papers with the title 'quantum spin dynamics (QSD)'.

NOTATION

Module One

Operator is entirely free of factor-ordering singularities and can be defined in (eb) symmetric and non-symmetric form.

G_{13} : Category one of Operator is entirely free of factor-ordering singularities defined

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of symmetric and non-symmetric form.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Authors work in the

Work is in real connection representation and obtains (eb) a well defined quantum theory.

The

action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting (e&eb) the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states.

G_{16} : Category one of Work

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of real connection representation and obtains (eb) a well defined quantum theory.

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting (e&eb) the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{20} : Category one of action of the Wheeler - DeWitt constraint; spin-network states is by annihilating, creating and rerouting (e&eb) the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of spin-network states is by annihilating, creating and rerouting (e&eb) the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states; action of the Wheeler - DeWitt constraint

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with (e&eb) the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{24} : Category one of action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting the quanta of angular momentum; edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states; action of the Wheeler - DeWitt constraint on (e&eb) spin-network states is by annihilating, creating and rerouting the quanta of angular momentum

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

action of the Wheeler - DeWitt constraint on spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{28} : Category one of underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of action of the Wheeler - DeWitt constraint on spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with the edges

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

action of the Wheeler - DeWitt constraint on spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with the edges of the underlying graph while the ADM energy is (=) essentially diagonalized by (e) the spin-network states

G_{32} : Category one of action of the Wheeler - DeWitt constraint on spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with the edges of the underlying graph while the ADM energy

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of essentially diagonalized by (e) the spin-network states

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

action of the Wheeler - DeWitt constraint on spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with the edges of the underlying graph while the ADM energy

is essentially diagonalized by (e) the spin-network states

G_{36} : Category one of spin-network states

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of action of the Wheeler - DeWitt constraint on spin-network states is by annihilating, creating and rerouting the quanta of angular momentum associated with the edges of the underlying graph while the ADM energy is essentially diagonalized

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Authors argue that the

Spin-network representation is (=) the 'nonlinear Fock representation' of quantum gravity, thus justifying the term 'quantum spin dynamics (QSD)'.

This paper is the first in a series of seven papers with the title 'quantum spin dynamics (QSD)'.

G_{40} : Category one of Spin-network representation

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of 'nonlinear Fock representation' of quantum gravity, thus justifying the term 'quantum spin dynamics (QSD)'.

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

G_{44} : Category one of nonlinear Fock representation' of quantum gravity; Wheeler - DeWitt constraint on spin-network states

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Wheeler - DeWitt constraint on spin-network states; nonlinear Fock representation' of quantum gravity

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}), t) =$ First detritions factor

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$, $(a''_{21})^{(3,3)}(T_{21}, t)$, $(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7)}(T_{37}, t)$, $(a''_{37})^{(7,7)}(T_{37}, t)$, $(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8)}(T_{41}, t)$, $(a''_{41})^{(8,8)}(T_{41}, t)$, $(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19}, t) & -(b'_{13})^{(1,1)}(G, t) & -(b''_{20})^{(3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{36})^{(7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19}, t) & -(b'_{14})^{(1,1)}(G, t) & -(b''_{21})^{(3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{37})^{(7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19}, t) & -(b'_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{38})^{(7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b'_{44})^{(9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9)}(G_{47}, t)$, $-(b'_{46})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21}, t) & +(a'_{16})^{(2,2,2)}(T_{17}, t) & +(a'_{13})^{(1,1,1)}(T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{36})^{(7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & + (a''_{21})^{(3)}(T_{21}, t) & + (a'_{17})^{(2,2,2)}(T_{17}, t) & + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{37})^{(7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)}(T_{21}, t), +(a'_{21})^{(3)}(T_{21}, t), +(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t), +(a'_{17})^{(2,2,2)}(T_{17}, t), +(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t), +(a'_{14})^{(1,1,1)}(T_{14}, t), +(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t), +(a'_{37})^{(7,7,7,7)}(T_{37}, t), +(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t), +(a'_{41})^{(8,8,8,8)}(T_{41}, t), +(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t), +(a'_{45})^{(9,9,9)}(T_{45}, t), +(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & - (b'_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{36})^{(7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & - (b'_{21})^{(3)}(G_{23}, t) & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{37})^{(7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} & - (b'_{22})^{(3)}(G_{23}, t) & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{38})^{(7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{22} \quad 72$$

$-(b'_{20})^{(3)}(G_{23}, t), -(b'_{21})^{(3)}(G_{23}, t), -(b'_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t), -(b'_{17})^{(2,2,2)}(G_{19}, t), -(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for

category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{c} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a''_{25})^{(4,4)}(T_{25}, t) & +(a''_{33})^{(6,6,6)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients

for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients

for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition

coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation

coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation

coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t)$, $-(b''_{33})^{(6)}(G_{35}, t)$, $-(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} \boxed{(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} \boxed{(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a'_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a'_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(UUU) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(VVV) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(WWW) \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(XXX)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T'_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(YYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(ZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with the constants $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(m) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right) \quad 181$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)} \right) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\bar{A}_{16})^{(2)} + (\bar{P}_{16})^{(2)} (\bar{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\bar{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

$$\textbf{Definition of } (\widetilde{G_{31}}, \widetilde{T_{31}}): (\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as 249
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{-(\hat{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\hat{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\hat{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\tilde{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\bar{G}_{40}^{(1)} - \bar{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a'_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a'_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a'_{40})^{(8)}$ and $(b'_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(\mu_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$$

317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$

318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions

319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}, t) - (b''_{25})^{(4)}((G_{27}, t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

$$\text{roots of the equations } (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)} , \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)} , \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$-\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(m) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]}, \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}, \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]}, \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]}, \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]}, \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]}, \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FOURTEEN

Spinorial Geometry, Horizons And Superconformal Symmetry

INTRODUCTION—VARIABLES USED

M Akyol and G Papadopoulos 2014 Class. Quantum Grav 31 123001 doi:10.1088/0264-9381/31/12/123001 Spinorial geometry, horizons and superconformal symmetry in six dimensions

- (1) In particular, it is explained how the method **of Spinorial geometry, horizons and superconformal symmetry in six dimensions** is used to identify (e&eb) both the fractions of supersymmetry preserved by (e) the geometry of all supersymmetric backgrounds.
- (2) Then two applications **Spinorial geometry, horizons and superconformal symmetry in six dimensions** are described to systems that exhibit (eb) superconformal symmetry.
- (3) The first is the proof that some six-dimensional black hole horizons are (=) locally isometric to $AdS_3 \times \Sigma_3$, where Σ_3 is diffeomorphic to S^3 .
- (4) The second one is a description of all supersymmetric solutions of (e) six-dimensional (1, 0) superconformal theories and in particular of their brane solitons. **M Akyol and G Papadopoulos 2014 Class. Quantum Grav 31 123001 doi:10.1088/0264-9381/31/12/123001 Spinorial geometry, horizons and superconformal symmetry in six dimensions.**

S S Y Chua et al 2014 Class. Quantum Grav 31 183001 doi:10.1088/0264-9381/31/18/183001 Quantum squeezed light in gravitational-wave detectors.

- (5) Field of squeezed states for gravitational-wave (GW) detector enhancement is rapidly maturing. In this review paper, **S S Y Chua1, B J J Slagmolen, D A Shaddock and D E McClelland** provide an analysis of the field circa 2013. They begin by outlining the concept and description of quantum squeezed states. This is followed by an overview of how quantum squeezed states can improve (eb) GW detection, and the requirements (e) on squeezed states to achieve such enhancement.
- (6) Next, an overview of current technology for producing squeezed states, using (e) atoms, optomechanical methods and nonlinear crystals, is provided. They finally highlight the milestone squeezing implementation experiments at the GEO600 and LIGO GW detectors. **S S Y Chua et al 2014 Class. Quantum Grav 31 183001 doi:10.1088/0264-9381/31/18/183001 Quantum squeezed light in gravitational-wave detectors**

NOTATION

Module One

In particular, it is explained how the method **of Spinorial geometry, horizons and superconformal symmetry in six dimensions** is used to identify (e&eb) both the fractions of supersymmetry preserved by (e) the geometry of all supersymmetric backgrounds

G_{13} : Category one of identification of both the fractions of supersymmetry preserved by (e) the geometry of all supersymmetric backgrounds

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **Spinorial geometry, horizons and superconformal symmetry in six dimensions**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

In particular, it is explained how the method of

Spinorial geometry, horizons and superconformal symmetry in six dimensions is used to identify both the fractions of supersymmetry preserved by (e) the geometry of all supersymmetric backgrounds

G_{16} : Category one of geometry of all supersymmetric backgrounds

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of **Spinorial geometry, horizons and superconformal symmetry in six dimensions** is used to identify both the fractions of supersymmetry preserved

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Then two applications

Spinorial geometry, horizons and superconformal symmetry in six dimensions are described to systems that exhibit superconformal symmetry

G_{20} : Category one of **Spinorial geometry, horizons and superconformal symmetry in six dimensions**; systems that exhibit superconformal symmetry

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of systems that exhibit superconformal symmetry ;**Spinorial geometry, horizons and superconformal symmetry in six dimensions**

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

The first is the proof that some six-dimensional black hole horizons are (=) locally isometric to $AdS_3 \times \Sigma_3$, where Σ_3 is diffeomorphic to

S3

G_{24} : Category one of six-dimensional black hole horizons

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of locally isometric to $AdS_3 \times \Sigma_3$, where Σ_3 is diffeomorphic to S^3

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The second one is a description of

all supersymmetric solutions of (e) six-dimensional (1, 0) superconformal theories and in particular of their brane solitons.

M Akyol and G Papadopoulos 2014 Class. Quantum Grav 31 123001 doi:10.1088/0264-9381/31/12/123001 Spinorial geometry, horizons and superconformal symmetry in six dimensions

G_{28} : Category one of six-dimensional (1, 0) superconformal theories and in particular of their brane solitons

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of all supersymmetric solutions

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

all supersymmetric solutions of six-dimensional (1, 0) superconformal theories and in particular of their brane solitons

G_{32} : Category one of all supersymmetric solutions of six-dimensional (1, 0) superconformal theories; brane solitons

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of brane solitons ;all supersymmetric solutions of six-dimensional (1, 0) superconformal theories

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Field of squeezed states for (e) gravitational-wave (GW) detector enhancement is rapidly maturing.

In this review paper, **S S Y Chua¹, B J J Slagmolen, D A Shaddock and D E McClelland** provide an analysis of the field circa 2013. They begin by outlining the concept and description of quantum squeezed states. This is followed by an overview of how quantum squeezed states can improve (eb) GW detection, and the requirements (e) on squeezed states to achieve such enhancement

G_{36} : Category one of squeezed states; gravitational-wave (GW) detector enhancement

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of gravitational-wave (GW) detector enhancement; squeezed states

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

In this review paper, **S S Y Chua¹, B J J Slagmolen, D A Shaddock and D E McClelland** provide an analysis of the field circa 2013. They begin by outlining the concept and description of quantum squeezed states.

This is followed by an overview of how

quantum squeezed states can improve (eb) GW detection, and the requirements (e) on squeezed states to achieve such enhancement

G_{40} : Category one of quantum squeezed states

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of GW detection, and the requirements (e) on squeezed states to achieve such enhancement

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

quantum squeezed states can improve GW detection, and the requirements on squeezed states to achieve (eb) such enhancement

G_{44} : Category one of quantum squeezed states can improve GW detection, and the requirements on

squeezed states

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of enhancements

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ + (a''_{20})^{(3)}(T_{21}, t) &= \text{First augmentation factor} \\ - (b''_{20})^{(3)}(G_{23}, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20 \\ \frac{dG_{26}}{dt} &= (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} & 21 \\ \frac{dT_{24}}{dt} &= (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} & 22 \\ \frac{dT_{25}}{dt} &= (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} & 23 \\ \frac{dT_{26}}{dt} &= (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} & 24 \\ + (a''_{24})^{(4)}(T_{25}, t) &= \text{First augmentation factor} \\ - (b''_{24})^{(4)}((G_{27}), t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\begin{aligned}\frac{dG_{28}}{dt} &= (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} & 25 \\ \frac{dG_{29}}{dt} &= (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} & 26 \\ \frac{dG_{30}}{dt} &= (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} & 27 \\ \frac{dT_{28}}{dt} &= (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} & 28 \\ \frac{dT_{29}}{dt} &= (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} & 29\end{aligned}$$

$$\begin{aligned}\frac{dT_{30}}{dt} &= (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \\ + (a''_{28})^{(5)}(T_{29}, t) &= \text{First augmentation factor} \\ - (b''_{28})^{(5)}((G_{31}), t) &= \text{First detritions factor}\end{aligned}$$

30

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\begin{aligned}\frac{dG_{32}}{dt} &= (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} & 31 \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} & 32 \\ \frac{dG_{34}}{dt} &= (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} & 33 \\ \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} & 34 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} & 35 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} & 36 \\ + (a''_{32})^{(6)}(T_{33}, t) &= \text{First augmentation factor}\end{aligned}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\begin{aligned}\frac{dG_{36}}{dt} &= (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} & 42 \\ + (a''_{36})^{(7)}(T_{37}, t) &= \text{First augmentation factor}\end{aligned}$$

Module Numbered Eight

The differential system of this model is now

$$\begin{aligned}\frac{dG_{40}}{dt} &= (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} & 44 \\ \frac{dG_{42}}{dt} &= (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} & 52\end{aligned}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - \left[\begin{array}{c} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{13})^{(1)}(G, t)}$, $\boxed{-(b'_{14})^{(1)}(G, t)}$, $\boxed{-(b'_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{37})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b'_{36})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} \boxed{+(a''_{16})^{(2)}(T_{17}, t)} \boxed{+(a''_{13})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} \boxed{+(a''_{17})^{(2)}(T_{17}, t)} \boxed{+(a''_{14})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{21})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} \boxed{+(a''_{18})^{(2)}(T_{17}, t)} \boxed{+(a''_{15})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{22})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{18} \quad 63$$

Where $\boxed{+(a'_{16})^{(2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2)}(T_{17}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{13})^{(1,1)}(T_{14}, t)}$, $\boxed{+(a'_{14})^{(1,1)}(T_{14}, t)}$, $\boxed{+(a'_{15})^{(1,1)}(T_{14}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{20})^{(3,3,3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3,3,3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)}(G_{19}, t) & -(b'_{13})^{(1,1)}(G, t) & -(b'_{20})^{(3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b'_{36})^{(7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)}(G_{19}, t) & -(b'_{14})^{(1,1)}(G, t) & -(b'_{21})^{(3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b'_{37})^{(7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)}(G_{19}, t) & -(b'_{15})^{(1,1)}(G, t) & -(b'_{22})^{(3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category

1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{l} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b'_{16})^{(2,2,2)}(G_{19}, t) - (b'_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{l} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) - (b'_{17})^{(2,2,2)}(G_{19}, t) - (b'_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$[-(b'_{20})^{(3)}(G_{23}, t)], [-(b'_{21})^{(3)}(G_{23}, t)], [-(b'_{22})^{(3)}(G_{23}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b'_{16})^{(2,2,2)}(G_{19}, t)], [-(b'_{17})^{(2,2,2)}(G_{19}, t)], [-(b'_{18})^{(2,2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b'_{13})^{(1,1,1)}(G, t)], [-(b'_{14})^{(1,1,1)}(G, t)], [-(b'_{15})^{(1,1,1)}(G, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b'_{36})^{(7,7,7,7)}(G_{39}, t)], [-(b'_{37})^{(7,7,7,7)}(G_{39}, t)], [-(b'_{38})^{(7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b'_{40})^{(8,8,8,8)}(G_{43}, t)], [-(b'_{41})^{(8,8,8,8)}(G_{43}, t)], [-(b'_{42})^{(8,8,8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b'_{46})^{(9,9,9)}(G_{47}, t)], [-(b'_{45})^{(9,9,9)}(G_{47}, t)], [-(b'_{44})^{(9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)}[+(a''_{24})^{(4)}(T_{25}, t)] & +(a'_{28})^{(5,5,)}(T_{29}, t) & +(a'_{32})^{(6,6,)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)}[+(a''_{25})^{(4)}(T_{25}, t)] & +(a'_{29})^{(5,5,)}(T_{29}, t) & +(a'_{33})^{(6,6,)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)}[+(a''_{26})^{(4)}(T_{25}, t)] & +(a'_{30})^{(5,5,)}(T_{29}, t) & +(a'_{34})^{(6,6,)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$[(a'_{24})^{(4)}(T_{25}, t)], [(a'_{25})^{(4)}(T_{25}, t)], [(a'_{26})^{(4)}(T_{25}, t)]$ are first augmentation coefficients category 1, 2 3

$[(a'_{28})^{(5,5,)}(T_{29}, t)], [(a'_{29})^{(5,5,)}(T_{29}, t)], [(a'_{30})^{(5,5,)}(T_{29}, t)]$ are second augmentation coefficient for category 1, 2 and 3

$[(a'_{32})^{(6,6,)}(T_{33}, t)], [(a'_{33})^{(6,6,)}(T_{33}, t)], [(a'_{34})^{(6,6,)}(T_{33}, t)]$ are third augmentation coefficient for category 1, 2 and 3

$[(a'_{13})^{(1,1,1,1)}(T_{14}, t)], [(a'_{14})^{(1,1,1,1)}(T_{14}, t)], [(a'_{15})^{(1,1,1,1)}(T_{14}, t)]$

are fourth augmentation coefficients for category 1, 2 and 3

$$+(a''_{16})^{(2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2)}(T_{17}, t)$$

are fifth augmentation coefficients for category 1, 2 and 3

$$+(a''_{20})^{(3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3)}(T_{21}, t)$$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a'_{24})^{(4,4)}(T_{25}, t) & + (a'_{32})^{(6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a'_{25})^{(4,4)}(T_{25}, t) & + (a'_{33})^{(6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a'_{26})^{(4,4)}(T_{25}, t) & + (a'_{34})^{(6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a'_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a'_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a'_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a'_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a'_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b'_{24})^{(4,4)}(G_{27}, t) & - (b'_{32})^{(6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1)}(G, t) & - (b'_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b'_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b'_{25})^{(4,4)}(G_{27}, t) & - (b'_{33})^{(6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1)}(G, t) & - (b'_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b'_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b'_{26})^{(4,4)}(G_{27}, t) & - (b'_{34})^{(6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1)}(G, t) & - (b'_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1, 2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1, 2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1, 2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1, 2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)} \text{ sixth augmentation}$$

coefficients

$$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)}$$

seventh augmentation coefficients

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$$

Eighth augmentation coefficients

$$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)} \text{ ninth augmentation}$$

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{l} \boxed{(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) - (b''_{28})^{(5,5,5)}(G_{31}, t) - (b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{l} \boxed{(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) - (b''_{29})^{(5,5,5)}(G_{31}, t) - (b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{l} \boxed{(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) - (b''_{30})^{(5,5,5)}(G_{31}, t) - (b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{34} \quad 90$$

$$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6)}(G_{35}, t)} \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t)} \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$$

are eighth detrition coefficients for category 1, 2, and 3

$$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \begin{bmatrix} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \begin{bmatrix} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}$, $(b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}$, $(\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}$, $(\hat{B}_{13})^{(1)}$, $(p_i)^{(1)}$, $(r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}$, $(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}$, $(\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31} - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(AAAA) (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(BBBB) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(CCCC) \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(DDDD)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T'_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(EEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(FFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(n) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$\left(1 + (a_{36})^{(7)} t \right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} =$$

$$\left(1 + (a_{40})^{(8)} t \right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

180

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

181

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t \right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose

182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is 188
analogous with the preceding one. An analogous property is true if G_{14} is bounded from below.

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})_1$ it follows $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \quad \text{If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), ((G_{27})^{(2)}, (T_{27})^{(2)}) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}\right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{-(\hat{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} \} ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
 $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
 $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
 $(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)}) - (S_2)^{(2)}} \right) \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}^{(4)}) = (a_{25}^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}^{(4)}) = (b_{25}^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case**.

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$** .

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(n) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]}, \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}, \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]}, \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]}, \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]}, \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]}, \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}) T_{13}^* \mathbb{G}_j \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}) T_{14}^* \mathbb{G}_j \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}) T_{15}^* \mathbb{G}_j \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a_{16}')^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a_{17}')^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a_{18}')^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b_{16}')^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b_{17}')^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b_{18}')^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a_{20}')^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a_{21}')^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a_{22}')^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b_{20}')^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)})$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)})$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTEEN

Holographic Space-Time And Matrix Theory

INTRODUCTION—VARIABLES USED

DOI: <http://dx.doi.org/10.1103/PhysRevD.84.086008> © 2011 American Physical Society **Fuzzy geometry via the spinor bundle, with applications to holographic space-time and matrix theory Phys. Rev. D 84, 086008 – Published 25 October 2011 Tom Banks and John Kehayias.**

- (1) **Tom Banks and John Kehayias** present a new framework for defining fuzzy approximations to (e) geometry in terms of (e&eb) a cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator.
- (2) This framework does not (e) require a symplectic form on the manifold, and is (=) completely rotation invariant on (eb) an arbitrary n-sphere.
- (3) The framework is motivated by the formalism of holographic space-time, whose fundamental variables are (=) sections of the spinor bundle over (e&eb) a compact Euclidean manifold.
- (4) The strong holographic principle requires (e) the space of these sections to be (=) finite dimensional.
- (5) They discuss applications of fuzzy spinor geometry to (e&eb) holographic space-time and to matrix theory. DOI: <http://dx.doi.org/10.1103/PhysRevD.84.086008> © 2011 American Physical Society **Fuzzy geometry via the spinor bundle, with applications to holographic space-time and matrix theory Phys. Rev. D 84, 086008 – Published 25 October 2011 Tom Banks and John Kehayias.**
- (6) **Chaolun Wu, Shao-Feng Wu** show that Horava-Lifshitz gravity theory can be employed (e) as a covariant framework to build (eb) an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry.
- (7) The key to this formalism is a set of correspondence relations that maps all the field degrees of freedom in the Horava-Lifshitz gravity theory to (e&eb) external background (source) fields among others in the effective action of the quantum Hall effect, according to their symmetry transformation properties.

NOTATION

Module One

Tom Banks and John Kehayias present a new framework for defining fuzzy approximations to (e) geometry in terms of (e&eb) a cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator.

G_{13} : Category one of geometry in terms of (e&eb) a cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of fuzzy approximations

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

fuzzy approximations to geometry in terms of (e&eb) a cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator

G_{16} : Category one of fuzzy approximations to geometry; cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator ;fuzzy approximations to geometry

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

fuzzy approximations to geometry in terms of a cutoff on the spectrum of (e) the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator

G_{20} : Category one of Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of fuzzy approximations to geometry in terms of a cutoff on the spectrum

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

fuzzy approximations to geometry in terms of a cutoff on the spectrum of the Dirac operator, and a generalization of (e&eb) it that we call the Dirac-flux operator

G_{24} : Category one of Dirac-flux operator

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of fuzzy approximations to geometry in terms of a cutoff on the spectrum of the Dirac operator, and a generalization

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

This

framework does not (e) require a symplectic form on the manifold, and is (=) completely rotation invariant on (eb) an arbitrary n-sphere

G_{28} : Category one of requirements of a symplectic form on the manifold, and is (=) completely rotation invariant on (eb) an arbitrary n-sphere

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of frameworks

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

framework does not require a symplectic form on the manifold, and is (=) completely rotation invariant on (eb) an arbitrary n-sphere

G_{32} : Category one of framework does not require a symplectic form on the manifold

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of completely rotation invariant on (eb) an arbitrary n-sphere

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

framework does not require a symplectic form on the manifold, and is completely rotation invariant on (eb) an arbitrary n-sphere

G_{36} : Category one of framework does not require a symplectic form on the manifold, and is completely rotation invariant

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of arbitrary n-sphere (and there could be many such n spheres)

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

The

framework is motivated by the formalism of holographic space-time, whose fundamental variables are (=) sections of the spinor bundle over (e&eb) a compact Euclidean manifold

G_{40} : Category one of framework is motivated by the formalism of holographic space-time, whose fundamental variables

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of sections of the spinor bundle over (e&eb) a compact Euclidean manifold

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

framework is motivated by the formalism of holographic space-time, whose fundamental variables are sections of the spinor bundle over (e&eb) a compact Euclidean manifold

G_{44} : Category one of framework is motivated by the formalism of holographic space-time, whose fundamental variables are sections of the spinor bundle; compact Euclidean manifold

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of compact Euclidean manifold ;framework is motivated by the formalism of holographic space-time, whose fundamental variables are sections of the spinor bundle

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}
 \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\
 \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\
 \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\
 \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\
 \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\
 \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\
 &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\
 &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}
 \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}
 \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\
 \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\
 \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\
 \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\
 \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\
 \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\
 &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\
 &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}
 \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} \left[\begin{array}{c} + (a''_{13})^{(1)}(T_{14}, t) \left[\begin{array}{c} + (a''_{16})^{(2,2)}(T_{17}, t) \left[\begin{array}{c} + (a''_{20})^{(3,3)}(T_{21}, t) \left[\begin{array}{c} + (a''_{24})^{(4,4,4,4)}(T_{25}, t) \left[\begin{array}{c} + (a''_{28})^{(5,5,5,5)}(T_{29}, t) \left[\begin{array}{c} + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \left[\begin{array}{c} + (a''_{36})^{(7,7)}(T_{37}, t) \left[\begin{array}{c} + (a''_{40})^{(8,8)}(T_{41}, t) \left[\begin{array}{c} + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b'_{13})^{(1)}(G, t)$, $-(b'_{14})^{(1)}(G, t)$, $-(b'_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19}, t) & -(b'_{13})^{(1,1)}(G, t) & -(b''_{20})^{(3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{36})^{(7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19}, t) & -(b'_{14})^{(1,1)}(G, t) & -(b''_{21})^{(3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{37})^{(7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19}, t) & -(b'_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b'_{38})^{(7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b'_{44})^{(9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9)}(G_{47}, t)$, $-(b'_{46})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21}, t) & +(a'_{16})^{(2,2,2)}(T_{17}, t) & +(a'_{13})^{(1,1,1)}(T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{36})^{(7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & + (a''_{21})^{(3)}(T_{21}, t) & + (a'_{17})^{(2,2,2)}(T_{17}, t) & + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ & + (a'_{37})^{(7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ & + (a'_{38})^{(7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)}(T_{21}, t), +(a'_{21})^{(3)}(T_{21}, t), +(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t), +(a'_{17})^{(2,2,2)}(T_{17}, t), +(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t), +(a'_{14})^{(1,1,1)}(T_{14}, t), +(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t), +(a'_{37})^{(7,7,7,7)}(T_{37}, t), +(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t), +(a'_{41})^{(8,8,8,8)}(T_{41}, t), +(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t), +(a'_{45})^{(9,9,9)}(T_{45}, t), +(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & - (b'_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ & - (b'_{36})^{(7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & - (b'_{21})^{(3)}(G_{23}, t) & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ & - (b'_{37})^{(7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} & - (b'_{22})^{(3)}(G_{23}, t) & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ & - (b'_{38})^{(7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b'_{20})^{(3)}(G_{23}, t), -(b'_{21})^{(3)}(G_{23}, t), -(b'_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t), -(b'_{17})^{(2,2,2)}(G_{19}, t), -(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for

category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category

1,2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition

coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition

coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients

for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for

category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for

category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{c} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients

category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation

coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation

coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$

are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$

are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a''_{25})^{(4,4)}(T_{25}, t) & +(a''_{33})^{(6,6,6)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients

for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients

for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition

coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{cccc} (a'_{32})^{(6)} & + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{cccc} (a'_{33})^{(6)} & + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{cccc} (a'_{34})^{(6)} & + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation

coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation

coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t)$, $-(b''_{33})^{(6)}(G_{35}, t)$, $-(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $(a'_{14})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $(a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$(a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \begin{bmatrix} -(b''_{38})^{(7)}(G_{39}, t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \end{bmatrix} \\ \begin{bmatrix} -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \end{bmatrix} \\ \begin{bmatrix} -(b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \begin{bmatrix} (a'_{40})^{(8)} \begin{bmatrix} +(a''_{40})^{(8)}(T_{41}, t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \end{bmatrix} \\ \begin{bmatrix} +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \end{bmatrix} \\ \begin{bmatrix} +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \begin{bmatrix} (a'_{41})^{(8)} \begin{bmatrix} +(a''_{41})^{(8)}(T_{41}, t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \end{bmatrix} \\ \begin{bmatrix} +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \end{bmatrix} \\ \begin{bmatrix} +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} \quad 96$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44}$$

$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(GGGG) (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(HHHH) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(IIII) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(JJJJ)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(KKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(LLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42$$

136

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$$

138

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)}$$

139

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)}$$

140

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)}$$

141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t}$$

142

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$$

143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(o) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right) \quad 181$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}\right), \left((G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})$ it follows $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \quad \text{If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$|\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} +$$

$$\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} \leq$$

$$\frac{1}{(\bar{M}_{20})^{(3)}} \left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)} \right) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{-(\hat{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)}s_{(28)}} e^{(\hat{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a'_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\bar{G}_{40}^{(1)} - \bar{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right)$, $t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)}) \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\begin{aligned} &\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations} \\ &(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0 \end{aligned}$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\begin{aligned} &\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations} \\ &(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0 \end{aligned}$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$$

317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$

318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions

319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)}) - (S_2)^{(6)}} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:-

$$\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(o) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a_{16}')^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a_{17}')^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a_{18}')^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b_{16}')^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b_{17}')^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b_{18}')^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a_{20}')^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a_{21}')^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a_{22}')^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b_{20}')^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)})$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)})$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ \left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \right. \\
 &\left. \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 &+ \left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \right. \\
 &\left. \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTEEN

Fuzzy Spinor Geometry And Holographic Space-Time

INTRODUCTION—VARIABLES USED

DOI: <http://dx.doi.org/10.1103/PhysRevD.84.086008> © 2011 American Physical Society **Fuzzy geometry via the spinor bundle, with applications to holographic space-time and matrix theory Phys. Rev. D 84, 086008 – Published 25 October 2011 Tom Banks and John Kehayias.**

- (1) The strong holographic principle requires (e) the space of these sections to be (=) finite dimensional.
- (2) They discuss applications of fuzzy spinor geometry to (e&eb) holographic space-time and to matrix theory. DOI: <http://dx.doi.org/10.1103/PhysRevD.84.086008> © 2011 American Physical Society **Fuzzy geometry via the spinor bundle, with applications to holographic space-time and matrix theory Phys. Rev. D 84, 086008 – Published 25 October 2011 Tom Banks and John Kehayias.**

Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect

- (3) **Chaolun Wu, Shao-Feng Wu** show that Horava-Lifshitz gravity theory can be employed (e) as a covariant framework to build (eb) an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry.
- (4) The key to this formalism is a set of correspondence relations that maps all the field degrees of freedom in the Horava-Lifshitz gravity theory to (e&eb) external background (source) fields among others in the effective action of the quantum Hall effect, according to their symmetry transformation properties.

NOTATION

Module One

The strong holographic principle requires (e) the space of these sections to be (=) finite dimensional
 G_{13} : Category one of space of these sections to be (=) finite dimensional
 G_{14} : Category two of SAS(same as superior/above)
 G_{15} : Category three of SAS
 T_{13} : Category one of strong holographic principle

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

strong holographic principle requires the space of these sections to be (=) finite dimensional

G_{16} : Category one of strong holographic principle requires the space of these sections

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of finite dimensional

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

They discuss applications of
 fuzzy spinor geometry to (e&eb) holographic space-time and to matrix theory.

DOI: <http://dx.doi.org/10.1103/PhysRevD.84.086008> © 2011 American Physical Society **Fuzzy geometry via the spinor bundle, with applications to holographic space-time and matrix theory Phys. Rev. D 84, 086008 – Published 25 October 2011 Tom Banks and John Kehayias.**

G_{20} : Category one of fuzzy spinor geometry; holographic space-time

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of fuzzy spinor geometry; holographic space-time

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

G_{24} : Category one of fuzzy spinor geometry; matrix theory.

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of matrix theory.; fuzzy spinor geometry

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Chaolun Wu, Shao-Feng Wu show that

Horava-Lifshitz gravity theory can be employed (e) as a covariant framework to build (eb) an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

G_{28} : Category one of covariant framework to build (eb) an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Horava-Lifshitz gravity theory

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Horava-Lifshitz gravity theory can be employed as a covariant framework to build (eb) an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

G_{32} : Category one of Horava-Lifshitz gravity theory can be employed as a covariant framework

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

G_{36} : Category one of Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

G_{40} : Category one of non-relativistic diffeomorphism invariance; anisotropic Weyl invariance as well as the gauge symmetry

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of anisotropic Weyl invariance as well as the gauge symmetry ;non-relativistic diffeomorphism invariance

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

non-relativistic diffeomorphism invariance and anisotropic Weyl invariance as well as the gauge symmetry

G_{44} : Category one of non-relativistic diffeomorphism invariance; gauge symmetry

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of gauge symmetry ;non-relativistic diffeomorphism invariance

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$

$$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$$

are Accentuation coefficients

$$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)},$$

$$(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$$

$$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$$

$$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$$

$$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$$

$$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$$

$$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad 6$$

$$+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor}$$

$$-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor}$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 56$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 57$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for

1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for

category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a''_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a''_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a''_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{c} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) - (b''_{13})^{(1,1)}(G, t) - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19}, t) & -(b''_{14})^{(1,1)}(G, t) & -(b''_{21})^{(3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ & -(b''_{37})^{(7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19}, t) & -(b''_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ & -(b''_{38})^{(7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b'_{36})^{(7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b'_{40})^{(8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b'_{44})^{(9,9)}(G_{47}, t)$, $-(b'_{46})^{(9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21}, t) & +(a''_{16})^{(2,2,2)}(T_{17}, t) & +(a''_{13})^{(1,1,1)}(T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ & +(a''_{36})^{(7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21}, t) & +(a''_{17})^{(2,2,2)}(T_{17}, t) & +(a''_{14})^{(1,1,1)}(T_{14}, t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ & +(a''_{37})^{(7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21}, t) & +(a''_{18})^{(2,2,2)}(T_{17}, t) & +(a''_{15})^{(1,1,1)}(T_{14}, t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ & +(a''_{38})^{(7,7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)}(T_{21}, t)$, $+(a'_{21})^{(3)}(T_{21}, t)$, $+(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$ are third augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$ are eight augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{20})^{(3)}(G_{23}, t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} \boxed{(b'_{21})^{(3)}(G_{23}, t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{22})^{(3)}(G_{23}, t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25}, t) & +(a''_{28})^{(5,5)}(T_{29}, t) & +(a''_{32})^{(6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25}, t) & +(a''_{29})^{(5,5)}(T_{29}, t) & +(a''_{33})^{(6,6)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25}, t) & +(a''_{30})^{(5,5)}(T_{29}, t) & +(a''_{34})^{(6,6)}(T_{33}, t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14}, t) & +(a''_{18})^{(2,2,2,2)}(T_{17}, t) & +(a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{l} (b'_{28})^{(5)}[-(b''_{28})^{(5)}(G_{31}, t) \quad -(b''_{24})^{(4,4)}(G_{27}, t) \quad -(b''_{32})^{(6,6,6)}(G_{35}, t)] \\ -(b''_{13})^{(1,1,1,1,1)}(G, t) \quad -(b''_{16})^{(2,2,2,2,2)}(G_{19}, t) \quad -(b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) \quad -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) \quad -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{l} (b'_{29})^{(5)}[-(b''_{29})^{(5)}(G_{31}, t) \quad -(b''_{25})^{(4,4)}(G_{27}, t) \quad -(b''_{33})^{(6,6,6)}(G_{35}, t)] \\ -(b''_{14})^{(1,1,1,1,1)}(G, t) \quad -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t) \quad -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) \quad -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) \quad -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{l} (b'_{30})^{(5)}[-(b''_{30})^{(5)}(G_{31}, t) \quad -(b''_{26})^{(4,4)}(G_{27}, t) \quad -(b''_{34})^{(6,6,6)}(G_{35}, t)] \\ -(b''_{15})^{(1,1,1,1,1)}(G, t) \quad -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t) \quad -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \quad -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) \quad -(b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & -(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & -(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & -(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(MMMM) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(NNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(OOOO) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(PPPP)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(QQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(RRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(p) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t}$$

376

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq$$

377

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(p) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b_{20}')^{(3)}(b_{21}')^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b_{20}')^{(3)}(b_{21}'')^{(3)}(G_{23}) + (b_{21}')^{(3)}(b_{20}'')^{(3)}(G_{23})] + (b_{20}'')^{(3)}(G_{23})(b_{21}'')^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b_{24}')^{(4)}(b_{25}')^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

$$\text{ASYMPTOTIC STABILITY ANALYSIS} \quad 531$$

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*\mathbb{G}_j \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*\mathbb{G}_j \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*\mathbb{G}_j \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*\mathbb{G}_j \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*\mathbb{G}_j \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*\mathbb{G}_j \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVENTEEN

Horava-Lifshitz Gravity

INTRODUCTION—VARIABLES USED

Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect

- (1) Chaolun Wu, Shao-Feng Wu originally derive the **map as (=) a holographic dictionary**, but its form is independent of (e) the existence of holographic duality.
- (2) This paves the way for the application of Horava-Lifshitz holography on (e&eb) fractional quantum Hall effect.

Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect

- (3) Using (e) the simplest holographic Chern-Simons model, authors compute (e&eb) the low energy effective action at leading orders and show that (eb) it captures universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations.
- (4) They identify the shift function in Horava-Lifshitz gravity theory as (=) minus of guiding center velocity and conjugate to guiding center momentum.
- (5) This enables us to distinguish guiding center angular momentum density from (e&eb) the internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles.
- (6) Effective action shows that Hall viscosity is (=) minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity is (=) half of the guiding center angular momentum density. ArXiv: 1409.1178 [hep-th] **Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect**

NOTATION

Module One

Chaolun Wu, Shao-Feng Wu originally derive the

map as (=) a holographic dictionary, but its form is independent of (e) the existence of holographic duality

G_{13} : Category one o **map**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **holographic dictionary**, but its form is independent of (e) the existence of holographic duality

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

map as a holographic dictionary, but its form is independent of (e) the existence of holographic duality

G_{16} : Category one of existence of holographic duality

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of **map as a holographic dictionary**, but its form is independent

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

This paves the way for the application of

Horava-Lifshitz holography on (e&eb) fractional quantum Hall effect

G_{20} : Category one of Horava-Lifshitz holography; fractional quantum Hall effect

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of fractional quantum Hall effect; Horava-Lifshitz holography

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Using (e) the simplest holographic Chern-Simons model, authors compute (e&eb) the low energy effective action at leading orders and show that (eb) it captures universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

G_{24} : Category one of simplest holographic Chern-Simons model

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of low energy effective action at leading orders and show that (eb) it captures universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Using the

simplest holographic Chern-Simons model, authors compute the low energy effective action at leading orders and show that (eb) model captures universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

G_{28} : Category one of simplest holographic Chern-Simons model, authors compute the low energy effective action(s)

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of model captures universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

simplest holographic Chern-Simons model, authors compute the low energy effective action at leading orders and show that model captures (e) universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

G_{32} : Category one of universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of simplest holographic Chern-Simons model, authors compute the low energy effective action at leading orders and show that model

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

simplest holographic Chern-Simons model, authors compute the low energy effective action at leading orders and show that model captures universal electromagnetic and geometric properties of Quantum Hall States, including (e) the Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

G_{36} : Category one of Wen-Zee shift (e&eb) Hall viscosity (e&eb) angular momentum density and their relations

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of simplest holographic Chern-Simons model, authors compute the low energy effective action at leading orders and show that model captures universal electromagnetic and geometric properties of Quantum Hall States

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

simplest holographic Chern-Simons model, authors compute the low energy effective action at leading orders and show that model captures universal electromagnetic and geometric properties of Quantum Hall States, including the Wen-Zee shift (e&eb) Hall viscosity angular momentum density and their relations

G_{40} : Category one of Wen-Zee shift; Hall viscosity angular momentum density

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Hall viscosity angular momentum density ;Wen-Zee shift

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

They identify the

shift function in Horava-Lifshitz gravity theory as (=) minus of guiding center velocity and conjugate to guiding center momentum

G_{44} : Category one of shift function in Horava-Lifshitz gravity theory

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of minus of guiding center velocity and conjugate to guiding center momentum

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)},$
 $(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &- (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &- (b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20\end{aligned}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & - (b''_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b''_{22})^{(3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{38})^{(7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) + (a''_{24})^{(4,4)}(T_{25}, t) + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}' - G_{31}|| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(SSSS) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(TTTT) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(UUUU) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(VVVV)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(WWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(XXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(q) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})_1$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{(\bar{M}_{16})^{(2)} s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{-(\bar{M}_{16})^{(2)} s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(9)}$ and $(b''_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t) :-$

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t) :-$

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484 A
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(q) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42}(s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION EIGHTEEN

Non-Perturbative Lorentzian Quantum Gravity

INTRODUCTION—VARIABLES USED

Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect

- (1) This shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density from (e&eb) the internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles.
- (2) Effective action shows that Hall viscosity is (=) minus half of the internal angular momentum

density and proportional to Wen-Zee shift, and Hall bulk viscosity is (=) half of the guiding center angular momentum density. ArXiv: 1409.1178 [hep-th] **Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect**

Nuclear Physics B Volume 536, Issues 1–2, 21 December 1998, Pages 407–434 Non-perturbative Lorentzian quantum gravity, causality and topology change J. Ambjørn, R. Loll doi:10.1016/S0550-3213(98)00692-0

- (3) Authors formulate a non-perturbative lattice model of (e) two-dimensional Lorentzian quantum gravity by performing (e&eb) the path integral over geometries with a causal structure.
- (4) The model can be solved exactly at (eb) the discretized level.
- (5) Its continuum limit coincides (=) with the theory obtained by (e) quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory.
- (6) By allowing topology change of the compact spatial slices (i.e. baby universe creation), one obtains (eb) agreement with the matrix models and Liouville theory

Supersymmetric Mechanics - Vol. 3 Lecture Notes in Physics Volume 755, 2008, pp 1-92 Black Holes, Black Rings, and their Microstates Iosif Bena, Nicholas P. Warner

In this review article, we describe some of the recent progress towards the construction and analysis of three-charge configurations in string theory and supergravity. We begin by describing the Born-Infeld construction of three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions in five dimensions. We explain in detail the use of these methods to construct black rings, black holes, as well as smooth microstate geometries with black hole and black ring charges, but with no horizon. We present arguments that many of these microstate geometries are dual to boundary states that belong to the same sector of the D1-D5-P CFT as the typical states. We end with an extended discussion of the implications of this work for the physics of black holes in string theory.

NOTATION

Module One

shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables (eb) us to distinguish guiding center angular momentum density from (e&eb) the internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

G_{13} : Category one of shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of distinguish guiding center angular momentum density from (e&eb) the internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding

center momentum enables us to distinguish guiding center angular momentum density from (e&eb) the internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

G_{16} : Category one of shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density; internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles; shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density from the internal one, which is (=) the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

G_{20} : Category one of shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density from the internal one

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density from the internal one, which is the sum of Landau orbit spin and (e&eb) intrinsic (topological) spin of the composite particles

G_{24} : Category one of shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density from the internal one, which is the sum of Landau orbit spin; intrinsic (topological) spin of the composite particles

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of intrinsic (topological) spin of the composite particles; shift function in Horava-Lifshitz gravity theory as minus of guiding center velocity and conjugate to guiding center momentum enables us to distinguish guiding center angular momentum density from the internal one, which is the sum of Landau orbit spin

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Effective action shows that Hall viscosity is (=) minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity is (=) half of the guiding center angular momentum density.

ArXiv: 1409.1178 [hep-th] **Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect**

G_{28} : Category one of Effective action shows that Hall viscosity

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity is (=) half of the guiding center angular momentum density.

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Effective action shows that Hall viscosity is minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity is (=) half of the guiding center angular momentum density.

G_{32} : Category one of Effective action shows that Hall viscosity is minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of half of the guiding center angular momentum density.

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Authors formulate a

non-perturbative lattice model of (e) two-dimensional Lorentzian quantum gravity by performing (e&eb) the path integral over geometries with a causal structure

G_{36} : Category one of two-dimensional Lorentzian quantum gravity by performing (e&eb) the path integral over geometries with a causal structure

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of non-perturbative lattice model

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

non-perturbative lattice model of two-dimensional Lorentzian quantum gravity by performing (e&eb) the path integral over geometries with a causal structure

G_{40} : Category one of non-perturbative lattice model of two-dimensional Lorentzian quantum gravity; path integral over geometries with a causal structure

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of path integral over geometries with a causal structure; non-perturbative lattice model of two-dimensional Lorentzian quantum gravity

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

non-perturbative lattice model of two-dimensional Lorentzian quantum gravity by performing the path integral over geometries with a causal structure

G_{44} : Category one of non-perturbative lattice model of two-dimensional Lorentzian quantum gravity by performing the path integral; geometries with a causal structure

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of geometries with a causal structure; non-perturbative lattice model of two-dimensional Lorentzian quantum gravity by performing the path integral

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \end{aligned}$$

$$\begin{aligned}\frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

12

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$- (b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+ (a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$- (b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+ (a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$- (b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\begin{aligned}\frac{dG_{32}}{dt} &= (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} & 31 \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} & 32 \\ \frac{dG_{34}}{dt} &= (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} & 33 \\ \frac{dT_{32}}{dt} &= (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} & 34 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} & 35 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} & 36 \\ &+ (a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor} \\ &\text{Module Numbered Seven:}\end{aligned}$$

The differential system of this model is now (Seventh Module)

$$\begin{aligned}\frac{dG_{36}}{dt} &= (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} & 42 \\ &+ (a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \\ &\text{Module Numbered Eight}\end{aligned}$$

The differential system of this model is now

$$\begin{aligned}\frac{dG_{40}}{dt} &= (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} & 44 \\ \frac{dG_{42}}{dt} &= (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} & 54 \\ &+ (a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor} \\ &- (b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}\end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a'_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a'_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a'_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{c} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients

for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for

category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients

for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition

coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition

coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31} - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(YYYY) (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(ZZZZ) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(AAAAA) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(BBBBB)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(CCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(DDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$, Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(r) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right] \quad 181$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is 188
analogous with the preceding one. An analogous property is true if G_{14} is bounded from below.

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}, i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{-(\hat{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)}t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(\mu_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} [e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(r) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)})$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)})$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION NINETEEN

Non-Perturbative Lorentzian Quantum Gravity

INTRODUCTION—VARIABLES USED

Nuclear Physics B Volume 536, Issues 1–2, 21 December 1998, Pages 407–434 Non-perturbative Lorentzian quantum gravity, causality and topology change J. Ambjørn, R. Loll doi:10.1016/S0550-3213(98)00692-0

- (1) The model can be solved exactly at (eb) the discretized level.
- (2) Its continuum limit coincides (=) with the theory obtained by (e) quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory.
- (3) By allowing topology change of the compact spatial slices (i.e. baby universe creation), one obtains (eb) agreement with the matrix models and Liouville theory

Supersymmetric Mechanics - Vol. 3 Lecture Notes in Physics Volume 755, 2008, pp 1-92 Black Holes, Black Rings, and their Microstates Iosif Bena, Nicholas P. Warner

- (4) In this review article, authors describe some of the recent progress towards the construction and analysis of three-charge configurations in (eb) string theory and supergravity.
- (5) Authors begin by describing the Born-Infeld construction of (e) three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions in (eb) five dimensions.
- (6) They explain in detail the use of these methods to construct (eb) black rings, black holes, as well as smooth microstate geometries with (e&eb) black hole and black ring charges, but with no horizon.
- (7) They present arguments that many of these microstate geometries are (=) dual to boundary states that belong (eb) to the same sector of the D1-D5-P CFT as the typical states.
- (8) Paper ends with an extended discussion of the implications of this work for the (e&eb) physics of black holes in string theory.

NOTATION

Module One

Model can be solved exactly at (eb) the discretized level.

G_{13} : Category one of model can be solved exactly

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of discretized level

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Its continuum limit coincides (=) with the theory obtained by (e) quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory

G_{16} : Category one of continuum limit

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of theory obtained by (e) quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Its continuum limit coincides with the theory obtained by (e) quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory

G_{20} : Category one of quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of continuum limit coincides with the theory obtained

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Its

continuum limit coincides with the theory obtained by quantizing 2d continuum gravity in (eb) proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory

G_{24} : Category one of continuum limit coincides with the theory obtained by quantizing 2d continuum gravity

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of proper-time gauge, but it disagrees with (e&eb) 2d gravity defined via matrix models or Liouville theory

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

continuum limit coincides with the theory obtained by quantizing 2d continuum gravity in proper-time gauge, but it disagrees with 2d gravity defined via matrix models or Liouville theory

G_{28} : Category one of continuum limit coincides with the theory obtained by quantizing 2d continuum gravity in proper-time; 2d gravity defined via matrix models or Liouville theory

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of 2d gravity defined via matrix models or Liouville theory; continuum limit coincides with the theory obtained by quantizing 2d continuum gravity in proper-time

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

By allowing topology change of the compact spatial slices (i.e. baby universe creation), one obtains (eb) agreement with the matrix models and Liouville theory

G_{32} : Category one of allowing topology change of the compact spatial slices (i.e. baby universe creation)

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of agreement with the matrix models and Liouville theory

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

G_{36} : Category one of matrix models

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Liouville theory

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

In this review article, authors describe some of the recent progress towards the construction and analysis of three-charge configurations in (eb) string theory and supergravity

G_{40} : Category one of construction and analysis of three-charge configurations

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of string theory and supergravity

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Authors begin by describing the

Born-Infeld constructions of three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions in (eb) five dimensions.

G_{44} : Category one of three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions in (eb) five dimensions

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Born-Infeld constructions

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$$\begin{aligned} & (a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)} \\ & , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)} \\ & (a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, \\ & (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)} \\ & (a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}, \\ & (a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}, \\ & (a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}, \end{aligned}$$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ & + (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ & - (b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\ & + (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ & - (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} & 16 \end{aligned}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 56$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 57$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for

1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for

category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1)}(G, t)} & \boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1)}(G, t)} & \boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{46})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a'_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a'_{21})^{(3)}(T_{21}, t)} & \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} \boxed{+(a'_{22})^{(3)}(T_{21}, t)} & \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \quad 69$$

$\boxed{+(a'_{20})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3)}(T_{21}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)}$ are second augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$ are third augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$ are eight augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{20})^{(3)}(G_{23}, t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} \boxed{(b'_{21})^{(3)}(G_{23}, t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{22})^{(3)}(G_{23}, t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(EEEE) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(FFFF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(GGGGG) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(HHHHH)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(IIII) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(JJJJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0} \right)} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\left(\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0} \right)} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(s) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{(\bar{M}_{16})^{(2)} s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{-(\bar{M}_{16})^{(2)} s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}\} (T_{25}(s_{(24)}), s_{(24)}) ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t})$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t) :-$

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t) :-$

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)}$ and $(b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(s) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b_{20}')^{(3)}(b_{21}')^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b_{20}')^{(3)}(b_{21}'')^{(3)}(G_{23}) + (b_{21}')^{(3)}(b_{20}'')^{(3)}(G_{23})] + (b_{20}'')^{(3)}(G_{23})(b_{21}'')^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b_{24}')^{(4)}(b_{25}')^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* \mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* \mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b_i'')^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* \mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^* \mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^* \mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)} T_{28}^* \mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)} T_{29}^* \mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)} T_{30}^* \mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)} \right) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY

Entanglement Renormalization Vis A Vis Holographic Principle

INTRODUCTION—VARIABLES USED

Supersymmetric Mechanics - Vol. 3 Lecture Notes in Physics Volume 755, 2008, pp 1-92 Black Holes, Black Rings, and their Microstates Iosif Bena, Nicholas P. Warner

- (1) They explain in detail the use of these methods to construct black rings, black holes, as well as smooth microstate geometries with (e&eb) black hole and black ring charges, but with no horizon.
- (2) They present arguments that many of these microstate geometries are (=) dual to boundary states that belong (eb) to the same sector of the D1-D5-P CFT as the typical states.

- (3) Paper ends with an extended discussion of the implications of this work for the (e&eb) physics of black holes in string theory.

**Phys. Rev. D 86, 065007 (2012) DOI: 10.1103/PhysRevD.86.065007 arXiv: 0905.1317 [cond-mat.str-el]
 Entanglement Renormalization and Holography Brian Swingle**

- (1) **Chaolun Wu, Shao-Feng Wu. Brian Swingle** shows how recent progress in real space renormalization group methods can be used to define (eb) a generalized notion of holography inspired by **holographic dualities in quantum gravity**.
- (2) The generalization is based upon organizing information in (eb) a quantum state in terms of scale and defining a higher dimensional geometry from (e) this structure.
- (3) While states with a finite correlation length typically give simple geometries, the state at a quantum critical point gives (eb) a discrete version of anti de Sitter space.
- (4) Some finite temperature quantum states include (e) black hole-like objects.
- (5) The gross features of equal time correlation functions are (=) also reproduced in this geometric framework. The relationship between this framework and better understood versions of holography is discussed: Phys. Rev. D 86, 065007 (2012) DOI: 10.1103/PhysRevD.86.065007 arXiv: 0905.1317 [cond-mat.str-el] **Entanglement Renormalization and Holography Brian Swingle**

NOTATION

Module One

Born-Infeld constructions of three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions in (eb) five dimensions

G_{13} : Category one of Born-Infeld constructions of three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of five dimensions

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

They explain in detail the use of these methods to construct black rings, black holes, as well as smooth microstate geometries with (e&eb) black hole and black ring charges, but with no horizon

G_{16} : Category one of Born-Infeld constructions of three-charge supertubes with two dipole charges and then discuss the general method of constructing three-charge solutions in five dimensions

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of black rings, black holes, as well as smooth microstate geometries with (e&eb) black hole and black ring charges, but with no horizon

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

black rings, black holes, as well as smooth microstate geometries with (e&eb) black hole and black ring charges, but with no horizon

G_{20} : Category one of black rings, black holes, as well as smooth microstate geometries; black hole and black ring charges, but with no horizon

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of black hole and black ring charges, but with no horizon; black rings, black holes, as well as smooth microstate geometries

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

They present arguments that

many of these microstate geometries are (=) dual to boundary states that belong (eb) to the same sector of the D1-D5-P CFT as the typical states

G_{24} : Category one of many of these microstate geometries

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of dual to boundary states that belong (eb) to the same sector of the D1-D5-P CFT as the typical states

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

many of these microstate geometries are dual to boundary states that belong to the same sector of the D1-D5-P CFT as the typical states

G_{28} : Category one of many of these microstate geometries are dual to boundary states

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of same sector of the D1-D5-P CFT as the typical states

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Paper ends with an extended discussion of the

implications of this work for the (e&eb) physics of black holes in string theory

G_{32} : Category one of many of these microstate geometries are dual to boundary states that belong to the same sector of the D1-D5-P CFT as the typical states; physics of black holes in string theory

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of physics of black holes in string theory; many of these microstate geometries are dual to boundary states that belong to the same sector of the D1-D5-P CFT as the typical states

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Chaolun Wu, Shao-Feng Wu. Brian Swingle shows how recent progress in real space renormalization group methods can be used to define a generalized notion of holography inspired by **holographic dualities in quantum gravity**.

G_{36} : Category one of real space renormalization group methods; define a generalized notion of holography inspired by **holographic dualities in quantum gravity**.

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of define a generalized notion of holography inspired by **holographic dualities in quantum gravity**.; real space renormalization group methods

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

The

generalization is based upon organizing information in (eb) a quantum state in terms of scale and defining a higher dimensional geometry from (e) this structure

G_{40} : Category one of generalization is based upon organizing information

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of quantum state in terms of scale and defining a higher dimensional geometry from (e) this structure

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

generalization is based upon organizing information in a quantum state in terms of scale and defining a higher dimensional geometry from (e) this structure

G_{44} : Category one of structure (there may be more structure for different experimentation; alternatively all the three categories can be taken as being equivalent)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of generalization is based upon organizing information in a quantum state in terms of scale and defining a higher dimensional geometry

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13}$$

$$\begin{aligned}\frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20 \\ \frac{dG_{26}}{dt} &= (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} & 21 \\ \frac{dT_{24}}{dt} &= (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} & 22 \\ \frac{dT_{25}}{dt} &= (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} & 23 \\ \frac{dT_{26}}{dt} &= (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} & 24\end{aligned}$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$, $(a''_{21})^{(3,3)}(T_{21}, t)$, $(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7)}(T_{37}, t)$, $(a''_{37})^{(7,7)}(T_{37}, t)$, $(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8)}(T_{41}, t)$, $(a''_{41})^{(8,8)}(T_{41}, t)$, $(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{c} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $[-(b''_{13})^{(1)}(G, t)]$, $[-(b''_{14})^{(1)}(G, t)]$, $[-(b''_{15})^{(1)}(G, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{16})^{(2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{20})^{(3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3)}(G_{23}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b''_{28})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b''_{29})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b''_{30})^{(5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b''_{36})^{(7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b''_{40})^{(8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8)}(G_{43}, t)]$, $[-(b''_{42})^{(8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17}, t)] \quad [+(a''_{13})^{(1,1)}(T_{14}, t)] \quad [+(a''_{20})^{(3,3,3)}(T_{21}, t)] \\ [+(a''_{24})^{(4,4,4,4)}(T_{25}, t)] \quad [+(a''_{28})^{(5,5,5,5)}(T_{29}, t)] \quad [+(a''_{32})^{(6,6,6,6)}(T_{33}, t)] \\ [+(a''_{36})^{(7,7,7)}(T_{37}, t)] \quad [+(a''_{40})^{(8,8,8)}(T_{41}, t)] \quad [+(a''_{44})^{(9,9)}(T_{45}, t)] \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}[+(a''_{17})^{(2)}(T_{17}, t)] \quad [+(a''_{14})^{(1,1)}(T_{14}, t)] \quad [+(a''_{21})^{(3,3,3)}(T_{21}, t)] \\ [+(a''_{25})^{(4,4,4,4)}(T_{25}, t)] \quad [+(a''_{29})^{(5,5,5,5)}(T_{29}, t)] \quad [+(a''_{33})^{(6,6,6,6)}(T_{33}, t)] \\ [+(a''_{37})^{(7,7,7)}(T_{37}, t)] \quad [+(a''_{41})^{(8,8,8)}(T_{41}, t)] \quad [+(a''_{45})^{(9,9)}(T_{45}, t)] \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b'_{20})^{(3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b'_{21})^{(3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{37})^{(7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b'_{22})^{(3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{38})^{(7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients

for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients

for category 1,2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for

category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for

category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category

1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients

for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients

for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficients for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation

coefficients for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation

coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients

for category 1, 2 and 3

$+(a'_{44})^{(9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23}, t) & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23}, t) & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23}, t) & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25}, t) & +(a''_{28})^{(5,5)}(T_{29}, t) & +(a''_{32})^{(6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25}, t) & +(a''_{29})^{(5,5)}(T_{29}, t) & +(a''_{33})^{(6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{c} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t), (a''_{25})^{(4)}(T_{25}, t), (a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t), +(a''_{29})^{(5,5)}(T_{29}, t), +(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t), +(a''_{33})^{(6,6)}(T_{33}, t), +(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{c} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) - (b''_{28})^{(5,5)}(G_{31}, t) - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{c} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) - (b''_{29})^{(5,5)}(G_{31}, t) - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{c} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) - (b''_{30})^{(5,5)}(G_{31}, t) - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a'_{24})^{(4,4)}(T_{25}, t) & + (a'_{32})^{(6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a'_{25})^{(4,4)}(T_{25}, t) & + (a'_{33})^{(6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a'_{26})^{(4,4)}(T_{25}, t) & + (a'_{34})^{(6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a'_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a'_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a'_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}[-(b''_{28})^{(5)}(G_{31}, t)] & -(b''_{24})^{(4,4)}(G_{27}, t) & -(b''_{32})^{(6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}[-(b''_{29})^{(5)}(G_{31}, t)] & -(b''_{25})^{(4,4)}(G_{27}, t) & -(b''_{33})^{(6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}[-(b''_{30})^{(5)}(G_{31}, t)] & -(b''_{26})^{(4,4)}(G_{27}, t) & -(b''_{34})^{(6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)}[+(a''_{32})^{(6)}(T_{33}, t)] & +(a''_{28})^{(5,5,5)}(T_{29}, t) & +(a''_{24})^{(4,4,4)}(T_{25}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)}[+(a''_{33})^{(6)}(T_{33}, t)] & +(a''_{29})^{(5,5,5)}(T_{29}, t) & +(a''_{25})^{(4,4,4)}(T_{25}, t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients

for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients

for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients

for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition

coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition

coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{36})^{(7)}(T_{37}, t)$, $(a''_{37})^{(7)}(T_{37}, t)$, $(a''_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) & + (a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)} + (a_{41}'')^{(8)}(T_{41}, t) & + (a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)} + (a_{42}'')^{(8)}(T_{41}, t) & + (a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41}, t)$, $+(a_{41}'')^{(8)}(T_{41}, t)$, $+(a_{42}'')^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

96

$$= (a_{44})^{(9)}G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}[+(a''_{45})^{(9)}(T_{45}, t)] & +(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & +(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}[+(a''_{46})^{(9)}(T_{37}, t)] & +(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & +(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}[-(b''_{44})^{(9)}(G_{47}, t)] & -(b'_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t) & -(b'_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) \quad - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) \quad - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47}, t)$, $-(b''_{45})^{(9)}(G_{47}, t)$, $-(b''_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b'_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a'_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b'_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition:

99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$:

106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition:

114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21}' - T_{21}| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23}' - G_{23}| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$:

115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26$$

117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

118

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition:

119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$:

120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30$$

122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

123

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

128

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33}' - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(KKKKK) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(LLLLL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(MMMMM) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(NNNNN)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ are positive constants and $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37}' - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(00000) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(PPPPPP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T'_{45}, t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47}) - (G_{47})'| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T'_{45}, t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(t) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{((\hat{P}_{13})^{(1)} + G_j^0)}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{((\hat{Q}_{13})^{(1)} + T_j^0)}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) &d((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}\} (T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $(\bar{M}_{44})^{(9)}_1, (\bar{M}_{44})^{(9)}_2$ and $(\bar{M}_{44})^{(9)}_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t})$$

$$T_{40}^0e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)}) - (S_2)^{(9)}} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 409

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(t) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b'_{14})^{(1)}(G) + (b'_{14})^{(1)}(b'_{13})^{(1)}(G)] + (b'_{13})^{(1)}(G)(b'_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b'_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b'_{16})^{(2)}(G_{19})] + (b'_{16})^{(2)}(G_{19})(b'_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b'_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b'_{20})^{(3)}(G_{23})] + (b'_{20})^{(3)}(G_{23})(b'_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}(G_{43}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)} \right) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY ONE

Topological Entanglement Entropy

INTRODUCTION—VARIABLES USED

Phys. Rev. D 86, 065007 (2012) DOI: 10.1103/PhysRevD.86.065007 arXiv: 0905.1317 [cond-mat.str-el]
Entanglement Renormalization and Holography Brian Swingle

- (1) Paul Fendley, 1 Matthew P. A. Fisher² and Chetan Nayak³, 4study the entropy of (e) chiral 2+1-dimensional topological phases, where there are (=) both gapped bulk excitations and gapless edge modes.
- (2) They show how the entanglement entropy of both types of excitations can be encoded in (eb) a

single partition function.

- (3) This partition function is holographic because (e) it can be expressed entirely in terms of the conformal field theory describing the edge modes.
- (4) They give a general expression for the holographic partition function, and discuss several examples in depth, including abelian and non-abelian fractional Quantum Hall States, and (e&eb) p + ip superconductors.
- (5) They extend these results to include a point contact allowing (eb) tunneling between two points on the edge, which causes (eb) thermodynamic entropy associated with (e&eb) the point contact to be lost with decreasing temperature.
- (6) Such a perturbation effectively breaks (e) the system in two, and we can identify (e&eb) the thermodynamic entropy loss with the loss of the edge entanglement entropy.
- (7) From these results, authors obtain a simple interpretation of (e) the non-integer ‘ground state degeneracy’ which is (=) obtained in 1+1-dimensional quantum impurity problems: its logarithm is (=) a 2+1-dimensional topological entanglement entropy **Topological Entanglement Entropy from the Holographic Partition Function**
- (8) **Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}. John Preskill** considers some promising future directions for quantum information theory that could influence the development of 21st century physics. Advances in the theory of the distinguishability of superoperators may lead to (eb) new strategies for improving the precision of quantum-limited measurements.
- (9) A better grasp of the properties of multi-partite quantum entanglement may lead to (eb) deeper understanding of (e) strongly-coupled dynamics in (eb) quantum many-body systems, quantum field theory, and quantum gravity.

NOTATION

Module One

Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}, 4study the

entropy of (e) chiral 2+1-dimensional topological phases, where there are (=) both gapped bulk excitations and gapless edge modes

G_{13} : Category one of chiral 2+1-dimensional topological phases, where there are (=) both gapped bulk excitations and gapless edge modes

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of entropy

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

entropy of chiral 2+1-dimensional topological phases, where there are (=) both gapped bulk excitations and gapless edge modes

G_{16} : Category one of entropy of chiral 2+1-dimensional topological phases,

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of both gapped bulk excitations and gapless edge modes

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

They show how the entanglement entropy of both types of excitations can be encoded in (eb) a single partition function
 G_{20} : Category one of entanglement entropy of both types of excitations can be encoded

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of single partition function(systemic differentiation, or alternately take all the three stratifications as equivalent)

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Partition function is holographic because (e) it can be expressed entirely in terms of the conformal field theory describing the edge modes
 G_{24} : Category one of expression entirely in terms of the conformal field theory describing the edge modes

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Partition function is holographic

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Partition function is holographic because it can be expressed entirely in terms of the conformal field theory describing the edge modes

G_{28} : Category one of Partition function is holographic because it can be expressed; conformal field theory describing the edge modes

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of conformal field theory describing the edge modes; Partition function is holographic because it can be expressed

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

They give a general expression for the holographic partition function, and discuss several examples in depth, including

abelian and non-abelian fractional Quantum Hall States, and $p + ip$ superconductors

G_{32} : Category one of abelian and non-abelian fractional Quantum Hall States; $p + ip$ superconductors

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of $p + ip$ superconductors ;abelian and non-abelian fractional Quantum Hall States

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

They

Extension of these results to include a point contact allowing $(e\&e_b)$ tunneling between two points on the edge, which causes (e_b) thermodynamic entropy associated with $(e\&e_b)$ the point contact to be lost with decreasing temperature

G_{36} : Category one of Extension of these results to include a point contact

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of tunneling between two points on the edge, which causes (e_b) thermodynamic entropy associated with $(e\&e_b)$ the point contact to be lost with decreasing temperature

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Extension of these results to include a point contact allowing tunneling between two points on the edge, which causes (e_b) thermodynamic entropy associated with $(e\&e_b)$ the point contact to be lost with decreasing temperature

G_{40} : Category one of Extension of these results to include a point contact allowing tunneling between two points on the edge

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of thermodynamic entropy associated with (e&eb) the point contact to be lost with decreasing temperature

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Extension of these results to include a point contact allowing tunneling between two points on the edge, which causes thermodynamic entropy associated with (e&eb) the point contact to be lost with decreasing temperature

G_{44} : Category one of Extension of these results to include a point contact allowing tunneling between two points on the edge, which causes thermodynamic entropy; point contact to be lost with decreasing temperature

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of point contact to be lost with decreasing temperature ;Extension of these results to include a point contact allowing tunneling between two points on the edge, which causes thermodynamic entropy

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &- (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &- (b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

42

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40}$$

43

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41}$$

44

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42}$$

45

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40}$$

46

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41}$$

47

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42}$$

48

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44}$$

49

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45}$$

50

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46}$$

51

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44}$$

52

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45}$$

53

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

55

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

56

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

57

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$, $(a''_{21})^{(3,3)}(T_{21}, t)$, $(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient

for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient

for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) \quad - (b''_{16})^{(2,2)}(G_{19}, t) \quad - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{28})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) \quad - (b''_{40})^{(8,8)}(G_{43}, t) \quad - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) \quad - (b''_{17})^{(2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8)}(G_{43}, t) \quad - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) \quad - (b''_{18})^{(2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8)}(G_{43}, t) \quad - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} (G_{19}, t) & -(b''_{15})^{(1,1)} (G, t) & -(b''_{22})^{(3,3,3)} (G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)} (G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)} (G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)} (G_{35}, t) \\ -(b''_{38})^{(7,7,7)} (G_{39}, t) & -(b''_{42})^{(8,8,8)} (G_{43}, t) & -(b''_{46})^{(9,9)} (G_{47}, t) \end{array} \right] T_{18}$$

where $-(b'_{16})^{(2)} (G_{19}, t)$, $-(b'_{17})^{(2)} (G_{19}, t)$, $-(b'_{18})^{(2)} (G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)} (G, t)$, $-(b'_{14})^{(1,1)} (G, t)$, $-(b'_{15})^{(1,1)} (G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)} (G_{23}, t)$, $-(b''_{21})^{(3,3,3)} (G_{23}, t)$, $-(b''_{22})^{(3,3,3)} (G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)} (G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)} (G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)} (G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)} (G_{39}, t)$, $-(b''_{37})^{(7,7,7)} (G_{39}, t)$, $-(b''_{38})^{(7,7,7)} (G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)} (G_{43}, t)$, $-(b''_{41})^{(8,8,8)} (G_{43}, t)$, $-(b''_{42})^{(8,8,8)} (G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)} (G_{47}, t)$, $-(b''_{46})^{(9,9)} (G_{47}, t)$, $-(b''_{45})^{(9,9)} (G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} (T_{21}, t) & +(a''_{16})^{(2,2,2)} (T_{17}, t) & +(a'_{13})^{(1,1,1)} (T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a''_{36})^{(7,7,7,7)} (T_{37}, t) & +(a''_{40})^{(8,8,8,8)} (T_{41}, t) & +(a''_{44})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} (T_{21}, t) & +(a''_{17})^{(2,2,2)} (T_{17}, t) & +(a'_{14})^{(1,1,1)} (T_{14}, t) \\ +(a''_{25})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a''_{37})^{(7,7,7,7)} (T_{37}, t) & +(a''_{41})^{(8,8,8,8)} (T_{41}, t) & +(a''_{45})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} (T_{21}, t) & +(a''_{18})^{(2,2,2)} (T_{17}, t) & +(a'_{15})^{(1,1,1)} (T_{14}, t) \\ +(a''_{26})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a''_{38})^{(7,7,7,7)} (T_{37}, t) & +(a''_{42})^{(8,8,8,8)} (T_{41}, t) & +(a''_{46})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)} (T_{21}, t)$, $+(a'_{21})^{(3)} (T_{21}, t)$, $+(a'_{22})^{(3)} (T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)} (T_{17}, t)$, $+(a'_{17})^{(2,2,2)} (T_{17}, t)$, $+(a'_{18})^{(2,2,2)} (T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)} (T_{14}, t)$, $+(a'_{14})^{(1,1,1)} (T_{14}, t)$, $+(a'_{15})^{(1,1,1)} (T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(QQQQQ) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(RRRRR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(SSSSS) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(TTTTT)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(UUUUU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(VVVVV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(u) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{31}}), (\widehat{T_{31}})$: $(\widehat{G_{31}}, \widehat{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)}) - (S_2)^{(9)}} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t} \right)$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of $(u_0)^{(5)}$** .

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(u) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b_i'')^{(5)}}{\partial G_j}(G_{31}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)} \right) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO

Thermodynamic Entropy Loss With The Loss Of The Edge Entanglement Entropy

INTRODUCTION—VARIABLES USED

Phys. Rev. D 86, 065007 (2012) DOI: 10.1103/PhysRevD.86.065007 arXiv: 0905.1317 [cond-mat.str-el]
Entanglement Renormalization and Holography Brian Swingle

- (1) Such a perturbation effectively breaks (e) the system in two, and we can identify (e&eb) the thermodynamic entropy loss with the loss of the edge entanglement entropy.
- (2) From these results, authors obtain a simple interpretation of (e) the non-integer ‘ground state degeneracy’ which is (=) obtained in 1+1-dimensional quantum impurity problems: its logarithm is

(=) a 2+1-dimensional topological entanglement entropy **Topological Entanglement Entropy from the Holographic Partition Function**

Spin Quantum information and physics: some future directions Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}, John Preskill

(3) Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}, John Preskill considers some promising future directions for quantum information theory that could influence the development of 21st century physics. Advances in the theory of the distinguishability of superoperators may lead to (eb) new strategies for improving the precision of quantum-limited measurements.

(4) A better grasp of the properties of multi-partite quantum entanglement may lead to (eb) deeper understanding of (e) strongly-coupled dynamics in (eb) quantum many-body systems, quantum field theory, and quantum gravity.

NOTATION

Module One

Such a

perturbation effectively breaks (e) the system in two, and we can identify (e&eb) the thermodynamic entropy loss with the loss of the edge entanglement entropy

G_{13} : Category one of system in two, and we can identify (e&eb) the thermodynamic entropy loss with the loss of the edge entanglement entropy

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of perturbation

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

perturbation effectively breaks the system in two, and we can identify (e&eb) the thermodynamic entropy loss with the loss of the edge entanglement entropy

G_{16} : Category one of perturbation effectively breaks the system in two, and we can identify; thermodynamic entropy loss with the loss of the edge entanglement entropy

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of thermodynamic entropy loss with the loss of the edge entanglement entropy; perturbation effectively breaks the system in two, and we can identify

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

From these results, authors obtain a simple interpretation of the non-integer ‘ground state degeneracy’ which is (=) obtained in 1+1-dimensional

quantum impurity problems: its logarithm is (=) a 2+1-dimensional topological entanglement entropy

Topological Entanglement Entropy from the Holographic Partition Function

G_{20} : Category one of simple interpretation of the non-integer 'ground state degeneracy'

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of obtained in 1+1-dimensional entanglement entropy quantum impurity problems: its logarithm is (=) a 2+1-dimensional topological

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

simple interpretation of the non-integer 'ground state degeneracy' which is (=) obtained in 1+1-dimensional quantum impurity problems: its logarithm is (=) a 2+1-dimensional topological entanglement entropy

G_{24} : Category one of simple interpretation of the non-integer 'ground state degeneracy' which is (=) obtained in 1+1-dimensional quantum impurity problems: its logarithm

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of 2+1-dimensional topological entanglement entropy

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}. John Preskill considers some promising future directions for quantum information theory that could influence the development of 21st century physics.

Advances in the theory of the distinguishability of superoperators may lead to (eb) new strategies for improving the precision of quantum-limited measurements

G_{28} : Category one of Advances in the theory of the distinguishability of superoperators

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of new strategies for improving the precision of quantum-limited measurements

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

A better grasp of the

properties of multi-partite quantum entanglement may lead to (eb) deeper understanding of (e) strongly-coupled dynamics in (eb) quantum many-body systems, quantum field theory, and quantum gravity

G_{32} : Category one of properties of multi-partite quantum entanglement

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of deeper understanding of (e) strongly-coupled dynamics in (eb) quantum many-body systems, quantum field theory, and quantum gravity

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

properties of multi-partite quantum entanglement may lead to deeper understanding of strongly-coupled dynamics in (eb) quantum many-body systems, quantum field theory, and quantum gravity

G_{36} : Category one of properties of multi-partite quantum entanglement may lead to deeper understanding of strongly-coupled dynamics

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of quantum many-body systems, quantum field theory, and quantum gravity

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Quantum mechanics and field theory with fractional spin and statistics Rev. Mod Phys. 64, 193 – Published 1 January 1992 Stefano Forte

- (1) Planar systems admit (e) quantum states that are neither bosons nor fermions, i.e., whose angular momentum is (=) neither integer nor half-integer.
- (2) After a discussion of some examples of familiar models in which fractional spin may arise, the relevant (nonrelativistic) quantum mechanics is developed from first principles. The appropriate generalization of statistics is also discussed. Some physical effects of fractional spin and statistics are worked out explicitly. The group theory underlying relativistic models with fractional spin and statistics is then introduced and applied to (e&eb) relativistic particle mechanics and field theory.
- (3) Field-theoretical models in 2+1 dimensions are presented which admit (e) solitons that carry fractional statistics, and are discussed in a semiclassical approach, in the functional integral approach, and in the canonical approach.

- (4) Finally, fundamental field theories whose Fock states carry (e&eb) fractional spin and statistics are discussed. DOI: <http://dx.doi.org/10.1103/RevModPhys.64.193>

Planar systems admit (e) quantum states that are neither bosons nor fermions, i.e., whose angular momentum is (=) neither integer nor half-integer

G_{40} : Category one of quantum states that are neither bosons nor fermions, i.e., whose angular momentum is (=) neither integer nor half-integer

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Planar systems

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

After a discussion of some examples of familiar models in which fractional spin may arise, the relevant (nonrelativistic) quantum mechanics is developed from first principles. The appropriate generalization of statistics is also discussed. Some physical effects of fractional spin and statistics are worked out explicitly.

The

group theory underlying relativistic models with fractional spin and statistics is then introduced and applied to (e&eb) relativistic particle mechanics and field theory

G_{44} : Category one of group theory underlying relativistic models with fractional spin and statistics; relativistic particle mechanics and field theory

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of relativistic particle mechanics and field theory; group theory underlying relativistic models with fractional spin and statistics

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$

$$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$$

are Accentuation coefficients

$$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)},$$

$$(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$$

$$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$$

$$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$$

$$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$$

$$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$$

$$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad 6$$

$$+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor}$$

$$-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor}$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 56$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 57$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for

1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for

category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1)}(G, t)} & \boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1)}(G, t)} & \boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{46})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a'_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a'_{21})^{(3)}(T_{21}, t)} & \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} \boxed{+(a'_{22})^{(3)}(T_{21}, t)} & \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \quad 69$$

$\boxed{+(a'_{20})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3)}(T_{21}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)}$ are second augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$ are third augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} \boxed{(b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{36})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$\begin{aligned} &= (a_{40})^{(8)} G_{41} \\ &- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \end{aligned}$$

$\frac{dG_{41}}{dt}$

$$\begin{aligned} &= (a_{41})^{(8)} G_{40} \\ &- \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \end{aligned}$$

$\frac{dG_{42}}{dt}$

$$\begin{aligned} &= (a_{42})^{(8)} G_{41} \\ &- \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} + \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T_{29}'| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} \|(G_{31}) - (G_{31})'\| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(WWWWW) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(XXXXX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(YYYYY) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(ZZZZZ)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(AAAAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(BBBBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(v) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$| (G_{19})^{(1)} - (G_{19})^{(2)} | e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e \left[- \int_0^t \{ (a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right] \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} &\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)} \right) d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(v) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b_{20}')^{(3)}(b_{21}')^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b_{20}')^{(3)}(b_{21}'')^{(3)}(G_{23}) + (b_{21}')^{(3)}(b_{20}'')^{(3)}(G_{23})] + (b_{20}'')^{(3)}(G_{23})(b_{21}'')^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b_{24}')^{(4)}(b_{25}')^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 & \left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY THREE

Spin Quantum Information

INTRODUCTION—VARIABLES USED

Spin Quantum information and physics: some future directions Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}, John Preskill

- (1) With the discovery of an apparent separation between the classical and quantum classifications of computational complexity [1], and (e&eb) of fault-tolerant schemes for quantum computation [2], quantum information theory has earned a lasting and prominent place at (eb) the foundations of computer science.

- (2) But at present this discipline seems rather isolated from most of the rest of physics. Will this change in the future? How might it change?
 One view is that thinking about information theory will lead us to a deeper understanding of the foundations of quantum mechanics. This vision has been vividly expressed by John Wheeler [3]; Bill Wootters [9] and Chris Fuchs [5] have been among its particularly eloquent spokespersons. But I am not convinced in my heart that we are supposed to understand the foundations of quantum mechanics much better than we currently do. **John Preskill** prefers to look in a different direction to anticipate where quantum information may have an impact on physics. Our deepening understanding of quantum information may lead to (eb) new strategies for pushing back the boundaries of quantum-limited measurements.
- (3) Quantum entanglement (e&eb) quantum error correction (e&eb) and quantum information processing might all be exploited to (e) improve the information-gathering capability of physics experiments.
- (4) The most challenging and interesting problems in quantum dynamics involve understanding the behavior of strongly-coupled many-body systems — systems with many degrees of freedom that undergo (e&eb) large quantum fluctuations.
- (5) Better ways of characterizing and classifying the features of many particle entanglements may lead to (eb) new and more effective methods for understanding the dynamical behavior of complex quantum system.
- (6) A watchword of quantum information theory is: “Entanglement is a Useful Resource.” It should not be a surprise if entanglement can extend the capabilities of the laboratory physicist. For example, the phenomenon of superdense coding illustrates that shared entanglement can enhance (eb) **classical communication between two parties [13].**
- (7) The same strategy can sometimes be used to exploit entanglement to improve (eb) the distinguishability among Hamiltonians (an idea suggested by Chris Fuchs [14]).
- (8) Suppose I wish to observe the precession of spin-1/2 objects to determine (eb) the value of an unknown magnetic field.

NOTATION

Module One

Field-theoretical models in 2+1 dimensions are presented which admit (e) solitons that carry fractional statistics, and are discussed in a semiclassical approach, in the functional integral approach, and in the canonical approach

G_{13} : Category one of solitons that carry fractional statistics, and are discussed in a semiclassical approach, in the functional integral approach, and in the canonical approach

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Field-theoretical models in 2+1 dimensions

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Finally,

fundamental field theories whose Fock states carry (e&eb) fractional spin and statistics
 are discussed

. DOI: <http://dx.doi.org/10.1103/RevModPhys.64.193>

G_{16} : Category one of fundamental field theories whose Fock states; fractional spin and statistics

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of fractional spin and statistics ;fundamental field theories whose Fock states

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Quantum entanglement (e&eb) quantum error correction (e&eb) and quantum information processing might
 all be exploited to (e) improve the information-gathering capability of physics experiments

G_{20} : Category one of Quantum entanglement; quantum error correction

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of quantum error correction; Quantum entanglement

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Quantum entanglement, quantum error correction (e&eb) and quantum information processing might all be
 exploited to (e) improve the information-gathering capability of physics experiments

G_{24} : Category one of quantum error correction; quantum information processing

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of quantum information processing; quantum error correction

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Quantum entanglement, quantum error correction and quantum information processing might all be
 exploited to improve the information-gathering capability of physics experiments

G_{28} : Category one of Quantum entanglement, quantum error correction and quantum information processing exploited

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of improve the information-gathering capability of physics experiments

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

The most challenging and interesting problems in quantum dynamics involve understanding the behavior of strongly-coupled many-body systems — systems with many degrees of freedom that undergo (e&eb) large quantum fluctuations

G_{32} : Category one of behavior of strongly-coupled many-body systems — systems with many degrees of freedom; large quantum fluctuations

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of large quantum fluctuations ;behavior of strongly-coupled many-body systems — systems with many degrees of freedom

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Better ways of characterizing and classifying the features of many particle entanglements may lead to (eb) new and more effective methods for understanding the dynamical behavior of complex quantum system

G_{36} : Category one of characterizing and classifying the features of many particle entanglements

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of effective methods for understanding the dynamical behavior of complex quantum system

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

A watchword of quantum information theory is: “Entanglement is a Useful Resource.” It should not be a surprise if entanglement can extend the capabilities of the laboratory physicist

. For example, the phenomenon of superdense coding illustrates that shared entanglement can enhance (eb) **classical communication between two parties**

G_{40} : Category one of phenomenon of superdense coding illustrates that shared entanglement

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **classical communication between two parties**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

The

same strategy can sometimes be used to exploit entanglement to improve (eb) the distinguishability among Hamiltonians (an idea suggested by Chris Fuchs)

G_{44} : Category one of same strategy can sometimes be used to exploit entanglement

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of distinguishability among Hamiltonians (an idea suggested by Chris Fuchs)

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

42

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40}$$

43

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41}$$

44

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42}$$

45

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40}$$

46

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41}$$

47

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42}$$

48

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44}$$

49

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45}$$

50

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46}$$

51

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44}$$

52

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45}$$

53

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

55

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

56

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

57

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$, $(a''_{21})^{(3,3)}(T_{21}, t)$, $(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient

for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient

for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} (G_{19}, t) & -(b''_{15})^{(1,1)} (G, t) & -(b''_{22})^{(3,3,3)} (G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)} (G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)} (G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)} (G_{35}, t) \\ -(b''_{38})^{(7,7,7)} (G_{39}, t) & -(b''_{42})^{(8,8,8)} (G_{43}, t) & -(b''_{46})^{(9,9)} (G_{47}, t) \end{array} \right] T_{18}$$

where $-(b'_{16})^{(2)} (G_{19}, t)$, $-(b'_{17})^{(2)} (G_{19}, t)$, $-(b'_{18})^{(2)} (G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)} (G, t)$, $-(b'_{14})^{(1,1)} (G, t)$, $-(b'_{15})^{(1,1)} (G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)} (G_{23}, t)$, $-(b''_{21})^{(3,3,3)} (G_{23}, t)$, $-(b''_{22})^{(3,3,3)} (G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)} (G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)} (G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)} (G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)} (G_{39}, t)$, $-(b''_{37})^{(7,7,7)} (G_{39}, t)$, $-(b''_{38})^{(7,7,7)} (G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)} (G_{43}, t)$, $-(b''_{41})^{(8,8,8)} (G_{43}, t)$, $-(b''_{42})^{(8,8,8)} (G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)} (G_{47}, t)$, $-(b''_{46})^{(9,9)} (G_{47}, t)$, $-(b''_{45})^{(9,9)} (G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} (T_{21}, t) & +(a''_{16})^{(2,2,2)} (T_{17}, t) & +(a'_{13})^{(1,1,1)} (T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a''_{36})^{(7,7,7,7)} (T_{37}, t) & +(a''_{40})^{(8,8,8,8)} (T_{41}, t) & +(a''_{44})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} (T_{21}, t) & +(a''_{17})^{(2,2,2)} (T_{17}, t) & +(a'_{14})^{(1,1,1)} (T_{14}, t) \\ +(a''_{25})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a''_{37})^{(7,7,7,7)} (T_{37}, t) & +(a''_{41})^{(8,8,8,8)} (T_{41}, t) & +(a''_{45})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} (T_{21}, t) & +(a''_{18})^{(2,2,2)} (T_{17}, t) & +(a'_{15})^{(1,1,1)} (T_{14}, t) \\ +(a''_{26})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a''_{38})^{(7,7,7,7)} (T_{37}, t) & +(a''_{42})^{(8,8,8,8)} (T_{41}, t) & +(a''_{46})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)} (T_{21}, t)$, $+(a'_{21})^{(3)} (T_{21}, t)$, $+(a'_{22})^{(3)} (T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)} (T_{17}, t)$, $+(a'_{17})^{(2,2,2)} (T_{17}, t)$, $+(a'_{18})^{(2,2,2)} (T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)} (T_{14}, t)$, $+(a'_{14})^{(1,1,1)} (T_{14}, t)$, $+(a'_{15})^{(1,1,1)} (T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{36})^{(7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{37})^{(7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$\begin{aligned} &= (a_{40})^{(8)} G_{41} \\ &- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \end{aligned}$$

$\frac{dG_{41}}{dt}$

$$\begin{aligned} &= (a_{41})^{(8)} G_{40} \\ &- \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a'_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \end{aligned}$$

$\frac{dG_{42}}{dt}$

$$\begin{aligned} &= (a_{42})^{(8)} G_{41} \\ &- \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a'_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} \|(G_{31}) - (G_{31})'\| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(CCCCC) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(DDDDDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(EEEEEE) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(FFFFFF)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(GGGGGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(HHHHHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\tilde{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\tilde{M}_{28})^{(5)}} \left[((\tilde{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\tilde{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\tilde{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\tilde{P}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\tilde{P}_{32})^{(6)}}{(\tilde{M}_{32})^{(6)}} \left(e^{(\tilde{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\tilde{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\tilde{M}_{32})^{(6)}} \left[((\tilde{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\tilde{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\tilde{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(w) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\tilde{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\tilde{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\tilde{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[((\tilde{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\tilde{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\tilde{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\tilde{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\tilde{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\tilde{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[((\tilde{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\tilde{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\tilde{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$| (G_{19})^{(1)} - (G_{19})^{(2)} | e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e \left[- \int_0^t \{ (a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right] \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{ (a'_i)^{(4)} - (a''_i)^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \} ds_{(24)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t}$$

376

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq$$

377

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $v^{(2)} = \frac{G_{16}}{G_{17}}$ 388

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t) :-$

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t) :-$

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(w) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY FOUR

Outcome Of The Measurement Of The First Spin Is Permitted To Influence The Choice Of The Measurement That Is Performed On The Second Spin

INTRODUCTION—VARIABLES USED

Paul Fendley,¹ Matthew P. A. Fisher² and Chetan Nayak^{3,4}. John Preskill **spin** Quantum information and physics: some future directions

- (1) If two spins are available, one way to estimate the value of the unknown field is to allow (eb) both spins to precess in the field independently, and then measure (eb) them separately.
- (2) An alternative method is to prepare an entangled Bell pair, expose (e&eb) one of the two spins to

the magnetic field while the other is carefully shielded (e&eb) from the field, and finally carry out (eb) a collective Bell measurement on the pair.

- (3) It turns out that in many cases (for example when we have no a priori knowledge about the field direction), the entangled strategy extracts (eb) more information about the unknown field than the strategy in which uncorrelated spins are (=) measured one at a time [6].
- (4) This separation still holds even if we allow the unentangled strategy to be (=) adaptive; that is, even if the **outcome of the measurement of the first spin is permitted to influence (e&eb) the choice of the measurement that is performed on the second spin** Quantum information and physics: some future directions John Preskill.

NOTATION

Module One

If two spins are available, one way to estimate the value of the unknown field is to allow (eb) both spins to precess in the field independently, and then measure (eb) them separately

G_{13} : Category one of two spins are available, one way to estimate the value of the unknown field is to allow

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of both spins to precess in the field independently, and then measure (eb) them separately

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

If

two spins are available, one way to estimate the value of the unknown field is to allow both spins to precess in the field independently, and then measure (eb) them separately

G_{16} : Category one of two spins are available, one way to estimate the value of the unknown field is to allow both spins to precess in the field independently, and then measure

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of them separately (there may be many such twosome; or take all the three categories as equivalent)

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

An alternative method is to

prepare an entangled Bell pair, expose (e&eb) one of the two spins to the magnetic field while the other is carefully shielded (e&eb) from the field, and finally carry out (eb) a collective Bell measurement on the pair

G_{20} : Category one of prepare an entangled Bell pair; one of the two spins to the magnetic field while the other is carefully shielded (e&eb) from the field, and finally carry out (eb) a collective Bell measurement on the pair

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of one of the two spins to the magnetic field while the other is carefully shielded (e&eb) from the field, and finally carry out (eb) a collective Bell measurement on the pair; prepare an entangled Bell pair

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

prepare an entangled Bell pair, expose one of the two spins to the magnetic field while the other is carefully shielded from the field, and finally carry out (eb) a collective Bell measurement on the pair

G_{24} : Category one of preparation of an entangled Bell pair, expose one of the two spins to the magnetic field while the other is carefully shielded from the field

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of collective Bell measurement on the pair

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

It turns out that in many cases (for example when we have no a priori knowledge about the field direction), the

Entangled strategy extracts (eb) more information about the unknown field than the strategy in which uncorrelated spins are (=) measured one at a time [6].

G_{28} : Category one of entangled strategy

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of more information about the unknown field than the strategy in which uncorrelated spins are (=) measured one at a time [6].(for references kindly see the original article)

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Entangled strategy extracts more information about the unknown field than the strategy in which uncorrelated spins are (=) measured one at a time [6].

G_{32} : Category one of Entangled strategy extracts more information about the unknown field than the strategy in which uncorrelated spins

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of measured one at a time

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

This

separation still holds even if we allow the unentangled strategy to be (=) adaptive; that is, even if the **outcome of the measurement of the first spin is permitted to influence (e&eb) the choice of the measurement that is performed on the second spin**

Quantum information and physics: some future directions John Preskill

G_{36} : Category one of separation still holds even if we allow the unentangled strategy

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of adaptive; that is, even if the **outcome of the measurement of the first spin is permitted to influence (e&eb) the choice of the measurement that is performed on the second spin**

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

separation still holds even if we allow the unentangled strategy to be adaptive; that is, even if the **outcome of the measurement of the first spin is permitted to influence (e&eb) the choice of the measurement that is performed on the second spin**

G_{40} : Category one of separation still holds even if we allow the unentangled strategy to be adaptive; that is, even if the **outcome of the measurement of the first spin; choice of the measurement that is performed on the second spin**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **choice of the measurement that is performed on the second spin** ;separation still holds even if we allow the unentangled strategy to be adaptive; that is, even if the **outcome of the measurement of the first spin**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

separation still holds even if we allow the unentangled strategy to be adaptive; that is, even if the **outcome of the measurement of the first spin is permitted to influence (e&eb) the choice of the measurement that is performed on the second spin**

G_{44} : Category one of **measurement of the first spin; measurement that is performed on the second spin**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **measurement that is performed on the second spin; measurement of the first spin**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)},$
 $(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &- (b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &- (b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20\end{aligned}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b'_{20})^{(3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{36})^{(7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b'_{21})^{(3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{37})^{(7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & - (b''_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b'_{22})^{(3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{38})^{(7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) + (a''_{24})^{(4,4)}(T_{25}, t) + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b''_{30})^{(5,5,5)}(G_{31}, t) & -(b''_{26})^{(4,4,4)}(G_{27}, t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t)$, $-(b''_{33})^{(6)}(G_{35}, t)$, $-(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}' - G_{31}|| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(IIIIII) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(JJJJJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(KKKKKK) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(LLLLLL)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(MMMMMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(NNNNNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(x) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{((\hat{P}_{13})^{(1)} + G_j^0)}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{((\hat{Q}_{13})^{(1)} + T_j^0)}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : (\widehat{G}_{31}), (\widehat{T}_{31}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $(\bar{M}_{44})^{(9)}_1, (\bar{M}_{44})^{(9)}_2$ and $(\bar{M}_{44})^{(9)}_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right]$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 409

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-

$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(x) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b'_{14})^{(1)}(G) + (b'_{14})^{(1)}(b'_{13})^{(1)}(G)] + (b'_{13})^{(1)}(G)(b'_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b'_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b'_{16})^{(2)}(G_{19})] + (b'_{16})^{(2)}(G_{19})(b'_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b'_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b'_{20})^{(3)}(G_{23})] + (b'_{20})^{(3)}(G_{23})(b'_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}(G_{43}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY FIVE

Compressible Quantum Phases

INTRODUCTION—VARIABLES USED

Compressible quantum phases from conformal field theories in 2+1 dimensions Subir Sachdev

- (1) Conformal field theories (CFTs) with a globally conserved U(1) charge Q can be deformed into (e&eb) compressible phases by modifying their Hamiltonian, H, by(e) a chemical potential H ! H “μQ.
- (2) Authors study 2+1 dimensional CFTs upon which an explicit (e&eb) S duality mapping can be performed.

- (3) **Subir Sachdev** finds that this construction leads (e) naturally to compressible phases which are (=) superfluids, solids, or non-Fermi liquids which are more appropriately called ‘Bose metals’ in the present context.
- (4) The Bose metal preserves all symmetries and has (e) Fermi surfaces of gauge-charged fermions, even in cases where the parent CFT can be expressed solely by (e) bosonic degrees of freedom.
- (5) Monopole operators are identified as (=) order parameters of the solids, and the product of their magnetic charge and Q determines (=) the area of the unit cell.
- (6) Author presents implications for holographic theories on (e&eb) asymptotically AdS_4 spacetimes: S duality and monopole/dyon fields play important roles in this connection **Compressible quantum phases from conformal field theories in 2+1 dimensions Subir Sachdev**

NOTATION

Module One

- Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q can be deformed into (e&eb) compressible phases by modifying their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$
- G_{13} : Category one of Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q ; compressible phases by modifying their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$
- G_{14} : Category two of SAS (same as superior/above)
- G_{15} : Category three of SAS
- T_{13} : Category one of compressible phases by modifying their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$; Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q
- T_{14} : Category two of SAS
- T_{15} : Category three of SAS

Module Two

- Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q can be deformed into compressible phases by modifying (e&eb) their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$
- G_{16} : Category one of Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q can be deformed into compressible phases; their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$
- G_{17} : Category two of SAS
- G_{18} : Category three of SAS
- T_{16} : Category one of their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$; Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q can be deformed into compressible phases
- T_{17} : Category two of SAS
- T_{18} : Category three of SAS

Module three

- Conformal field theories (CFTs) with a globally conserved $U(1)$ charge Q can be deformed into compressible phases by modifying their Hamiltonian, H , by (e) a chemical potential $H \rightarrow H - \mu Q$
- G_{20} : Category one of chemical potential $H \rightarrow H - \mu Q$

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Conformal field theories (CFTs) with a globally conserved U (1) charge Q can be deformed into compressible phases by modifying their Hamiltonian, H

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Authors study

2+1 dimensional CFTs upon which an explicit (e&eb) S duality mapping can be performed.

G_{24} : Category one of 2+1 dimensional CFTs; S duality mapping

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of S duality mapping ;2+1 dimensional CFTs

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Subir Sachdev finds that this construction leads (eb) naturally to compressible phases which are (=) superfluids, solids, or non-Fermi liquids which are more appropriately called ‘Bose metals’ in the present context

G_{28} : Category one of 2+1 dimensional CFTs upon which an explicit S duality mapping can be performed

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of compressible phases which are (=) superfluids, solids, or non-Fermi liquids which are more appropriately called ‘Bose metals’ in the present context

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

The Bose metal preserves all symmetries and has (e) Fermi surfaces of gauge-charged fermions, even in cases where the parent CFT can be expressed solely by (e) bosonic degrees of freedom

G_{32} : Category one of Fermi surfaces of gauge-charged fermions, even in cases where the parent CFT can be expressed solely by (e) bosonic degrees of freedom

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Bose metal preserves all symmetries

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Bose metal preserves all symmetries and has Fermi surfaces of gauge-charged fermions, even in cases where the parent CFT can be expressed solely by (e) bosonic degrees of freedom

G_{36} : Category one of bosonic degrees of freedom

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Bose metal preserves all symmetries and has Fermi surfaces of gauge-charged fermions, even in cases where the parent CFT can be expressed

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Monopole operators are identified as (=) order parameters of the solids, and the product of their magnetic charge and Q determines (=) the area of the unit cell

G_{40} : Category one of Monopole operators are identified

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of order parameters of the solids, and the product of their magnetic charge and Q determines (=) the area of the unit cell

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Monopole operators are identified as order parameters of the solids, and the product of their magnetic charge and Q determines (=) the area of the unit cell

G_{44} : Category one of Monopole operators are identified as order parameters of the solids, and the product of their magnetic charge and Q

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of area of the unit cell

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 7$$

$$\begin{aligned}\frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} & 20 \\ \frac{dG_{26}}{dt} &= (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} & 21 \\ \frac{dT_{24}}{dt} &= (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} & 22 \\ \frac{dT_{25}}{dt} &= (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} & 23 \\ \frac{dT_{26}}{dt} &= (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} & 24 \\ &+ (a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor} \\ &-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\begin{aligned}\frac{dG_{28}}{dt} &= (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} & 25 \\ \frac{dG_{29}}{dt} &= (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} & 26 \\ \frac{dG_{30}}{dt} &= (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} & 27 \\ \frac{dT_{28}}{dt} &= (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} & 28 \\ \frac{dT_{29}}{dt} &= (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} & 29 \\ \frac{dT_{30}}{dt} &= (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} & 30\end{aligned}$$

$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$

$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$, $(a''_{17})^{(2,2)}(T_{17}, t)$, $(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$, $(a''_{21})^{(3,3)}(T_{21}, t)$, $(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7)}(T_{37}, t)$, $(a''_{37})^{(7,7)}(T_{37}, t)$, $(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8)}(T_{41}, t)$, $(a''_{41})^{(8,8)}(T_{41}, t)$, $(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] & -(b''_{18})^{(2,2)}(G_{19}, t) & -(b''_{22})^{(3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7)}(G_{39}, t) & -(b''_{42})^{(8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $[-(b'_{13})^{(1)}(G, t)]$, $[-(b'_{14})^{(1)}(G, t)]$, $[-(b'_{15})^{(1)}(G, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b'_{16})^{(2,2)}(G_{19}, t)]$, $[-(b'_{17})^{(2,2)}(G_{19}, t)]$, $[-(b'_{18})^{(2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b'_{20})^{(3,3)}(G_{23}, t)]$, $[-(b'_{21})^{(3,3)}(G_{23}, t)]$, $[-(b'_{22})^{(3,3)}(G_{23}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b'_{24})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b'_{25})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b'_{26})^{(4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b'_{28})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b'_{29})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b'_{30})^{(5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b'_{32})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b'_{33})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b'_{34})^{(6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b'_{37})^{(7,7)}(G_{39}, t)]$, $[-(b'_{36})^{(7,7)}(G_{39}, t)]$, $[-(b'_{38})^{(7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b'_{40})^{(8,8)}(G_{43}, t)]$, $[-(b'_{41})^{(8,8)}(G_{43}, t)]$, $[-(b'_{42})^{(8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17}, t)] & +(a''_{13})^{(1,1)}(T_{14}, t) & +(a''_{20})^{(3,3,3)}(T_{21}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{36})^{(7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)}[+(a''_{17})^{(2)}(T_{17}, t)] & +(a''_{14})^{(1,1)}(T_{14}, t) & +(a''_{21})^{(3,3,3)}(T_{21}, t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{37})^{(7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}[+(a''_{18})^{(2)}(T_{17}, t)] & +(a''_{15})^{(1,1)}(T_{14}, t) & +(a''_{22})^{(3,3,3)}(T_{21}, t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{38})^{(7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $[(a'_{16})^{(2)}(T_{17}, t)]$, $[(a'_{17})^{(2)}(T_{17}, t)]$, $[(a'_{18})^{(2)}(T_{17}, t)]$ are first augmentation coefficients for category 1, 2 and 3

$[(a'_{13})^{(1,1)}(T_{14}, t)]$, $[(a'_{14})^{(1,1)}(T_{14}, t)]$, $[(a'_{15})^{(1,1)}(T_{14}, t)]$ are second augmentation coefficient for category 1, 2 and 3

$[(a'_{20})^{(3,3,3)}(T_{21}, t)]$, $[(a'_{21})^{(3,3,3)}(T_{21}, t)]$, $[(a'_{22})^{(3,3,3)}(T_{21}, t)]$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a''_{16})^{(2,2,2)}(T_{17}, t) + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a''_{17})^{(2,2,2)}(T_{17}, t) + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a''_{18})^{(2,2,2)}(T_{17}, t) + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{l} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{16})^{(2,2,2)}(G_{19}, t) - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{l} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) - (b''_{17})^{(2,2,2)}(G_{19}, t) - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$[-(b'_{20})^{(3)}(G_{23}, t)], [-(b'_{21})^{(3)}(G_{23}, t)], [-(b'_{22})^{(3)}(G_{23}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b'_{16})^{(2,2,2)}(G_{19}, t)], [-(b'_{17})^{(2,2,2)}(G_{19}, t)], [-(b'_{18})^{(2,2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b'_{13})^{(1,1,1)}(G, t)], [-(b'_{14})^{(1,1,1)}(G, t)], [-(b'_{15})^{(1,1,1)}(G, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b'_{36})^{(7,7,7,7)}(G_{39}, t)], [-(b'_{37})^{(7,7,7,7)}(G_{39}, t)], [-(b'_{38})^{(7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b'_{40})^{(8,8,8,8)}(G_{43}, t)], [-(b'_{41})^{(8,8,8,8)}(G_{43}, t)], [-(b'_{42})^{(8,8,8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b'_{46})^{(9,9,9)}(G_{47}, t)], [-(b'_{45})^{(9,9,9)}(G_{47}, t)], [-(b'_{44})^{(9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)}[+(a''_{24})^{(4)}(T_{25}, t)] & +(a'_{28})^{(5,5)}(T_{29}, t) & +(a'_{32})^{(6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)}[+(a''_{25})^{(4)}(T_{25}, t)] & +(a'_{29})^{(5,5)}(T_{29}, t) & +(a'_{33})^{(6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)}[+(a''_{26})^{(4)}(T_{25}, t)] & +(a'_{30})^{(5,5)}(T_{29}, t) & +(a'_{34})^{(6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3)}(T_{21}, t) \\ +(a'_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$[(a'_{24})^{(4)}(T_{25}, t)], [(a'_{25})^{(4)}(T_{25}, t)], [(a'_{26})^{(4)}(T_{25}, t)]$ are first augmentation coefficients category 1, 2 3

$[(a'_{28})^{(5,5)}(T_{29}, t)], [(a'_{29})^{(5,5)}(T_{29}, t)], [(a'_{30})^{(5,5)}(T_{29}, t)]$ are second augmentation coefficient for category 1, 2 and 3

$[(a'_{32})^{(6,6)}(T_{33}, t)], [(a'_{33})^{(6,6)}(T_{33}, t)], [(a'_{34})^{(6,6)}(T_{33}, t)]$ are third augmentation coefficient for category 1, 2 and 3

$[(a'_{13})^{(1,1,1,1)}(T_{14}, t)], [(a'_{14})^{(1,1,1,1)}(T_{14}, t)], [(a'_{15})^{(1,1,1,1)}(T_{14}, t)]$

are fourth augmentation coefficients for category 1, 2 and 3

$$+(a''_{16})^{(2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2)}(T_{17}, t)$$

are fifth augmentation coefficients for category 1, 2 and 3

$$+(a''_{20})^{(3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3)}(T_{21}, t)$$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$$

are sixth detrition coefficients for category 1, 2 and 3

$$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1, 2 and 3

$$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2 and 3

$$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a''_{25})^{(4,4)}(T_{25}, t) & + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1, 2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1, 2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1, 2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1, 2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 86$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 87$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t)$, $+(a''_{33})^{(6)}(T_{33}, t)$, $+(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)} \text{ sixth augmentation}$$

coefficients

$$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)}$$

seventh augmentation coefficients

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$$

Eighth augmentation coefficients

$$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)} \text{ ninth augmentation}$$

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} \boxed{(b'_{32})^{(6)}} \boxed{-(b''_{32})^{(6)}(G_{35}, t)} \boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)} \boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} \boxed{(b'_{33})^{(6)}} \boxed{-(b''_{33})^{(6)}(G_{35}, t)} \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)} \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} \boxed{(b'_{34})^{(6)}} \boxed{-(b''_{34})^{(6)}(G_{35}, t)} \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{34} \quad 90$$

$$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6)}(G_{35}, t)} \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t)} \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$$

are eighth detrition coefficients for category 1, 2, and 3

$$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(000000) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(PPPPPP) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(QQQQQQ) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(RRRRRR)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(SSSSSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(TTTTTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(y) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right] \quad 181$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})_1$ it follows $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \quad \text{If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$|\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} +$$

$$\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} \leq$$

$$\frac{1}{(\bar{M}_{20})^{(3)}} \left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)} \right) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ & G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}\right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a'_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a'_{36})^{(7)}$ and $(b'_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\bar{G}_{40}^{(1)} - \bar{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$\begin{aligned} (m_2)^{(1)} &= (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ (m_2)^{(1)} &= (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} (\mu_2)^{(1)} &= (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ (\mu_2)^{(1)} &= (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)}) - (S_2)^{(2)}} \right) \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

$$\text{roots of the equations } (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq \quad 334$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$: 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$ $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case**.

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$** .

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(y) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}) T_{13}^* \mathbb{G}_j \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}) T_{14}^* \mathbb{G}_j \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}) T_{15}^* \mathbb{G}_j \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*\mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*\mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*\mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)})$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)})$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ \left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \right. \\
 &\left. \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 &+ \left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \right. \\
 &\left. \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY SIX

Quantum Gravity As The Natural Regulator Of The Hamiltonian Constraint

INTRODUCTION—VARIABLES USED

The quantization of topology, from quantum Hall effect to quantum gravity Andrei T. Patrascu.

- (1) **Andrei T. Patrascu** extends the **notion of quantization** from the standard interpretation focused on non-commuting observables defined starting from classical analogues, to the (eb) topological equivalents defined in terms of (e&eb) coefficient groups in (co)homology.
- (2) It is shown that the commutation relations between quantum observables become (=) (non)compatibility relations between coefficient groups.
- (3) Main result is the construction of a new, higher-level form of quantization, as seen from (e) the perspective of the universal coefficient theorem.

The quantization of topology, from quantum Hall effect to quantum gravity Andrei T. Patrascu.

- (4) This idea brings us closer to a consistent quantization of gravity, allows for a (eb) systematic description of topology changing string interactions but also gives (eb) new, quantum-topological degrees of freedom in discussions involving (e&eb) quantum information.
- (5) On the practical side, a possible connection to the fractional quantum Hall effect is explored. ArXiv: 1411.4475 [physics.gen-ph]: **The quantization of topology, from quantum Hall effect to quantum gravity Andrei T. Patrascu.**

T Thiemann 1998 Class Quantum Grav.15 1281 doi:10.1088/0264-9381/15/5/012 Quantum spin dynamics (QSD): V. Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories

- (6) It is an old speculation and prognostication in physics that, once the gravitational field is successfully quantized, it should serve as the natural regulator (e&eb) of infrared and ultraviolet singularities that plague (e) quantum field theories in a background metric.
- (7) **T Thiemann** demonstrates that at least part of this idea is implemented in a precise sense within the framework of (e) four-dimensional canonical Lorentzian quantum gravity in the continuum.
- (8) Specifically, he shows that the Hamiltonian of the standard model supports (eb) a representation in which finite linear combinations of (e) Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operators.
- (9) This Hamiltonian, surprisingly, does not (e) suffer from any singularities; it is (=) completely finite without renormalization.

- (10) This property is shared by string theory. In contrast to string theory, however, we are dealing with a particular phase of the standard model coupled to (e&eb) gravity which is entirely non-perturbatively defined and **second quantized**.
- (11) Of course, to show that the entire theory is finite requires more: one would need to know what the physical observables are, apart from the Hamiltonian constraint, and whether they are also finite. However, with the results given in this paper this question can now be answered, at least in principle. **T Thiemann 1998 Class Quantum Grav.15 1281 doi:10.1088/0264-9381/15/5/012 Quantum spin dynamics (QSD): V. Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories.**

NOTATION

Module One

Andrei T. Patrascu extends the

notion of quantization from the standard interpretation focused on non-commuting observables defined starting from classical analogues, to the (eb) topological equivalents defined in terms of (e&eb) coefficient groups in (co)homology

G_{13} : Category one of **notion of quantization** from the standard interpretation focused on non-commuting observables defined starting from classical analogues

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of topological equivalents defined in terms of (e&eb) coefficient groups in (co)homology

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Andrei T. Patrascu extends the

notion of quantization from the standard interpretation focused on non-commuting observables defined starting from classical analogues, to the topological equivalents defined in terms of (e&eb) coefficient groups in (co)homology

G_{16} : Category one of **notion of quantization** from the standard interpretation focused on non-commuting observables defined starting from classical analogues, to the topological equivalents; coefficient groups in (co)homology

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of coefficient groups in (co)homology **notion of quantization** from the standard interpretation focused on non-commuting observables defined starting from classical analogues, to the topological equivalents

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

It is shown that the

commutation relations between quantum observables become (=) (non)compatibility relations between coefficient groups

G_{20} : Category one of commutation relations between quantum observables

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of (non)compatibility relations between coefficient groups

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Main result is the

construction of a new, higher-level form of quantization, as seen from (e) the perspective of the universal coefficient theorem

G_{24} : Category one of perspective of the universal coefficient theorem; construction of a new, higher-level form of quantization

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of construction of a new, higher-level form of quantization; perspective of the universal coefficient theorem

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

This idea brings us closer to a

consistent quantization of gravity, allows for a (eb) systematic description of topology changing string interactions but also gives (eb) new, quantum-topological degrees of freedom in discussions involving (e&eb) quantum information

G_{28} : Category one of consistent quantization of gravity

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of systematic description of topology changing string interactions but also gives (eb) new, quantum-topological degrees of freedom in discussions involving (e&eb) quantum information

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

consistent quantization of gravity, allows for a systematic description of topology changing string interactions but also gives (eb) new, quantum-topological degrees of freedom in discussions involving (e&eb) quantum information

G_{32} : Category one of consistent quantization of gravity, allows for a systematic description of topology changing string interactions

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of new, quantum-topological degrees of freedom involving (e&eb) quantum information

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

consistent quantization of gravity, allows for a systematic description of topology changing string interactions but also gives new, quantum-topological degrees of freedom involving (e&eb) quantum information

G_{36} : Category one of consistent quantization of gravity, allows for a systematic description of topology changing string interactions but also gives new, quantum-topological degrees of freedom; quantum information

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of quantum information; consistent quantization of gravity, allows for a systematic description of topology changing string interactions but also gives new, quantum-topological degrees of freedom

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

On the practical side, a possible connection to the fractional quantum Hall effect is explored

. ArXiv: 1411.4475 [physics.gen-ph]: **The quantization of topology, from quantum Hall effect to quantum gravity** Andrei T. Patrascu

G_{40} : Category one of **quantization of topology**; fractional quantum Hall effect

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of fractional quantum Hall effect ;**quantization of topology**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

It is an old speculation and prognostication in physics that, once the

gravitational field successfully quantized, it should serve as the natural regulator (e&eb) of infrared and ultraviolet singularities that plague (e) quantum field theories in a background metric

G_{44} : Category one of gravitational field successfully quantized; infrared and ultraviolet singularities that plague (e) quantum field theories in a background metric

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of infrared and ultraviolet singularities that plague (e) quantum field theories in a background metric ;gravitational field successfully quantized

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

42

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40}$$

43

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41}$$

44

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42}$$

45

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40}$$

46

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41}$$

47

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42}$$

48

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44}$$

49

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45}$$

50

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46}$$

51

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44}$$

52

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45}$$

53

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

55

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

56

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

57

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient

for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient

for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} (G_{19}, t) & -(b''_{18})^{(2)} (G_{19}, t) & -(b'_{15})^{(1,1)} (G, t) & -(b''_{22})^{(3,3,3)} (G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)} (G_{27}, t) & -(b'_{30})^{(5,5,5,5,5)} (G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)} (G_{35}, t) & \\ -(b'_{38})^{(7,7,7)} (G_{39}, t) & -(b''_{42})^{(8,8,8)} (G_{43}, t) & -(b'_{46})^{(9,9)} (G_{47}, t) & \end{array} \right] T_{18}$$

where $-(b'_{16})^{(2)} (G_{19}, t)$, $-(b'_{17})^{(2)} (G_{19}, t)$, $-(b'_{18})^{(2)} (G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)} (G, t)$, $-(b'_{14})^{(1,1)} (G, t)$, $-(b'_{15})^{(1,1)} (G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b'_{20})^{(3,3,3)} (G_{23}, t)$, $-(b'_{21})^{(3,3,3)} (G_{23}, t)$, $-(b'_{22})^{(3,3,3)} (G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b'_{24})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4)} (G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b'_{28})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5)} (G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b'_{32})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6)} (G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b'_{36})^{(7,7,7)} (G_{39}, t)$, $-(b'_{37})^{(7,7,7)} (G_{39}, t)$, $-(b'_{38})^{(7,7,7)} (G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b'_{40})^{(8,8,8)} (G_{43}, t)$, $-(b'_{41})^{(8,8,8)} (G_{43}, t)$, $-(b'_{42})^{(8,8,8)} (G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b'_{44})^{(9,9)} (G_{47}, t)$, $-(b'_{46})^{(9,9)} (G_{47}, t)$, $-(b'_{45})^{(9,9)} (G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} (T_{21}, t) & +(a''_{20})^{(3)} (T_{21}, t) & +(a'_{16})^{(2,2,2)} (T_{17}, t) & +(a'_{13})^{(1,1,1)} (T_{14}, t) \\ +(a'_{24})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)} (T_{29}, t) & +(a'_{32})^{(6,6,6,6,6)} (T_{33}, t) & \\ +(a'_{36})^{(7,7,7,7)} (T_{37}, t) & +(a'_{40})^{(8,8,8,8)} (T_{41}, t) & +(a'_{44})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{20}$$

67

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} (T_{21}, t) & +(a'_{17})^{(2,2,2)} (T_{17}, t) & +(a'_{14})^{(1,1,1)} (T_{14}, t) \\ +(a'_{25})^{(4,4,4,4,4)} (T_{25}, t) & +(a'_{29})^{(5,5,5,5,5)} (T_{29}, t) & +(a'_{33})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a'_{37})^{(7,7,7,7)} (T_{37}, t) & +(a'_{41})^{(8,8,8,8)} (T_{41}, t) & +(a'_{45})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{21}$$

68

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} (T_{21}, t) & +(a'_{18})^{(2,2,2)} (T_{17}, t) & +(a'_{15})^{(1,1,1)} (T_{14}, t) \\ +(a'_{26})^{(4,4,4,4,4)} (T_{25}, t) & +(a'_{30})^{(5,5,5,5,5)} (T_{29}, t) & +(a'_{34})^{(6,6,6,6,6)} (T_{33}, t) \\ +(a'_{38})^{(7,7,7,7)} (T_{37}, t) & +(a'_{42})^{(8,8,8,8)} (T_{41}, t) & +(a'_{46})^{(9,9,9)} (T_{45}, t) \end{array} \right] G_{22}$$

69

$+(a'_{20})^{(3)} (T_{21}, t)$, $+(a'_{21})^{(3)} (T_{21}, t)$, $+(a'_{22})^{(3)} (T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)} (T_{17}, t)$, $+(a'_{17})^{(2,2,2)} (T_{17}, t)$, $+(a'_{18})^{(2,2,2)} (T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)} (T_{14}, t)$, $+(a'_{14})^{(1,1,1)} (T_{14}, t)$, $+(a'_{15})^{(1,1,1)} (T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b''_{16})^{(2,2,2)}(G_{19}, t) & -(b''_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b''_{17})^{(2,2,2)}(G_{19}, t) & -(b''_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b''_{18})^{(2,2,2)}(G_{19}, t) & -(b''_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(UUUUUU) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

$$(VVVVVV) \quad \text{The functions } (a_i'')^{(7)}, (b_i'')^{(7)} \text{ are positive continuous increasing and bounded.}$$

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(WWWWW) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(XXXXXX)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(YYYYYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(ZZZZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(z) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d \left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}) \right) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{31}}), (\widehat{T_{31}})$: $(\widehat{G_{31}}, \widehat{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on
 (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \right) \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t) :-$

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t) :-$

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (\bar{v}_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)}$ and $(b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(z) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

$$\text{ASYMPTOTIC STABILITY ANALYSIS} \quad 531$$

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* T_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* T_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* \mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* \mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* \mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^* \mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^* \mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)} T_{28}^* \mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)} T_{29}^* \mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)} T_{30}^* \mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + \left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \{ (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \} \\ & \left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + \left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) \left((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)} \right) \{ (\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)} \} \\ & \left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY SEVEN

Hamiltonian Constraint Of Matter Quantum Field Theories

INTRODUCTION—VARIABLES USED

T Thiemann 1998 Class Quantum Grav.15 1281 doi:10.1088/0264-9381/15/5/012 Quantum spin dynamics (QSD): V. Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories

- (1) It is an old speculation and prognostication in physics that, once the gravitational field is successfully quantized, it should serve as the natural regulator (e&eb) of infrared and ultraviolet

singularities that plague (e) quantum field theories in a background metric.

- (2) **T Thiemann** demonstrates that at least part of this idea is implemented in a precise sense within the framework of (e) four-dimensional canonical Lorentzian quantum gravity in the continuum.
- (3) Specifically, he shows that the Hamiltonian of the standard model supports (eb) a representation in which finite linear combinations of (e) Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operators.
- (4) This Hamiltonian, surprisingly, does not (e) suffer from any singularities; it is (=) completely finite without renormalization.
- (5) This property is shared by string theory. In contrast to string theory, however, we are dealing with a particular phase of the standard model coupled to (e&eb) gravity which is entirely non-perturbatively defined and **second quantized**.
- (6) Of course, to show that the entire theory is finite requires more: one would need to know what the physical observables are, apart from the Hamiltonian constraint, and whether they are also finite. However, with the results given in this paper this question can now be answered, at least in principle. **T Thiemann 1998 Class Quantum Grav.15 1281 doi:10.1088/0264-9381/15/5/012 Quantum spin dynamics (QSD): V. Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories.**

NOTATION

Module One

It is an old speculation and prognostication in physics that, once the gravitational field successfully quantized, it should serve as the natural regulator of infrared and ultraviolet singularities that plague (e) quantum field theories in a background metric.

G_{13} : Category one of quantum field theories in a background metric

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of gravitational field successfully quantized, it should serve as the natural regulator of infrared and ultraviolet singularities

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

T Thiemann demonstrates that at least part of this idea is implemented in a precise sense within the framework of (e) four-dimensional canonical Lorentzian quantum gravity in the continuum

G_{16} : Category one of gravitational field successfully quantized, it should serve as the natural regulator of infrared and ultraviolet singularities that plague quantum field theories in a background metric; four-dimensional canonical Lorentzian quantum gravity in the continuum

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of four-dimensional canonical Lorentzian quantum gravity in the continuum; gravitational field successfully quantized, it should serve as the natural regulator of infrared and ultraviolet singularities that plague quantum field theories in a background metric

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Specifically, he shows that the

Hamiltonian of the standard model supports (eb) a representation in which finite linear combinations of (e) Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operators

G_{20} : Category one of Hamiltonian of the standard model

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of representation in which finite linear combinations of (e) Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operators

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Hamiltonian of the standard model supports a representation in which finite linear combinations of (e) Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operators

G_{24} : Category one of Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operator

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Hamiltonian of the standard model supports a representation in which finite linear combinations

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Hamiltonian of the standard model supports a representation in which finite linear combinations of Wilson loop functionals around closed loops, as well as along open lines with (e&eb) fermionic and Higgs field insertions at the end points, are (=) densely defined operators

G_{28} : Category one of Hamiltonian of the standard model supports a representation in which finite linear combinations of Wilson loop functionals around closed loops, as well as along open lines; fermionic and Higgs field insertions at the end points, are (=) densely defined operators

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of fermionic and Higgs field insertions at the end points, are (=) densely defined operators ;Hamiltonian of the standard model supports a representation in which finite linear combinations of Wilson loop functionals around closed loops, as well as along open lines

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Hamiltonian of the standard model supports a representation in which finite linear combinations of Wilson loop functionals around closed loops, as well as along open lines with fermionic and Higgs field insertions at the end points, are (=) densely defined operators

G_{32} : Category one of Hamiltonian of the standard model supports a representation in which finite linear combinations of Wilson loop functionals around closed loops, as well as along open lines with fermionic and Higgs field insertions at the end points

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of densely defined operators

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

This

Hamiltonian, surprisingly, does not (e) suffer from any singularities; it is (=) completely finite without renormalization

G_{36} : Category one of suffer age from any singularities; it is (=) completely finite without renormalization

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Hamiltonian,

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Hamiltonian, surprisingly, does not suffer from any singularities; it is (=) completely finite without

renormalization

G_{40} : Category one of Hamiltonian, surprisingly, does not suffer from any singularities

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of completely finite without renormalization

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

This property is shared by string theory. In contrast to string theory, however, we are dealing with a particular

phase of the standard model coupled to (e&b) gravity which is entirely non-perturbatively defined and

second quantized

G_{44} : Category one of phase of the standard model; gravity which is entirely non-perturbatively defined and **second quantized**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of gravity which is entirely non-perturbatively defined and **second quantized**; phase of the standard model

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

42

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40}$$

43

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41}$$

44

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42}$$

45

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40}$$

46

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41}$$

47

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42}$$

48

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44}$$

49

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45}$$

50

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46}$$

51

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44}$$

52

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45}$$

53

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

55

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

56

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

57

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient

for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient

for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) \quad - (b''_{16})^{(2,2)}(G_{19}, t) \quad - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{28})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) \quad - (b''_{40})^{(8,8)}(G_{43}, t) \quad - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) \quad - (b''_{17})^{(2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8)}(G_{43}, t) \quad - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) \quad - (b''_{18})^{(2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8)}(G_{43}, t) \quad - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} (G_{19}, t) & -(b''_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18}$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} (T_{21}, t) & +(a''_{16})^{(2,2,2)}(T_{17}, t) & +(a'_{13})^{(1,1,1)}(T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} (T_{21}, t) & +(a''_{17})^{(2,2,2)}(T_{17}, t) & +(a'_{14})^{(1,1,1)}(T_{14}, t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} (T_{21}, t) & +(a''_{18})^{(2,2,2)}(T_{17}, t) & +(a'_{15})^{(1,1,1)}(T_{14}, t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a'_{20})^{(3)}(T_{21}, t)$, $+(a'_{21})^{(3)}(T_{21}, t)$, $+(a'_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b''_{16})^{(2,2,2)}(G_{19}, t) & -(b''_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b''_{17})^{(2,2,2)}(G_{19}, t) & -(b''_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b''_{18})^{(2,2,2)}(G_{19}, t) & -(b''_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} \boxed{(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(AAAAAAA) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(BBBBBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(CCCCCCC) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(DDDDDDDD)$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(EEEEEEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(FFFFFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}{}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}{}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(aa) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{(\bar{M}_{16})^{(2)} s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{-(\bar{M}_{16})^{(2)} s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as 249
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on
 (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right.$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:-

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(aa) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and

512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]}$$

517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}(s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}(s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}(s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b'_i)^{(5)}}{\partial G_j}(G_{31}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}(G_{43}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY EIGHT

Gauge Field Theory Coherent States

INTRODUCTION—VARIABLES USED

T Thiemann and O Winkler 2001 Class Quantum Grav.18 2561 doi:10.1088/0264-9381/18/14/301
Gauge field theory coherent states (GCS): II. Peakedness properties

- (1) **T Thiemann and O Winkler** apply the methods outlined in the previous paper of this series to the particular set of states obtained by choosing the complexifier to be a Laplace operator for each edge of a graph.
- (2) The corresponding coherent state transform was introduced by Hall for one edge and

generalized by Ashtekar, Lewandowski, Marolf, Mourão and Thiemann to arbitrary, finite, piecewise-analytic graphs.

- (3) However, both of these works were incomplete with respect to the following two issues. The focus was on the unitarity of the transform and left (e) the properties of the corresponding coherent states themselves untouched.
- (4) While these states depend in some sense on (eb) **complexified connections**, it remained unclear what the complexification was in terms of (e&eb) the coordinates of the underlying real phase space.
- (5) In this paper authors complement these results: first, they explicitly derive the complexification of the configuration space underlying (e&eb) these heat kernel coherent states and, secondly, prove that this family of states satisfies (eb) all the usual properties. (i) **Peakedness in the configuration, momentum and (e&eb) phase space (or Bargmann-Segal) representation.** (ii) **Saturation of (e) the unquenched Heisenberg uncertainty bound.** (iii) (Over) completeness.
- (6) These states therefore comprise (e) a candidate family for the semiclassical analysis of canonical quantum gravity and quantum gauge theory coupled to (e&eb) quantum gravity.
- (7) They also enable (eb) error-controlled approximations to difficult analytical calculations and therefore set a new starting point for numerical, semiclassical canonical quantum general relativity and gauge theory. The text is supplemented and accentuated with an appendix which contains extensive graphics in order to give a feeling for the so far unknown peakedness properties of the states constructed. **T Thiemann and O Winkler 2001 Class Quantum Grav.18 2561 doi:10.1088/0264-9381/18/14/301 Gauge field theory coherent states (GCS): II. Peakedness properties.**

NOTATION

Module One

T Thiemann and O Winkler apply the methods outlined in the previous paper of this series to the (e&eb) particular set of states obtained by (e) choosing the complexifier to be a Laplace operator for each edge of a graph

G_{13} : Category one of **Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories**; particular set of states obtained by (e) choosing the complexifier to be a Laplace operator for each edge of a graph

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of particular set of states obtained by (e) choosing the complexifier to be a Laplace operator for each edge of a graph ;**Quantum gravity as the natural regulator of the Hamiltonian constraint of matter quantum field theories**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

The corresponding coherent state transform was introduced by Hall for (e) one edge and generalized by Ashtekar, Lewandowski, Marolf, Mourão and Thiemann to arbitrary, finite, piecewise-analytic graphs

G_{16} : Category one of one edge and generalized to arbitrary, finite, piecewise-analytic graphs

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of corresponding coherent state transform

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

However, both of these works were incomplete with respect to the following two issues. The focus was on the unitarity of the transform and left (e) the properties of the corresponding coherent states themselves untouched

G_{20} : Category one of unitarity of the transform; properties of the corresponding coherent states

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of properties of the corresponding coherent states ;unitarity of the transform

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

While these states depend in some sense on (eb) **complexified connections**, it remained unclear what the complexification was in terms of (e&eb) the coordinates of the underlying real phase space

G_{24} : Category one of these states depend

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **complexified connections**, it remained unclear what the complexification was in terms of (e&eb) the coordinates of the underlying real phase space

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

While these states depend in some sense on **complexified connections**, it remained unclear what the complexification was in terms of (e&eb) the coordinates of the underlying real phase space

We assume the complexification on phase space and give model

G_{28} : Category one of complexification; coordinates of the underlying real phase space

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of coordinates of the underlying real phase space ;complexification

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

In this paper authors complement these results: first, they explicitly derive the complexification of the configuration space underlying (e&eb) these heat kernel coherent states and, secondly, prove that this

family of states satisfies (eb) all the usual properties. (i) **Peakedness in the configuration, momentum and (e&eb) phase space (or Bargmann-Segal) representation.** (ii) **Saturation of (e) the unquenched Heisenberg uncertainty bound.** (iii) (Over) completeness

G_{32} : Category one of complexification of the configuration space; heat kernel coherent states

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of heat kernel coherent states ;complexification of the configuration space

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

family of states satisfies (eb) all the usual properties. (i) **Peakedness in the configuration, momentum and (e&eb) phase space (or Bargmann-Segal) representation.** (ii) **Saturation of (e) the unquenched Heisenberg uncertainty bound.** (iii) (Over) completeness

G_{36} : Category one of family of states

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of **Peakedness in the configuration, momentum and (e&eb) phase space (or Bargmann-Segal) representation.** (ii) **Saturation of (e) the unquenched Heisenberg uncertainty bound.** (iii) (Over) completeness

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

G_{40} : Category one of **Peakedness in the configuration, momentum; phase space (or Bargmann-Segal) representation.**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **phase space (or Bargmann-Segal) representation.;Peakedness in the configuration, momentum**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Saturation of (e) the unquenched Heisenberg uncertainty bound. (iii) (Over) completeness

G_{44} : Category one of **unquenched Heisenberg uncertainty bound**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **Saturation**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ + (a''_{20})^{(3)}(T_{21}, t) &= \text{First augmentation factor} \\ - (b''_{20})^{(3)}(G_{23}, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20\end{aligned}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & - (b''_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b''_{22})^{(3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{38})^{(7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) + (a''_{24})^{(4,4)}(T_{25}, t) + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31}, t) & -(b'_{24})^{(4,4)}(G_{27}, t) & -(b'_{32})^{(6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1)}(G, t) & -(b'_{16})^{(2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31}, t) & -(b'_{25})^{(4,4)}(G_{27}, t) & -(b'_{33})^{(6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1)}(G, t) & -(b'_{17})^{(2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31}, t) & -(b'_{26})^{(4,4)}(G_{27}, t) & -(b'_{34})^{(6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b'_{32})^{(6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b'_{13})^{(1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b'_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b'_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b'_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b'_{42})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b'_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(GGGGGGG) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(HHHHHHH) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(IIIIII) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(IIIIIIII)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(KKKKKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(LLLLLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{40})^{(8)}, T_i \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(bb) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : (\widehat{G}_{31}), (\widehat{T}_{31}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widetilde{A}_{36})^{(7)} + (\widetilde{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widetilde{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widetilde{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{t \in \mathbb{R}_+} \{ \max_i |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \right.$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \right)$$

355

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)}) - (S_2)^{(9)}} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(bb) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and

512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]}$$

517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{21})^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b''_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b''_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* \mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* \mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* \mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^* \mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^* \mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)} T_{28}^* \mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)} T_{29}^* \mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)} T_{30}^* \mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY NINE

Emergent Space-Time

INTRODUCTION—VARIABLES USED

A potential foundation for emergent space-time Kevin H. Knuth^{1, a)} and Newshaw Bahreyni¹.

- (1) Kevin H. Knuth^{1,a)} and Newshaw Bahreyni¹ present a novel derivation of both the Minkowski metric and Lorentz transformations from (e) the consistent quantification of a causally ordered set of events with respect (e&eb) to an **embedded observer**.
- (2) Unlike past derivations, which have relied on assumptions such as the existence of a 4-dimensional manifold, symmetries of space-time, or the constant speed of light, **Kevin H. Knuth^{1,a)} and**

Newshaw Bahreyni demonstrate that this now familiar mathematics can be derived as (=) the unique means to consistently quantify a network of events.

- (3) This suggests (eb) that **space-time need not be physical, but instead the mathematics of space and time emerges as (=) the unique way in which an observer can consistently quantify (eb) events and their relationships to one another.**
- (4) **The result is a potential foundation for (e) emergent space-time.** A potential foundation for emergent space-time **Kevin H. Knuth**, a) and **Newshaw Bahreyni**.

Alejandro Perez 2003 Class. Quantum Grav. 20 R43 doi:10.1088/0264-9381/20/6/202 Spin foam models for quantum gravity

- (5) In this topical review, **Alejandro Perez** reviews the present status of the **spin foam formulation** of (e) non-perturbative (background-independent) quantum gravity.
- (6) The topical review is divided into two parts. In the first part, **Alejandro Perez** presents a general introduction to the main ideas emphasizing their motivation from various perspectives. Riemannian three-dimensional gravity is used as a simple example to illustrate (eb) conceptual issues and the main goals of the approach to **spin foam formulation**.
- (7) The main features of the various existing models for four-dimensional gravity are also presented here. Authors conclude with a discussion of important questions to be addressed in four dimensions (gauge invariance (e&eb) discretization independence, etc).
- (8) In the second part, Alejandro concentrates on the definition of the **Barrett–Crane model**. He presents the main results obtained in this framework from a critical perspective. Finally, they review the combinatorial formulation of spin foam models based on (e) the dual group field theory technology. Authors present the Barrett–Crane model in this framework and review the finiteness results obtained for (e) both its Riemannian and its Lorentzian variants. **Alejandro Perez 2003 Class. Quantum Grav. 20 R43 doi:10.1088/0264-9381/20/6/202 Spin foam models for quantum gravity**

NOTATION

Module One

- (1) These states therefore comprise (e) a candidate family for the semiclassical analysis of canonical quantum gravity and quantum gauge theory coupled to (e&eb) quantum gravity.
- (2) They also enable (eb) error-controlled approximations to difficult analytical calculations and therefore set a new starting point for numerical, semiclassical canonical quantum general relativity and gauge theory. The text is supplemented and accentuated with an appendix which contains extensive graphics in order to give a feeling for the so far unknown peakedness properties of the states constructed. **T Thiemann and O Winkler 2001 Class Quantum Grav.18 2561 doi:10.1088/0264-9381/18/14/301 Gauge field theory coherent states (GCS): II. Peakedness properties.**

G_{13} : Category one of candidate family for the semiclassical analysis of canonical quantum gravity and quantum gauge theory coupled to (e&eb) quantum gravity

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of states therefore comprise

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

canonical quantum gravity and quantum gauge theory coupled to (e&eb) quantum gravity

G_{16} : Category one of canonical quantum gravity and quantum gauge theory; quantum gravity

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of quantum gravity; canonical quantum gravity and quantum gauge theory

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Kevin H. Knuth^{1,a)} and Newshaw Bahreyni¹ present a novel derivation of both the

Minkowski metric and Lorentz transformations from (e) the consistent quantification of a causally ordered set of events with respect (e&eb) to an **embedded observer**

G_{20} : Category one of consistent quantification of a causally ordered set of events with respect (e&eb) to an **embedded observer**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Minkowski metric and Lorentz transformations

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Minkowski metric and Lorentz transformations from the consistent quantification of a causally ordered set of events with respect (e&eb) to an **embedded observer**

G_{24} : Category one of Minkowski metric and Lorentz transformations from the consistent quantification of a causally ordered set of events; **embedded observer**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **embedded observer**; Minkowski metric and Lorentz transformations from the consistent quantification of a causally ordered set of events

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Unlike past derivations, which have relied on assumptions such as the existence of a 4-dimensional manifold, symmetries of space-time, or the constant speed of light, **Kevin H. Knuth^{1,a} and Newshaw Bahreyni¹** demonstrate that this now familiar mathematics can be derived as (=) the unique means to consistently quantify a network of events.

G_{28} : Category one of existence of a 4-dimensional manifold, symmetries of space-time, or the constant speed of light can be derived

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of unique means to consistently quantify a network of events.

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

This suggests that space-time need not be physical, but instead the mathematics of space and time emerges as (=) the unique way in which an observer can consistently quantify (eb) events and their relationships to one another

G_{32} : Category one of existence of a 4-dimensional manifold, symmetries of space-time, or the constant speed of light can be derived as unique means to consistently quantify a network of events

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of space-time need not be physical, but instead the mathematics of space and time emerges as (=) the unique way in which an observer can consistently quantify (eb) events and their relationships to one another

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

space-time need not be physical, but instead the mathematics of space and time emerges as (=) the unique way in which an observer can consistently quantify (eb) events and their relationships to one another

G_{36} : Category one of space-time need not be physical, but instead the mathematics of space and time emerges

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of **unique way in which an observer can consistently quantify (eb) events and their relationships to one another**

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

space-time need not be physical, but instead the mathematics of space and time emerges as the unique way in which an observer can consistently quantify (eb) events and their relationships to one another

G_{40} : Category one of **space-time need not be physical, but instead the mathematics of space and time emerges as the unique way in which an observer can consistently quantify**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **events and their relationships to one another**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

The result is a potential foundation for (e) emergent space-time. A potential foundation for emergent space-time Kevin H. Knuth1, a) and Newshaw Bahreyni1

G_{44} : Category one of **space-time need not be physical, but instead the mathematics of space and time emerges as the unique way in which an observer can consistently quantify events and their relationships to one another**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **potential foundation for (e) emergent space-time**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY

Spin Foam Formulation Of Non-Perturbative (Background-Independent) Quantum Gravity

INTRODUCTION—VARIABLES USED

Alejandro Perez 2003 Class. Quantum Grav. 20 R43 doi:10.1088/0264-9381/20/6/202 Spin foam

models for quantum gravity

- (1) In this topical review, **Alejandro Perez** reviews the present status of the **spin foam formulation** of (e) non-perturbative (background-independent) quantum gravity.
- (2) The topical review is divided into two parts. In the first part, **Alejandro Perez** presents a general introduction to the main ideas emphasizing their motivation from various perspectives. Riemannian three-dimensional gravity is used as a simple example to illustrate (eb) conceptual issues and the main goals of the approach to **spin foam formulation**.
- (3) The main features of the various existing models for four-dimensional gravity are also presented here. Authors conclude with a discussion of important questions to be addressed in four dimensions (gauge invariance (e&eb) discretization independence, etc).
- (4) In the second part, Alejandro concentrates on the definition of the **Barrett–Crane model**. He presents the main results obtained in this framework from a critical perspective. Finally, they review the combinatorial formulation of spin foam models based on (e) the dual group field theory technology. Authors present the Barrett–Crane model in this framework and review the finiteness results obtained for (e) both its Riemannian and its Lorentzian variants. **Alejandro Perez 2003 Class. Quantum Grav. 20 R43 doi:10.1088/0264-9381/20/6/202 Spin foam models for quantum gravity**

T Thiemann 1998 Class Quantum Grav 15 1249 doi:10.1088/0264-9381/15/5/011 Quantum spin dynamics (QSD): IV. Euclidean quantum gravity as a model to test Lorentzian quantum gravity

- (5) The quantization of Lorentzian or Euclidean $2 + 1$ gravity by (e) canonical methods is a well studied problem.
- (6) However, the constraints of $2 + 1$ gravity are (=) those of a topological field theory and therefore resemble (e&eb) very little those of the corresponding Lorentzian $3 + 1$ constraint.
- (7) In this paper authors canonically quantize Euclidean $2 + 1$ gravity for (e) an arbitrary genus of the spacelike hypersurface with (e&eb) new, classically equivalent constraints that maximally probe the Lorentzian $3 + 1$ situation.
- (8) They choose the signature to be Euclidean because (e) this implies that the gauge group is (=) as in the $3 + 1$ case, $SU(2)$ rather than.
- (9) They employ, and carry out to full completion, the new quantization method introduced in (eb) preceding papers of this series which resulted in (eb) a finite $3 + 1$ Lorentzian quantum field theory for gravity.
- (10) The space of solutions to all constraints turns out to be (=) much larger than that obtained by traditional approaches; however, it is fully included.
- (11) Thus, by a suitable restriction of the solution space, we can recover (eb) all former results which give confidence in the new quantization methods. The meaning of the remaining 'spurious solutions' is discussed.

NOTATION

Module One

In this topical review, **Alejandro Perez** reviews the present status of the **spin foam formulation** of (e) non-perturbative (background-independent) quantum gravity

G_{13} : Category one of non-perturbative (background-independent) quantum gravity

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of spin foam formulation

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

The topical review is divided into two parts. In the first part, **Alejandro Perez** presents a general introduction to the main ideas emphasizing their motivation from various perspectives.

Riemannian three-dimensional gravity is used as a simple example to illustrate (eb) conceptual issues and the main goals of the approach to spin foam formulation

G_{16} : Category one of Riemannian three-dimensional gravity is used as a simple example

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of goals of the approach to spin foam formulation

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

The main features of the various existing models for four-dimensional gravity are also presented here.

Authors conclude with a discussion of important questions to be addressed in four dimensions (gauge invariance (e&eb) discretization independence, etc)

G_{20} : Category one of gauge invariance; discretization independence

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of discretization independence ;gauge invariance

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

In the second part, Alejandro concentrates on the definition of the Barrett–Crane model. He presents the main results obtained in this framework from a critical perspective. Finally, they review the

combinatorial formulation of spin foam models based on (e) the dual group field theory technology.

Authors present the Barrett–Crane model in this framework and review the finiteness results obtained for (e) both its Riemannian and its Lorentzian variants. **Alejandro Perez 2003 Class. Quantum Grav. 20 R43**

doi:10.1088/0264-9381/20/6/202 Spin foam models for quantum gravity

G_{24} : Category one of dual group field theory technology

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of combinatorial formulation of spin foam models

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The

quantization of Lorentzian or Euclidean $2 + 1$ gravity by (e) canonical methods

is a well studied problem

G_{28} : Category one of canonical methods

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of quantization of Lorentzian or Euclidean $2 + 1$ gravity

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

However, the

constraints of $2 + 1$ gravity are (=) those of a topological field theory and therefore resemble (e&eb) very little those of the corresponding Lorentzian $3 + 1$ constraint

G_{32} : Category one of constraints of $2 + 1$ gravity

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of topological field theory and therefore resemble (e&eb) very little those of the corresponding Lorentzian $3 + 1$ constraint

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

constraints of $2 + 1$ gravity are those of a topological field theory and therefore resemble (e&eb) very little

those of the corresponding Lorentzian 3 + 1 constraint

G_{36} : Category one of constraints of 2 + 1 gravity are those of a topological field theory; corresponding Lorentzian 3 + 1 constraint

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of corresponding Lorentzian 3 + 1 constraint ;constraints of 2 + 1 gravity are those of a topological field theory

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

In this paper authors

canonically quantization of Euclidean 2 + 1 gravity for (e) an arbitrary genus of the spacelike hypersurface with (e&eb) new, classically equivalent constraints that maximally probe the Lorentzian 3 + 1 situation

G_{40} : Category one of arbitrary genus of the spacelike hypersurface with (e&eb) new, classically equivalent constraints that maximally probe the Lorentzian 3 + 1 situation

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of canonically quantization of Euclidean 2 + 1 gravity

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

canonically quantization of Euclidean 2 + 1 gravity for an arbitrary genus of the spacelike hypersurface with (e&eb) new, classically equivalent constraints that maximally probe the Lorentzian 3 + 1 situation

G_{44} : Category one of canonically quantization of Euclidean 2 + 1 gravity for an arbitrary genus of the spacelike hypersurface; classically equivalent constraints that maximally probe the Lorentzian 3 + 1 situation

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of classically equivalent constraints that maximally probe the Lorentzian 3 + 1 situation ;canonically quantization of Euclidean 2 + 1 gravity for an arbitrary genus of the spacelike hypersurface

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \end{aligned}$$

$$\begin{aligned}\frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

12

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$- (b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+ (a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$- (b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+ (a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$- (b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\begin{aligned}\frac{dG_{32}}{dt} &= (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} & 31 \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} & 32 \\ \frac{dG_{34}}{dt} &= (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} & 33 \\ \frac{dT_{32}}{dt} &= (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} & 34 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} & 35 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} & 36 \\ &+ (a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor} \\ &\text{Module Numbered Seven:}\end{aligned}$$

The differential system of this model is now (Seventh Module)

$$\begin{aligned}\frac{dG_{36}}{dt} &= (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} & 42 \\ &+ (a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \\ &\text{Module Numbered Eight}\end{aligned}$$

The differential system of this model is now

$$\begin{aligned}\frac{dG_{40}}{dt} &= (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} & 44 \\ \frac{dG_{42}}{dt} &= (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} & 54 \\ &+ (a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor} \\ &- (b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}\end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a'_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a'_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a'_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t), +(a''_{21})^{(3)}(T_{21}, t), +(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{l} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{l} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{l} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{l} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{l} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients

for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for

category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients

for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition

coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition

coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)} G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41}, t)} & \boxed{+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41}, t)} & \boxed{+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30$$

122

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

123

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(MMMMMM) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(NNNNNNN) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(OOOOOOO) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(PPPPPPP)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(QQQQQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(RRRRRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42$$

136

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$$

138

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)}$$

139

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)}$$

140

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)}$$

141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t}$$

142

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$$

143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(cc) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$\left(1 + (a_{36})^{(7)} t \right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 180

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} =$$

$$\left(1 + (a_{40})^{(8)} t \right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t \right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}\right), \left((G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})$ it follows $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), ((G_{27})^{(2)}, (T_{27})^{(2)}) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}\right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} &((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t(t), t)) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{-(\hat{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\hat{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\hat{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\tilde{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} ((b_{38}'')^{(7)}((G_{39})(t), t)) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\text{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\bar{G}_{40}^{(1)} - \bar{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} & (\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ & -(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ & -(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} & \text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ & (m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ & (m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} & (\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ & (\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

$$\text{roots of the equations } (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-
$$v^{(2)} = \frac{G_{16}}{G_{17}}$$
 388

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines $(\nu_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{[-(a_{41})^{(8)}((\nu_1)^{(8)} - (\nu_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((\nu_1)^{(8)} - (\nu_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get

422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (\bar{C})^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (\bar{C})^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(cc) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}) T_{13}^* \mathbb{G}_j \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}) T_{14}^* \mathbb{G}_j \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}) T_{15}^* \mathbb{G}_j \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}(G_{47}^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY ONE

Signature To Be Euclidean

INTRODUCTION—VARIABLES USED

T Thiemann 1998 Class Quantum Grav 15 1249 doi:10.1088/0264-9381/15/5/011 Quantum spin dynamics (QSD): IV. Euclidean quantum gravity as a model to test Lorentzian quantum gravity

- (1) They choose the signature to be Euclidean because (e) this implies that the gauge group is (=) as in the 3 + 1 case, SU (2) rather than.
- (2) They employ, and carry out to full completion, the new quantization method introduced in (eb) preceding papers of this series which resulted in (eb) a finite 3 + 1 Lorentzian quantum field theory for gravity.
- (3) The space of solutions to all constraints turns out to be (=) much larger than that obtained by traditional approaches; however, it is fully included.
- (4) Thus, by a suitable restriction of the solution space, we can recover (eb) all former results which give confidence in the new quantization methods. The meaning of the remaining 'spurious solutions' is discussed.

NOTATION

Module One

They choose the signature to be Euclidean because this implies that the gauge group is (=) as in the 3 + 1 case, SU (2) rather than.

G_{13} : Category one of signature to be Euclidean is chosen because by implication

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of gauge group is (=) as in the 3 + 1 case, SU (2)

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

They employ, and carry out to full completion, the new quantization method introduced in (eb) preceding

papers of this series which resulted in (eb) a finite 3 + 1 Lorentzian quantum field theory for gravity

G_{16} : Category one of employment and completion of new quantization method

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of finite 3 + 1 Lorentzian quantum field theory for gravity

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

The

space of solutions to all constraints turns out to be (=) much larger than that obtained by traditional approaches; however, it is fully included

G_{20} : Category one of space of solutions to all constraints

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of much larger than that obtained by traditional approaches; however, it is fully included

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Thus, by a suitable restriction of the solution space, we can recover (eb) all former results which give confidence in the new quantization methods.

The meaning of the remaining 'spurious solutions' is discussed.

G_{24} : Category one of suitable restriction of the solution space

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of all former results which give confidence in the new quantization methods.

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Massive Dirac fermions and spin physics in an ultrathin film of topological insulator Phys. Rev. B 81, 115407 – Published 3 March 2010 Hai-Zhou Lu, Wen-Yu Shan, Wang Yao, Qian Niu, and Shun-Qing

Shen

- (1) Authors study transport and optical properties of the surface states, which lie in (eb) the bulk energy gap of a thin-film topological insulator.
- (2) When the film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas.
- (3) Spin-dependent physics emerges in (eb) the surface bands, which are (=) vastly different from the bulk behavior. Spin-dependent physics include (e) the surface spin Hall effects, spin-dependent orbital magnetic moment, and spin-dependent optical transition selection rule, which allow (eb) optical spin injection.
- (4) They show a topological quantum phase transition where the Chern number of the surface bands changes (e&eb) when varying the thickness of the thin film. DOI: <http://dx.doi.org/10.1103/PhysRevB.81.115407>

Authors study

transport and optical properties of the surface states, which lie in (eb) the bulk energy gap of a thin-film topological insulator

G_{28} : Category one of transport and optical properties of the surface states

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of bulk energy gap of a thin-film topological insulator

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

When the

film thickness is comparable with (e&eb) the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas

G_{32} : Category one of film thickness; surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas ;film thickness

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas

G_{36} : Category one of film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top; bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas ;film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas

G_{40} : Category one of film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of energy gap and form (eb) two degenerate massive Dirac hyperbolas

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap and form (eb) two degenerate massive Dirac hyperbolas

G_{44} : Category one of film thickness is comparable with the surface-state decay length into the bulk, the tunneling between the top and bottom surfaces opens an energy gap

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of two degenerate massive Dirac hyperbolas

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \end{aligned}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor}$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t), +(a''_{21})^{(3)}(T_{21}, t), +(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & +(a''_{24})^{(4,4)}(T_{25}, t) & +(a''_{32})^{(6,6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{c} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients

for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for

category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients

for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition

coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition

coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$\begin{aligned} & - \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86 \end{aligned}$$

$$\begin{aligned} \frac{dG_{34}}{dt} &= (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87 \end{aligned}$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third

augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth

augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31} - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(SSSSSS) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(TTTTTTTT) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(UUUUUUUU) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(VVVVVVVV)$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - G_{39}| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(WWWWWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(XXXXXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = 169$$

$$\left(1 + (a_{16})^{(2)} t \right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t \right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(dd) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 180

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)} \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\bar{A}_{16})^{(2)} + (\bar{P}_{16})^{(2)} (\bar{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)} \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\bar{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad \text{205}$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as 227
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{-(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} | (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(\hat{M}_{32})^{(6)} s_{(32)}} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} \right) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} ((b_{38}'')^{(7)}((G_{39})(t), t)) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

266

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)}$$

267

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)}$$

268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\}$$

269

Indeed if we denote

270

Definition of $(\bar{G}_{43}), (\bar{T}_{43})$: $(\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results

271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned}$$

273

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

274

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)})_1 \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations}$$

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(\mu_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(\mu_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} [e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)}) - (S_2)^{(6)}} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-
$$v^{(2)} = \frac{G_{16}}{G_{17}}$$
 388

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)} :-$
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)} :-$

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t) :-$

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines $(\nu_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{[-(a_{41})^{(8)}((\nu_1)^{(8)} - (\nu_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((\nu_1)^{(8)} - (\nu_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get

422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t) \right) - (a_{45}'')^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(dd) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]}, \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}, \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]}, \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]}, \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]}, \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]}, \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_{47}'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ \left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \right. \\
 &\left. \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ \left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 &+ \left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \right. \\
 &\left. \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY TWO

Topological Strings, And Supersymmetric Gauge Theories

INTRODUCTION—VARIABLES USED

Nuclear Physics B Volume 644, Issues 1–2, 11 November 2002, Pages 3–20 Matrix models, topological strings, and supersymmetric gauge theories Robbert Dijkgraafa Cumrun Vafa doi:10.1016/S0550-3213(02)00766-6.

- (1) **Robbert Dijkgraafa Cumrun Vafa** show that B-model topological strings on (eb) local Calabi–Yau threefolds are (=) large-N duals of matrix models, which in the planar limit naturally give rise to (eb) special geometry.
- (2) These matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by (e) deforming $\mathcal{N} = 2$ theories by (e) superpotential term that can be directly identified with the potential of the matrix model.
- (3) Moreover by tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models coupled to (e&eb) 2d gravity, thereby relating non-critical string theories to (e&eb) type II superstrings on Calabi–Yau backgrounds. Copyright © 2002 Published by Elsevier B.V. **Nuclear Physics B Volume 644, Issues 1–2, 11 November 2002, Pages 3–20 Matrix models, topological strings, and supersymmetric gauge theories Robbert Dijkgraafa Cumrun Vafa doi:10.1016/S0550-3213(02)00766-6.**

T W B Kibble 1976 J. Phys A: Math. Gen. 9 1387 doi:10.1088/0305-4470/9/8/029 Topology of cosmic domains and strings

- (4) The possible domain structures which can arise in (eb) the universe in a spontaneously broken gauge theory are studied.
- (5) It is shown that the formation of domain wall, strings or monopoles depends on (=) the homotopy groups of the manifold of degenerate vacua.
- (6) The subsequent evolution of these structures is investigated. It is argued that while theories generating domain walls can probably be eliminated (because of their unacceptable gravitational effects), a cosmic network of strings may well have been formed and may have had important cosmological effects. **T W B Kibble 1976 J. Phys A: Math. Gen. 9 1387 doi:10.1088/0305-4470/9/8/029 Topology of cosmic domains and strings**

NOTATION

Module One

Robbert Dijkgraafa Cumrun Vafa show that

B-model topological strings on (eb) local Calabi–Yau threefolds are (=) large-N duals of matrix models, which in the planar limit naturally give rise to (eb) special geometry.

G_{13} : Category one of B-model topological strings

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of local Calabi–Yau threefolds are (=) large-N duals of matrix models, which in the planar limit naturally give rise to (eb) special geometry

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

B-model topological strings on local Calabi–Yau threefolds are (=) large-N duals of matrix models, which in the planar limit naturally give rise to (eb) special geometry

G_{16} : Category one of B-model topological strings on local Calabi–Yau threefolds

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of large-N duals of matrix models, which in the planar limit naturally give rise to (eb) special geometry

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

B-model topological strings on local Calabi–Yau threefolds are large-N duals of matrix models, which in the planar limit naturally give rise to (eb) special geometry

G_{20} : Category one of B-model topological strings on local Calabi–Yau threefolds are large-N duals of matrix models, which in the planar limit

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of special geometry

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

These

matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by (e) deforming $\mathcal{N} = 2$ theories by (e) superpotential term that can be directly identified with the potential of the matrix model

G_{24} : Category one of deforming $\mathcal{N} = 2$ theories by (e) superpotential term that can be directly identified with the potential of the matrix model

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by deforming $\mathcal{N} = 2$ theories by (e) superpotential term that can be directly identified with the potential of the matrix model

G_{28} : Category one of superpotential term that can be directly identified with the potential of the matrix model

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by deforming $\mathcal{N} = 2$ theories

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by deforming $\mathcal{N} = 2$ theories by superpotential term that can be directly identified with the potential of the matrix model

G_{32} : Category one of matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by deforming $\mathcal{N} = 2$ theories by superpotential term; potential of the matrix model

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of potential of the matrix model; matrix models directly compute F-terms in an associated $\mathcal{N} = 1$ supersymmetric gauge theory, obtained by deforming $\mathcal{N} = 2$ theories by superpotential term

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Moreover by

tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models coupled to (e&eb) 2d gravity, thereby relating non-critical string theories to (e&eb) type II superstrings on Calabi–Yau backgrounds.

Copyright © 2002 Published by Elsevier B.V. **Nuclear Physics B Volume 644, Issues 1–2, 11 November 2002, Pages 3–20 Matrix models, topological strings, and supersymmetric gauge theories Robbert Dijkgraaf Cumrun Vafa doi:10.1016/S0550-3213(02)00766-6.**

G_{36} : Category one of tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models; 2d gravity, thereby relating non-critical string theories to (e&eb) type II superstrings on Calabi–Yau backgrounds

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of 2d gravity, thereby relating non-critical string theories to (e&eb) type II superstrings on Calabi–Yau backgrounds ;tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models coupled to 2d gravity, thereby relating non-critical string theories to (e&eb) type II superstrings on Calabi–Yau backgrounds.

G_{40} : Category one of tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models coupled to 2d gravity, thereby relating non-critical string theories; **type II superstrings on Calabi–Yau backgrounds.**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **type II superstrings on Calabi–Yau backgrounds**; tuning some of the parameters of the geometry in a double scaling limit we recover (p,q) conformal minimal models coupled to 2d gravity, thereby relating non-critical string theories

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

The

possible domain structures which can arise in (eb) the universe in a spontaneously broken gauge theory
 are studied

G_{44} : Category one of possible domain structures

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of universe in a spontaneously broken gauge theory

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ + (a''_{20})^{(3)}(T_{21}, t) &= \text{First augmentation factor} \\ - (b''_{20})^{(3)}(G_{23}, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20 \\ \frac{dG_{26}}{dt} &= (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} & 21 \\ \frac{dT_{24}}{dt} &= (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} & 22 \\ \frac{dT_{25}}{dt} &= (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} & 23\end{aligned}$$

$$\begin{aligned}\frac{dT_{26}}{dt} &= (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \\ + (a''_{24})^{(4)}(T_{25}, t) &= \text{First augmentation factor} \\ - (b''_{24})^{(4)}((G_{27}), t) &= \text{First detritions factor}\end{aligned}$$

24

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\begin{aligned}\frac{dG_{28}}{dt} &= (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} & 25 \\ \frac{dG_{29}}{dt} &= (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} & 26 \\ \frac{dG_{30}}{dt} &= (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} & 27 \\ \frac{dT_{28}}{dt} &= (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} & 28 \\ \frac{dT_{29}}{dt} &= (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} & 29 \\ \frac{dT_{30}}{dt} &= (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} & 30 \\ + (a''_{28})^{(5)}(T_{29}, t) &= \text{First augmentation factor} \\ - (b''_{28})^{(5)}((G_{31}), t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\begin{aligned}\frac{dG_{32}}{dt} &= (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} & 31 \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} & 32 \\ \frac{dG_{34}}{dt} &= (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} & 33 \\ \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} & 34 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} & 35 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} & 36 \\ + (a''_{32})^{(6)}(T_{33}, t) &= \text{First augmentation factor}\end{aligned}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\begin{aligned}\frac{dG_{36}}{dt} &= (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} & 42 \\ + (a''_{36})^{(7)}(T_{37}, t) &= \text{First augmentation factor}\end{aligned}$$

Module Numbered Eight

The differential system of this model is now

$$\begin{aligned}\frac{dG_{40}}{dt} &= (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} & 44\end{aligned}$$

$$\begin{aligned}\frac{dG_{42}}{dt} &= (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} & 54 \\ &+ (a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor} \\ &- (b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}\end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a'_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a'_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a'_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3
 $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth
augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{c} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{c} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $[-(b''_{13})^{(1)}(G, t)]$, $[-(b''_{14})^{(1)}(G, t)]$, $[-(b''_{15})^{(1)}(G, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{16})^{(2,2)}(G_{19}, t)]$, $[-(b''_{17})^{(2,2)}(G_{19}, t)]$, $[-(b''_{18})^{(2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{20})^{(3,3)}(G_{23}, t)]$, $[-(b''_{21})^{(3,3)}(G_{23}, t)]$, $[-(b''_{22})^{(3,3)}(G_{23}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4,4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b''_{28})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b''_{29})^{(5,5,5,5)}(G_{31}, t)]$, $[-(b''_{30})^{(5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b''_{36})^{(7,7)}(G_{39}, t)]$, $[-(b''_{37})^{(7,7)}(G_{39}, t)]$, $[-(b''_{38})^{(7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b''_{40})^{(8,8)}(G_{43}, t)]$, $[-(b''_{41})^{(8,8)}(G_{43}, t)]$, $[-(b''_{42})^{(8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$, $[-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17}, t)] \quad [+(a''_{13})^{(1,1)}(T_{14}, t)] \quad [+(a''_{20})^{(3,3,3)}(T_{21}, t)] \\ [+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)] \quad [+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)] \quad [+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)] \\ [+(a''_{36})^{(7,7,7)}(T_{37}, t)] \quad [+(a''_{40})^{(8,8,8)}(T_{41}, t)] \quad [+(a''_{44})^{(9,9)}(T_{45}, t)] \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & - (b''_{18})^{(2)}(G_{19}, t) & - (b''_{15})^{(1,1)}(G, t) & - (b''_{22})^{(3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{38})^{(7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b'_{16})^{(2)}(G_{19}, t)$, $-(b'_{17})^{(2)}(G_{19}, t)$, $-(b'_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)}(G, t)$, $-(b'_{14})^{(1,1)}(G, t)$, $-(b'_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category

1,2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients

for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}, t) & +(a''_{28})^{(5,5)}(T_{29}, t) & +(a''_{32})^{(6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a'_{29})^{(5,5)}(T_{29}, t) & + (a'_{33})^{(6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1)}(T_{14}, t) & + (a'_{17})^{(2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a'_{30})^{(5,5)}(T_{29}, t) & + (a'_{34})^{(6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t), (a'_{25})^{(4)}(T_{25}, t), (a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a'_{28})^{(5,5)}(T_{29}, t), +(a'_{29})^{(5,5)}(T_{29}, t), +(a'_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6)}(T_{33}, t), +(a'_{33})^{(6,6)}(T_{33}, t), +(a'_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1)}(T_{14}, t), +(a'_{14})^{(1,1,1,1)}(T_{14}, t), +(a'_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2)}(T_{17}, t), +(a'_{17})^{(2,2,2,2)}(T_{17}, t), +(a'_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3)}(T_{21}, t), +(a'_{21})^{(3,3,3,3)}(T_{21}, t), +(a'_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a'_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a'_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a'_{46})^{(9,9,9,9)}(T_{45}, t), +(a'_{45})^{(9,9,9,9)}(T_{45}, t), +(a'_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b'_{32})^{(6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1)}(G, t) & - (b'_{16})^{(2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b'_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b'_{33})^{(6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1)}(G, t) & - (b'_{17})^{(2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b'_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b'_{34})^{(6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1)}(G, t) & - (b'_{18})^{(2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b'_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a''_{25})^{(4,4)}(T_{25}, t) & + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1,2,3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1,2,3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{c} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{c} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation

coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{c} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) - (b''_{28})^{(5,5,5)}(G_{31}, t) - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{c} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) - (b''_{29})^{(5,5,5)}(G_{31}, t) - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - \left[\begin{array}{c} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) - (b''_{30})^{(5,5,5)}(G_{31}, t) - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients

for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(YYYYYY) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(ZZZZZZ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(AAAAA) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(BBBBBB) \lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(CCCCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(DDDDDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ee) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[- \int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-

$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(ee) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b_{20}')^{(3)}(b_{21}')^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b_{20}')^{(3)}(b_{21}'')^{(3)}(G_{23}) + (b_{21}')^{(3)}(b_{20}'')^{(3)}(G_{23})] + (b_{20}'')^{(3)}(G_{23})(b_{21}'')^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b_{24}')^{(4)}(b_{25}')^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{21})^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY THREE

Topology Of Cosmic Domains And Strings

INTRODUCTION—VARIABLES USED

T W B Kibble 1976 J. Phys A: Math. Gen. 9 1387 doi:10.1088/0305-4470/9/8/029 Topology of cosmic domains and strings

- (1) It is shown that the formation of domain wall, strings or monopoles depends on (=) the homotopy groups of the manifold of degenerate vacua.
- (2) The subsequent evolution of these structures is investigated. It is argued that while theories generating domain walls can probably be eliminated (because of their unacceptable gravitational

effects), a cosmic network of strings may well have been formed and may have had important cosmological effects. **T W B Kibble 1976 J. Phys A: Math. Gen. 9 1387 doi:10.1088/0305-4470/9/8/029 Topology of cosmic domains and strings**

Horava-Lifshitz Gravity and Effective Theory of the Fractional Quantum Hall Effect Chaolun Wu, Shao-Feng Wu

- (3) Authors show that Horava-Lifshitz gravity theory can be employed as a covariant framework to build an (eb) effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and (e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry.
- (4) The key to this formalism is a set of correspondence relations that maps all the field degrees of freedom in the Horava-Lifshitz gravity theory to (e&eb) external background (source) fields among others in the effective action of the quantum Hall effect, according to their symmetry transformation properties.
- (5) They originally derive the map as a holographic dictionary, but its form is (=) independent of the existence of holographic duality.
- (6) This paves the way for the application of Horava-Lifshitz holography on (e&eb) fractional quantum Hall effect.
- (7) Using the simplest holographic Chern-Simons model, they compute (eb) the low energy effective action at leading orders and show that it captures (e) universal electromagnetic and geometric properties of Quantum Hall States, including the Wen-Zee shift, Hall viscosity, angular momentum density and their relations.
- (8) They identify the shift function in Horava-Lifshitz gravity theory as (=) minus of guiding center velocity and conjugate to (e) guiding center momentum.
- (9) This enables us to distinguish guiding center angular momentum density from the internal one, which is the sum of Landau orbit spin and intrinsic (topological) spin of the composite particles. Our effective action shows that Hall viscosity is minus half of the internal angular momentum density and proportional to Wen-Zee shift, and Hall bulk viscosity is half of the guiding center angular momentum density. arXiv:1409.1178 [hep-th]

NOTATION

Module One

It is shown that the

formation of domain wall, strings or monopoles depends on (=) the homotopy groups of the manifold of degenerate vacua

G_{13} : Category one of formation of domain wall, strings or monopoles

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of homotopy groups of the manifold of degenerate vacua

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

The subsequent evolution of these structures is investigated. It is argued that while theories generating domain walls can probably be eliminated (because of their unacceptable gravitational effects), a cosmic network of strings may well have been formed and may have had important cosmological effects. **T W B Kibble 1976 J. Phys A: Math. Gen. 9 1387 doi:10.1088/0305-4470/9/8/029 Topology of cosmic domains and strings**

G_{16} : Category one of theories generating domain walls; cosmic network of strings

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of cosmic network of strings; theories generating domain walls

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Authors show that

Horava-Lifshitz gravity theory can be employed as a covariant framework to build an (eb) effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and (e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

G_{20} : Category one of Horava-Lifshitz gravity theory can be employed as a covariant framework

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and (e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects (eb) all the spacetime symmetries such as non-relativistic diffeomorphism invariance and (e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

G_{24} : Category one of Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of all the spacetime symmetries such as non-relativistic diffeomorphism invariance and

(e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects all the spacetime symmetries such as non-relativistic diffeomorphism invariance and (e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

G_{28} : Category one of Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects all the spacetime symmetries such as non-relativistic diffeomorphism invariance; anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of anisotropic Weyl invariance as well as (e&eb) the gauge symmetry ;Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects all the spacetime symmetries such as non-relativistic diffeomorphism invariance

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Horava-Lifshitz gravity theory can be employed as a covariant framework to build an effective field theory for the fractional quantum Hall effect that respects all the spacetime symmetries such as non-relativistic diffeomorphism invariance and (e&eb) anisotropic Weyl invariance as well as (e&eb) the gauge symmetry

G_{32} : Category one of anisotropic Weyl invariance; gauge symmetry

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of gauge symmetry; anisotropic Weyl invariance

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

G_{36} : Category one of non-relativistic diffeomorphism invariance ; gauge symmetry

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of gauge symmetry ;non-relativistic diffeomorphism invariance

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

They originally derive the map as a holographic dictionary, but its form is (=) independent of the existence of holographic duality

G_{40} : Category one of form of map as a holographic dictionary

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of independent of the existence of holographic duality

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Using the simplest holographic Chern-Simons model, they compute (eb) the low energy effective action at leading orders and show that it captures (e) universal electromagnetic and geometric properties of Quantum Hall States, including the Wen-Zee shift, Hall viscosity, angular momentum density and their relations

G_{44} : Category one of Wen-Zee shift; Hall viscosity

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Hall viscosity ;Wen-Zee shift

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$

$$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$$

are Accentuation coefficients

$$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, \\ (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}, \\ (a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, \\ (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}, \\ (a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}, \\ (a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}, \\ (a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor} \end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \end{aligned}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor} \quad 55$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 56$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 57$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for

category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for

1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for

category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17}, t)$, $+(a''_{17})^{(2)}(T_{17}, t)$, $+(a''_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1)}(G, t)} & \boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1)}(G, t)} & \boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b'_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{46})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a'_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a'_{21})^{(3)}(T_{21}, t)} & \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} \boxed{+(a'_{22})^{(3)}(T_{21}, t)} & \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a'_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \quad 69$$

$\boxed{+(a'_{20})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3)}(T_{21}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)}$ are second augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$ are third augmentation coefficients

for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} \boxed{(b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} \boxed{(b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23}, t)}} & \boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{36})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7,7,7,7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7,7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25}, t) & +(a''_{28})^{(5,5)}(T_{29}, t) & +(a''_{32})^{(6,6)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{cccc} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25}, t) & +(a''_{29})^{(5,5)}(T_{29}, t) & +(a''_{33})^{(6,6)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{cccc} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25}, t) & +(a''_{30})^{(5,5)}(T_{29}, t) & +(a''_{34})^{(6,6)}(T_{33}, t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14}, t) & +(a''_{18})^{(2,2,2,2)}(T_{17}, t) & +(a''_{22})^{(3,3,3,3)}(T_{21}, t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} \boxed{(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} + \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35}) - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(EEEEEEEE) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(FFFFFFF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(GGGGGGGG) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(HHHHHHHH)$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(IIIIIIII) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(JJJJJJJJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0} \right)} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\left(\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0} \right)} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ff) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$| (G_{19})^{(1)} - (G_{19})^{(2)} | e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e \left[- \int_0^t \{ (a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right] \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} | (a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as 249
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on
 (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t}$$

376

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq$$

377

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (\bar{C})^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (\bar{C})^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(ff) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

$$\text{ASYMPTOTIC STABILITY ANALYSIS} \quad 531$$

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}(s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}(s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}(s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY FOUR

Condition Of Phase-Conjugate-Adaptive-Resonance Is Necessary To Completely Specify The Act Of Perception

INTRODUCTION—VARIABLES USED

National Institute for Discovery Science: Nature's Mind: the Quantum Hologram by Edgar Mitchell,
Ph.D. Nature's Mind: the Quantum Hologram Fax: 561-641-5242, edgarmitchell@msn.com

- (1) These different types of mind/mind, mind/matter experiments have been rigorously and routinely conducted for decades with statistically compelling results but just as routinely dismissed or ignored by main stream science because the implications of non-local action are so foreign to the

classical paradigm.

- (2) However, if we consider that the **condition of phase-conjugate-adaptive-resonance is necessary to completely specify the act of perception** as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as locked in a resonant feedback loop.
- (3) The incoming wave front carrying information may be labeled as "perception" from the point of view of the percipient, and the return path required by the resonant relationship may be labeled "attention" (or for subsequent discussion, "intention").
- (4) It is a well established principle in the meditative practices of esoteric disciplines that prolonged focused attention on a object of meditation causes the percipient and the object to appear to merge so that a deeper level of information about the object is obtained; information such as history or internal functioning, that would not be available through classical space/time information.
- (5) The concept of the quantum hologram adequately and completely describes how this phenomenon might take place
- (6) Further, it is accepted that the mind/brain is a massively parallel processor, capable of performing many tasks simultaneously and subconsciously (in the right, intuitive part of the brain).
- (7) Attention (meaning conscious, focused attention) is a unique and singular task that must take place sequentially, mostly in the left cognitive part of the brain.
- (8) The condition of attention deficit disorder (ADD) is precisely the problem of a percipient being unable to maintain a singular focus for sufficient time to complete a desired task or observation.
- (9) Thus, the action of focusing attention by a percipient may be construed as a necessary condition for pcar to be established with the perceived object.

NOTATION

Module One

However, if we consider that the

condition of phase-conjugate-adaptive-resonance is necessary to (e) completely specify the act of perception as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as locked in a resonant feedback loop

G_{13} : Category one of **condition of phase-conjugate-adaptive-resonance is necessary**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **completely specify the act of perception** as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as locked in a resonant feedback loop

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

condition of phase-conjugate-adaptive-resonance is necessary to completely specify the act of perception as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as (=) locked in a resonant feedback loop

G_{16} : Category one of **condition of phase-conjugate-adaptive-resonance is necessary to completely**

specify the act of perception as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of locked in a resonant feedback loop

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

condition of phase-conjugate-adaptive-resonance is necessary to completely specify the act of perception as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as locked in a resonant feedback loop

G_{20} : Category one of perceived object; percipient's perceptual system

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of percipient's perceptual system; perceived object

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

condition of phase-conjugate-adaptive-resonance is necessary to completely specify the act of perception as described in the mathematical formalism of the non-local quantum hologram by Marcer, then we may also consider the perceived object and the percipient's perceptual system as locked in a resonant feedback loop

G_{24} : Category one of **complete specification of the act of perception**; percipient's perceptual system

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of percipient's perceptual system ; **complete specification of the act of perception**

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

G_{28} : Category one of percipient's perceptual system; **condition of phase-conjugate-adaptive-resonance**

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of **condition of phase-conjugate-adaptive-resonance**; percipient's perceptual system

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

The

incoming wave front carrying information may be labeled as (=) "perception" from the point of view of the percipient, and the return path required by the resonant relationship may be labeled "attention" (or for subsequent discussion, "intention").

G_{32} : Category one of incoming wave front carrying information

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of "perception" from the point of view of the percipient

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

incoming wave front carrying information may be labeled as (=) "perception" from the point of view of the percipient, and the

return path required by the resonant relationship may be labeled "attention" (or for subsequent discussion, "intention").

G_{36} : Category one of return path required by the resonant relationship

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of "attention" (or for subsequent discussion, "intention").

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

It is a well established principle in the meditative practices of esoteric disciplines that

prolonged focused attention on a object of meditation causes (eb) the percipient and the object to appear to merge so that a deeper level of information about the object is obtained formation such as history or internal

functioning, that would not be available through classical space/time information

G_{40} : Category one of prolonged focused attention on a object of meditation

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of percipient and the object to appear to merge so that a deeper level of information about the object is obtained

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

prolonged focused attention on a object of meditation causes (eb) the **percipient** and the object to appear to merge so that a deeper level of information about the object is obtained

G_{44} : Category one of **percipient**; object

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **object** ;**percipient**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ + (a''_{20})^{(3)}(T_{21}, t) &= \text{First augmentation factor} \\ - (b''_{20})^{(3)}(G_{23}, t) &= \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\begin{aligned}\frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} & 19 \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} & 20\end{aligned}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)}[-(b''_{13})^{(1)}(G, t)] \quad [-(b''_{16})^{(2,2)}(G_{19}, t)] \quad [-(b''_{20})^{(3,3)}(G_{23}, t)] \\ [-(b''_{24})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{28})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{32})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{36})^{(7,7)}(G_{39}, t)] \quad [-(b''_{40})^{(8,8)}(G_{43}, t)] \quad [-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)}[-(b''_{14})^{(1)}(G, t)] \quad [-(b''_{17})^{(2,2)}(G_{19}, t)] \quad [-(b''_{21})^{(3,3)}(G_{23}, t)] \\ [-(b''_{25})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{29})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{33})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{37})^{(7,7)}(G_{39}, t)] \quad [-(b''_{41})^{(8,8)}(G_{43}, t)] \quad [-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)}[-(b''_{15})^{(1)}(G, t)] \quad [-(b''_{18})^{(2,2)}(G_{19}, t)] \quad [-(b''_{22})^{(3,3)}(G_{23}, t)] \\ [-(b''_{26})^{(4,4,4,4)}(G_{27}, t)] \quad [-(b''_{30})^{(5,5,5,5)}(G_{31}, t)] \quad [-(b''_{34})^{(6,6,6,6)}(G_{35}, t)] \\ [-(b''_{38})^{(7,7)}(G_{39}, t)] \quad [-(b''_{42})^{(8,8)}(G_{43}, t)] \quad [-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)] \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{36})^{(7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7)}(T_{37}, t)$, $+(a'_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{40})^{(8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a'_{44})^{(9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b'_{20})^{(3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b'_{21})^{(3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{37})^{(7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b'_{22})^{(3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b'_{38})^{(7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9)}(G_{47}, t) \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a'_{16})^{(2,2,2)}(T_{17}, t) + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b'_{16})^{(2,2,2)}(G_{19}, t) & -(b'_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b'_{17})^{(2,2,2)}(G_{19}, t) & -(b'_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b'_{18})^{(2,2,2)}(G_{19}, t) & -(b'_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) + (a''_{24})^{(4,4)}(T_{25}, t) + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} & + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) & \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} & - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} & - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} \boxed{(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T_{29}'| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} \|(G_{31}) - (G_{31})'\| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35}) - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(KKKKKKKK) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(LLLLLLLLL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(MMMMMMMM) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(NNNNNNNN)$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(OOOOOOOO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(PPPPPPPP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166
A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(gg) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) &d((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}\} (T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (\bar{C})^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (\bar{C})^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)}$ and $(b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(gg) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b_{20}')^{(3)}(b_{21}')^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b_{20}')^{(3)}(b_{21}'')^{(3)}(G_{23}) + (b_{21}')^{(3)}(b_{20}'')^{(3)}(G_{23})] + (b_{20}'')^{(3)}(G_{23})(b_{21}'')^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b_{24}')^{(4)}(b_{25}')^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{21})^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

586
A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} & ((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ & \left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ & ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ & + ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ & ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ & ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ & ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ & + ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ & ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ & + \\ & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ & ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY FIVE

Non-Locality, Near And Far

INTRODUCTION—VARIABLES USED

National Institute for Discovery Science: Nature's Mind: the Quantum Hologram by Edgar Mitchell, Ph.D. Nature's Mind: the Quantum Hologram Fax: 561-641-5242, edgarmitchell@msn.com

- (1) The concept of the quantum hologram adequately and completely describes how this phenomenon might take place
- (2) Further, it is accepted that the mind/brain is a massively parallel processor, capable of performing many tasks simultaneously and subconsciously (in the right, intuitive part of the brain).

- (3) Attention (meaning conscious, focused attention) is a unique and singular task that must take place sequentially, mostly in the left cognitive part of the brain.
- (4) The condition of attention deficit disorder (ADD) is precisely the problem of a percipient being unable to maintain a singular focus for sufficient time to complete a desired task or observation.
- (5) Thus, the action of focusing attention by a percipient may be construed as a necessary condition for pcar to be established with the perceived object.
- (6) Marcer has presented the case for the pcar requirement in normal sensory perception (visual and acoustic).
- (7) A frequent modality used by psychic sensitive individuals to gain information is to physically touch an object. Touching an object satisfies the pcar requirement and presumably allows the percipient access to information about the object not available from space/time information
- (8) Police agencies frequently use this modality with psychic sensitives to gain information about a crime scene, much as they utilize a bloodhound to track the scent of an individual, often with considerable success
- (9) If, as in the theory of the quantum hologram, the object has been in the presence of the individual about whom information is desired, the event history of the object and that of the individual intersect.

NOTATION

Module One

The concept of the

quantum hologram adequately and completely describes how this phenomenon might take place

G_{13} : Category one of quantum hologram

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **percipient** and the object to appear to merge so that a deeper level of information about the object is obtained

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Further, it is accepted that the

mind/brain is a massively parallel processor, capable of performing many tasks simultaneously and subconsciously (in the right, intuitive part of the brain)

G_{16} : Category one of mind/brain

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of massively parallel processor, capable of performing many tasks simultaneously and subconsciously (in the right, intuitive part of the brain)

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Attention (meaning conscious, focused attention) is a unique and singular task that must take place sequentially, mostly in the left cognitive part of the brain

G_{20} : Category one of Attention (meaning conscious, focused attention)

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of unique and singular task that must take place **sequentially**, mostly in the left cognitive part of the brain

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

The

condition of attention deficit disorder (ADD) is (=) precisely the problem of a percipient being unable to maintain a singular focus for sufficient time to complete a desired task or observation

G_{24} : Category one of condition of attention deficit disorder (ADD)

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of problem of a percipient being unable to maintain a singular focus for sufficient time to complete a desired task or observation

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Thus, the

action of focusing attention by a percipient may be construed as a necessary condition for pcar to be established with the perceived object

G_{28} : Category one of action of focusing attention by a percipient is a necessary condition

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of pcar to be established with the perceived object

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

A frequent modality used by psychic sensitive individuals to gain information is to physically touch an object.

Touching an object satisfies the pcar requirement and presumably allows the percipient access to information about the object not available from space/time information

G_{32} : Category one of Touching an object

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of pcar requirement

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

G_{36} : Category one of pcar requirement

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of percipient access to information about the object not available from space/time information

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

If, as in the theory of the quantum hologram, the object has been in the presence of the individual about whom information is desired, the event history of the object and that of the individual intersect.

G_{40} : Category one of object has been in the presence of the individual about whom information is desired

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of event history of the object and that of the individual intersect

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

event history of the object and that of the individual intersect

G_{44} : Category one of event history of the object; event history of individual

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of event history of individual; event history of the object

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b'_{44})^{(9)}((G_{47}), t) =$ **First detriction factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a'_{16})^{(2,2)}(T_{17}, t) + (a'_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a'_{17})^{(2,2)}(T_{17}, t) + (a'_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a'_{18})^{(2,2)}(T_{17}, t) + (a'_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - \left[\begin{array}{c} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b'_{16})^{(2,2)}(G_{19}, t) - (b'_{20})^{(3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4)}(G_{27}, t) - (b'_{28})^{(5,5,5,5)}(G_{31}, t) - (b'_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7)}(G_{39}, t) - (b'_{40})^{(8,8)}(G_{43}, t) - (b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} \boxed{+(a''_{16})^{(2)}(T_{17}, t)} \boxed{+(a''_{13})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{20})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} \boxed{+(a''_{17})^{(2)}(T_{17}, t)} \boxed{+(a''_{14})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{21})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} \boxed{+(a''_{18})^{(2)}(T_{17}, t)} \boxed{+(a''_{15})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{22})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19}, t) & -(b'_{13})^{(1,1)}(G, t) & -(b''_{20})^{(3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b''_{36})^{(7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19}, t) & -(b'_{14})^{(1,1)}(G, t) & -(b''_{21})^{(3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b''_{37})^{(7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19}, t) & -(b'_{15})^{(1,1)}(G, t) & -(b''_{22})^{(3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t) & \\ -(b''_{38})^{(7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19}, t)$, $-(b''_{17})^{(2)}(G_{19}, t)$, $-(b''_{18})^{(2)}(G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$, $-(b''_{14})^{(1,1)}(G, t)$, $-(b''_{15})^{(1,1)}(G, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & + (a''_{20})^{(3)}(T_{21}, t) & + (a'_{16})^{(2,2,2)}(T_{17}, t) & + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & + (a''_{21})^{(3)}(T_{21}, t) & + (a'_{17})^{(2,2,2)}(T_{17}, t) & + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & - (b''_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b''_{17})^{(2,2,2)}(G_{19}, t) & -(b''_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b''_{18})^{(2,2,2)}(G_{19}, t) & -(b''_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$[-(b''_{20})^{(3)}(G_{23}, t)], [-(b''_{21})^{(3)}(G_{23}, t)], [-(b''_{22})^{(3)}(G_{23}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{16})^{(2,2,2)}(G_{19}, t)], [-(b''_{17})^{(2,2,2)}(G_{19}, t)], [-(b''_{18})^{(2,2,2)}(G_{19}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{13})^{(1,1,1)}(G, t)], [-(b''_{14})^{(1,1,1)}(G, t)], [-(b''_{15})^{(1,1,1)}(G, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)], [-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)]$ are fourth detrition coefficients for category 1, 2 and 3

$[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)], [-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)]$ are fifth detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)], [-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)]$ are sixth detrition coefficients for category 1, 2 and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)], [-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)], [-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2 and 3

$[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)], [-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)], [-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eight detrition coefficients for category 1, 2 and 3

$[-(b''_{46})^{(9,9,9)}(G_{47}, t)], [-(b''_{45})^{(9,9,9)}(G_{47}, t)], [-(b''_{44})^{(9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)}[+(a''_{24})^{(4)}(T_{25}, t)] & +(a''_{28})^{(5,5,)}(T_{29}, t) & +(a''_{32})^{(6,6,)}(T_{33}, t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)}[+(a''_{25})^{(4)}(T_{25}, t)] & +(a''_{29})^{(5,5,)}(T_{29}, t) & +(a''_{33})^{(6,6,)}(T_{33}, t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)}[+(a''_{26})^{(4)}(T_{25}, t)] & +(a''_{30})^{(5,5,)}(T_{29}, t) & +(a''_{34})^{(6,6,)}(T_{33}, t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients
category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation
coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation
coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$
are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$
are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$
are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$
are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$
are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for
category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27}, t) & -(b''_{28})^{(5,5)}(G_{31}, t) & -(b''_{32})^{(6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{cccc} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27}, t) & -(b''_{29})^{(5,5)}(G_{31}, t) & -(b''_{33})^{(6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{cccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$
are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$
are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$$

are seventh detrition coefficients for category 1, 2 and 3

$$[-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)]$$

are eighth detrition coefficients for category 1, 2 and 3

$$[-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)] \text{ are ninth detrition coefficients for}$$

category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a''_{25})^{(4,4)}(T_{25}, t) & + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}[-(b''_{29})^{(5)}(G_{31}, t)] & -(b''_{25})^{(4,4)}(G_{27}, t) & -(b''_{33})^{(6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1)}(G, t) & -(b'_{17})^{(2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}[-(b''_{30})^{(5)}(G_{31}, t)] & -(b'_{26})^{(4,4)}(G_{27}, t) & -(b'_{34})^{(6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $[-(b''_{28})^{(5)}(G_{31}, t)]$, $[-(b''_{29})^{(5)}(G_{31}, t)]$, $[-(b''_{30})^{(5)}(G_{31}, t)]$ are first detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4)}(G_{27}, t)]$, $[-(b''_{25})^{(4,4)}(G_{27}, t)]$, $[-(b''_{26})^{(4,4)}(G_{27}, t)]$ are second detrition coefficients for category 1, 2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{33})^{(6,6,6)}(G_{35}, t)]$, $[-(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients for category 1, 2 and 3

$[-(b'_{13})^{(1,1,1,1,1)}(G, t)]$, $[-(b'_{14})^{(1,1,1,1,1)}(G, t)]$, $[-(b'_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for category 1, 2, and 3

$[-(b'_{16})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b'_{17})^{(2,2,2,2,2)}(G_{19}, t)]$, $[-(b'_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients for category 1, 2, and 3

$[-(b'_{20})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b'_{21})^{(3,3,3,3,3)}(G_{23}, t)]$, $[-(b'_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients for category 1, 2, and 3

$[-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$, $[-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition coefficients for category 1, 2, and 3

$[-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$, $[-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition coefficients for category 1, 2, and 3

$[-(b'_{46})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b'_{45})^{(9,9,9,9,9)}(G_{47}, t)]$, $[-(b'_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)}[+(a''_{32})^{(6)}(T_{33}, t)] & +(a''_{28})^{(5,5,5)}(T_{29}, t) & +(a''_{24})^{(4,4,4)}(T_{25}, t) \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)}[+(a''_{33})^{(6)}(T_{33}, t)] & +(a''_{29})^{(5,5,5)}(T_{29}, t) & +(a''_{25})^{(4,4,4)}(T_{25}, t) \\ +(a'_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}[+(a''_{34})^{(6)}(T_{33}, t)] & +(a''_{30})^{(5,5,5)}(T_{29}, t) & +(a''_{26})^{(4,4,4)}(T_{25}, t) \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a'_{32})^{(6)}(T_{33}, t)$, $+(a'_{33})^{(6)}(T_{33}, t)$, $+(a'_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)}$

Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)}$ ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} \boxed{(b'_{32})^{(6)}(G_{35}, t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} \boxed{(b'_{33})^{(6)}(G_{35}, t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} \boxed{(b'_{34})^{(6)}(G_{35}, t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{34} & 90 \end{aligned}$$

$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6)}(G_{35}, t)}$ are first detrution coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)}$ are second detrution coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)}$ are third detrution coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)}$ are fourth detrution coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)}$ are fifth detrution coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)}$ are sixth detrution

coefficients for category 1, 2, and 3

$$[-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39}, t)] \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$[-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)]$$

are eighth detrition coefficients for category 1, 2, and 3

$$[-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)] \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a'_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation

coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) & + (a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)} + (a_{41}'')^{(8)}(T_{41}, t) & + (a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)} + (a_{42}'')^{(8)}(T_{41}, t) & + (a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41}, t)$, $+(a_{41}'')^{(8)}(T_{41}, t)$, $+(a_{42}'')^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

96

$$= (a_{44})^{(9)}G_{45}$$

$$- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)} - (b'_{44})^{(9)}(G_{47}, t) & - (b'_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t) \quad - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t) \quad - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) \quad - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47}, t)$, $-(b''_{45})^{(9)}(G_{47}, t)$, $-(b''_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b'_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a'_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b'_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition:

99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$:

106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition:

114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21}' - T_{21}| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} \|G_{23} - G_{23}'\| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$:

115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26$$

117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

118

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition:

119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$:

120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30$$

122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

123

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

128

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33}' - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(QQQQQQQQ) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38$$

131

(RRRRRRRR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(SSSSSSSS) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(TTTTTTTTT)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ are positive constants and $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37}' - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(UUUUUUUU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(VVVVVVVV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40, 41, 42$$

136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||G_{43} - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T'_{45}, t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47}) - (G_{47})'| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T'_{45}, t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153
B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(hh) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{((\hat{P}_{13})^{(1)} + G_j^0)}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{((\hat{Q}_{13})^{(1)} + T_j^0)}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{(\bar{M}_{16})^{(2)} s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{-(\bar{M}_{16})^{(2)} s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$| (G_{19})^{(1)} - (G_{19})^{(2)} | e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}\} (T_{25}(s_{(24)}), s_{(24)}) ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)}) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 409

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-

$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(8)}$ and $(b'_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(hh) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b'_{14})^{(1)}(G) + (b'_{14})^{(1)}(b'_{13})^{(1)}(G)] + (b'_{13})^{(1)}(G)(b'_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b'_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b'_{16})^{(2)}(G_{19})] + (b'_{16})^{(2)}(G_{19})(b'_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b'_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b'_{20})^{(3)}(G_{23})] + (b'_{20})^{(3)}(G_{23})(b'_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + \left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \{ (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \} \\ & \left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + \left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) \left((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)} \right) \{ (\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)} \} \\ & \left[\left(((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 & \left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY SIX

Attention And Intention

INTRODUCTION—VARIABLES USED

file:///C:/Documents%20and%20Settings/Judy/My%20...m%20Hologram%20by%20Edgar%20Mitchell,%20Ph_D.htm (8 of 18) [6/25/2011 9:21:06 PM] National Institute for Discovery Science: Nature's Mind: the Quantum Hologram by Edgar Mitchell, Ph.D.

- (1) The **Berry phase information of the object** contains its journey in three dimensional space and time, as well as the quantum states through which it has passed on this journey. The sensitive individual, with a honed talent, seems often able to decode useful Berry phase information from the object

about the individual sought.

- (2) It may also be the case with the blood hound, which additional non-local information has been gained about the subject, even though the classical explanation is that the animal is operating only with heightened olfactory sensing.
- (3) Although perception in the three dimensional world requires and utilizes pcar, most humans, however, do not bring to conscious awareness non-local information when we are routinely operating in three-dimensional reality. We perceive objects as presented by space/time information, that is, shape, color, function (tree, chair, table, etc) but are not usually aware of the additional non-local information
- (4) A powerful and telling series of experiments conducted by Dean Radin (1997) at University of Nevada at Las Vegas following a decade long set of equally significant experiments by Brenda Dunne and Robert Jahn at Princeton University (1988) provide insight as to the subtleties involved in this level of mind/brain functioning. Jahn and Dunne provided overwhelming evidence that subjects could intentionally produce statistically skewed results in mechanical processes normally thought to be driven by random processes.
- (5) Radin went further; he discovered that audiences watching a stage performance would skew the output of nearby random number generators during periods of high emotional content in the stage performance.
- (6) Further, in a wide-ranging audience participation experiment, he recorded the output of computer random number generators during the television broadcasts of the O.J. Simpson murder trial. Most television media reported this event for weeks on end and tens of millions of humans were watching the results. Results of the random number generators were skewed (e&eb) corresponding to emotional peaks during the trial drama and corresponding to the number of people watching television
- (7) **The thesis in the Princeton experiments was that participant intentionality created a non-random effect to bias the skewed distribution.**
- (8) In the Radin experiments the results were not intentional, as the participants were unaware of the experiment, but the hypothesis was that attention (in particular, rapt attention) drove (eb) the system away from chaos (randomness) and toward (e&eb) greater order (reduced entropy).
- (9) These results suggest that attention and intention provide closely correlated outcomes, further, that randomness may not be a general property of nature, but that what is perceived as random noise in a system may be information (a pattern of energy) that is not in resonance at that moment with the particular perceptual system.
- (10) William Tiller, emeritus professor at Stanford also has performed experiments (1997) that are consistent with these results, though his interpretation of the operating mechanism is somewhat different.

NOTATION

Module One

The

Berry phase information of the object contains its journey in three dimensional space and time, as well as the quantum states through which it has passed on this journey

G_{13} : Category one of journey in three dimensional space and time, as well as the quantum states through which it has passed on this journey **(note this is what we have called individual karma: the individual general ledger)**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **Berry phase information of the object**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

It may also be the case with the

blood hound garners additional non-local information about the subject, even though the classical explanation is that the animal is (=) operating only with heightened olfactory sensing

I have slightly changed the sentence

G_{16} : Category one of blood hound garners additional non-local information about the subject, even though the classical explanation is that the animal

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of operating only with heightened olfactory sensing

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Although

perception in the three dimensional world requires and utilizes pcar,

most humans, however, do not (e) bring to conscious awareness non-local information when we are routinely operating in three-dimensional reality.

We perceive objects as presented by space/time information, that is, shape, color, function (tree, chair, table, etc) but are not usually aware of the additional non-local information

G_{20} : Category one of pcar

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of perception in the three dimensional world

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

most humans, however, do not (e) bring to conscious awareness non-local information when we are routinely operating in three-dimensional reality

G_{24} : Category one of conscious awareness non-local information when we are routinely operating in three-dimensional reality

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of most humans

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

A powerful and telling series of experiments conducted by Dean Radin (1997) at University of Nevada at Las Vegas following a decade long set of equally significant experiments by Brenda Dunne and Robert Jahn at Princeton University (1988) provide insight as to the subtleties involved in this level of mind/brain functioning. Jahn and Dunne provided overwhelming evidence that

subjects could intentionally produce statistically skewed results in mechanical processes normally thought to be driven by random processes.

G_{28} : Category one of subjects could intentionally

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of statistically skewed results in mechanical processes normally thought to be driven by random processes

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Radin went further; he discovered that audiences watching a stage performance would skew (e&eb) the output of nearby random number generators during periods of high emotional content in the stage performance
 G_{32} : Category one of audiences watching a stage performance; output of nearby random number generators during periods of high emotional content in the stage performance

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of output of nearby random number generators during periods of high emotional content in the stage performance; audiences watching a stage performance

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Further, in a wide-ranging audience participation experiment, he recorded the output of computer random number generators during the television broadcasts of the O.J. Simpson murder trial. Most television media reported this event for weeks on end and tens of millions of humans were watching the results.

Results of the random number generators were skewed (e&eb) corresponding to emotional peaks during the trial drama and corresponding to the number of people watching television

G_{36} : Category one of Results of the random number generators; corresponding to emotional peaks during the trial drama and corresponding to the number of people watching television

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of corresponding to emotional peaks during the trial drama and corresponding to the number of people watching television; Results of the random number generators

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

The thesis in the Princeton experiments was that participant intentionality created a non-random effect to bias the skewed distribution

G_{40} : Category one of participant intentionality

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of non-random effect to bias the skewed distribution

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

In the Radin experiments the results were not intentional, as the participants were unaware of the experiment, but the hypothesis was that

attention (in particular, rapt attention) drove (eb) the system away from chaos (randomness) and toward (e&eb) greater order (reduced entropy)

G_{44} : Category one of attention (in particular, rapt attention)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of system away from chaos (randomness) and toward (e&eb) greater order (reduced entropy)

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY SEVEN

Suppression By Cultural Conditioning In Childhood And Subsequent Lack Of Practice Cause The Natural Ability For Conscious, Intuitive Perceptions To Atrophy

INTRODUCTION—VARIABLES USED

National Institute for Discovery Science: Nature's Mind: the Quantum Hologram by Edgar Mitchell, Ph.D. Nature's Mind: the Quantum Hologram Fax: 561-641-5242, edgarmitchell@msn.com

- (6) It takes training as provided by many of the esoteric traditions and/or certain naturally sensitive individuals to routinely perceive the non-local holographic information associated with a particular object.
- (7) There is massive evidence to suggest, however, that the brain has these latter capabilities at birth. **Suppression by cultural conditioning in childhood and subsequent lack of practice cause (eb) the natural ability for conscious, intuitive perceptions to atrophy.**
- (8) Particularly in western tradition, educational interest has been on the left brain, rational functions rather than right brain, intuitive functions. However, mystic adepts and natural psychics routinely demonstrate that non-local information is perceptible from physical objects by focusing attention, quieting the left brain and allowing intuitive perceptions to appear.
- (9) It is the left brain cognitive ability in humans that provides canonical labeling of the intuitive and artistic processes taking place in the right brain.
- (10) The fact that with training and practice, individuals can recover, deepen and label their individual cognitive access to intuitive, non-local information demonstrates that learning is taking place within the whole brain itself and involves enhanced coherence and coordination between the hemispheres. This process is different and distinct from the left brain function of extending and extrapolating factual data and logical deduction to leap to (e&eb) an "intuitive" conclusion, while omitting the intermediate steps leading to that conclusion.

NOTATION

Module One

attention (in particular, rapt attention) drove the system away from chaos (randomness) and toward (e&eb) greater order (reduced entropy)

G_{13} : Category one of attention (in particular, rapt attention) drove the system away from chaos (randomness); greater order (reduced entropy)

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of greater order (reduced entropy); attention (in particular, rapt attention) drove the system away from chaos (randomness)

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

These results suggest that

attention and intention provide (eb) closely correlated outcomes

further, that

randomness may not be a general property of nature, but that what is perceived as random noise in a system may be information (a pattern of energy) that is (=) not in resonance at that moment with the particular perceptual system

G_{16} : Category one of attention and intention

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of closely correlated outcomes

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

randomness may not be a general property of nature, but that what is perceived as random noise in a system may be information (a pattern of energy) that is (=) dissonance at that moment with the particular perceptual system

G_{20} : Category one of randomness may not be a general property of nature, but that what is perceived as random noise in a system may be information (a pattern of energy)

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of dissonance at that moment with the particular perceptual system

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

It takes training as provided by many of the esoteric traditions and/or certain naturally sensitive individuals to routinely perceive (e) the non-local holographic information associated with a particular object

G_{24} : Category one of non-local holographic information associated with a particular object

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of training as provided by many of the esoteric traditions and/or certain naturally sensitive individuals

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

There is massive evidence to suggest, however, that the brain has these latter capabilities at birth.

Suppression by cultural conditioning in childhood and subsequent lack of practice cause (eb) the natural ability for conscious, intuitive perceptions to atrophy

G_{28} : Category one of **natural ability for conscious, intuitive perceptions to atrophy**

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of **Suppression by cultural conditioning in childhood and subsequent lack of practice**

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Particularly in western tradition, educational interest has been on the left brain, rational functions rather than right brain, intuitive functions. However, mystic adepts and natural psychics routinely demonstrate that

non-local information is perceptible from physical objects by (e) focusing attention, quieting the left brain and allowing intuitive perceptions to appear

G_{32} : Category one of focusing attention, quieting the left brain and allowing intuitive perceptions to appear

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of non-local information is perceptible from physical objects

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

quieting the left brain and allowing intuitive perceptions to appear

G_{36} : Category one of quieting the left brain

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of intuitive perceptions to appear

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

It is the

left brain cognitive ability in humans that provides canonical labeling of the intuitive and artistic processes taking place in the right brain

G_{40} : Category one of left brain cognitive ability in humans

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of canonical labeling of the intuitive and artistic processes taking place in the right brain

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

The fact that with training and practice, individuals can recover, deepen and label their individual cognitive access to intuitive, non-local information demonstrates that learning is taking place within the whole brain itself and involves enhanced coherence and coordination between the hemispheres.

enhanced coherence and coordination between the hemispheres process is different and distinct from the left brain function of extending and extrapolating factual data and logical deduction to leap to (and) an "intuitive" conclusion, while omitting the intermediate steps leading to that conclusion.

G_{44} : Category one of enhanced coherence and coordination between the hemispheres process is different and distinct from the left brain function of extending and extrapolating factual data and logical deduction; "intuitive" conclusion, while omitting the intermediate steps leading to that conclusion

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of "intuitive" conclusion, while omitting the intermediate steps leading to that conclusion; enhanced coherence and coordination between the hemispheres process is different and distinct from the left brain function of extending and extrapolating factual data and logical deduction

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &- (b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \end{aligned}$$

$$\begin{aligned}\frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor}\end{aligned}$$

12

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20}$$

13

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21}$$

14

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22}$$

15

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20}$$

16

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21}$$

17

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22}$$

18

$$+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$- (b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24}$$

19

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25}$$

20

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26}$$

21

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24}$$

22

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25}$$

23

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26}$$

24

$$+ (a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$- (b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28}$$

25

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29}$$

26

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30}$$

27

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28}$$

28

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29}$$

29

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30}$$

30

$$+ (a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$- (b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\begin{aligned}\frac{dG_{32}}{dt} &= (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} & 31 \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} & 32 \\ \frac{dG_{34}}{dt} &= (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} & 33 \\ \frac{dT_{32}}{dt} &= (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} & 34 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} & 35 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} & 36 \\ &+ (a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor} \\ &\text{Module Numbered Seven:}\end{aligned}$$

The differential system of this model is now (Seventh Module)

$$\begin{aligned}\frac{dG_{36}}{dt} &= (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)] T_{38} & 42 \\ &+ (a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \\ &\text{Module Numbered Eight}\end{aligned}$$

The differential system of this model is now

$$\begin{aligned}\frac{dG_{40}}{dt} &= (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} & 44 \\ \frac{dG_{42}}{dt} &= (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)] T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)] T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)] T_{42} & 48\end{aligned}$$

Module Numbered Nine

The differential system of this model is now

$$\begin{aligned}\frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)] T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)] T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)] T_{46} & 54 \\ &+ (a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor} \\ &- (b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}\end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[\begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[\begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where $(a'_{13})^{(1)}(T_{14}, t)$, $(a'_{14})^{(1)}(T_{14}, t)$, $(a'_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{16})^{(2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{20})^{(3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{24})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{25})^{(4,4,4,4)}(T_{25}, t)$, $(a'_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$(a'_{28})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5,5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6,6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$(a'_{38})^{(7,7)}(T_{37}, t)$, $(a'_{37})^{(7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$(a'_{40})^{(8,8)}(T_{41}, t)$, $(a'_{41})^{(8,8)}(T_{41}, t)$, $(a'_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$(a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $(a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) - (b''_{18})^{(2,2)}(G_{19}, t) - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) - (b''_{42})^{(8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{16})^{(2)} + (a'_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{36})^{(7,7,7)}(T_{37}, t) + (a'_{40})^{(8,8,8)}(T_{41}, t) + (a'_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{l} (a'_{17})^{(2)} + (a'_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{37})^{(7,7,7)}(T_{37}, t) + (a'_{41})^{(8,8,8)}(T_{41}, t) + (a'_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{l} (a'_{18})^{(2)} + (a'_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a'_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a'_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a'_{38})^{(7,7,7)}(T_{37}, t) + (a'_{42})^{(8,8,8)}(T_{41}, t) + (a'_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} \boxed{(b'_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} \boxed{(b'_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} & 67 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} & 68 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} & 69 \end{aligned}$$

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} & 70 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} & 71 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} & 72 \end{aligned}$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{c} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{c} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$

are sixth augmentation coefficients for category 1, 2 and 3

$$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$$

are seventh augmentation coefficients for category 1, 2 and 3

$$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$$

are eighth augmentation coefficients for category 1, 2 and 3

$$+(a''_{46})^{(9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9)}(T_{45}, t) \text{ are ninth detrition coefficients for category 1 2 3}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27}, t), -(b''_{25})^{(4)}(G_{27}, t), -(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t), -(b''_{29})^{(5,5)}(G_{31}, t), -(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t), -(b''_{33})^{(6,6)}(G_{35}, t), -(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1)}(G, t)$

are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3)}(G_{23}, t)$

are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$

are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29}, t)$, $+(a''_{29})^{(5)}(T_{29}, t)$, $+(a''_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{c} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[\begin{array}{c} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) - (b''_{26})^{(4,4)}(G_{27}, t) - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients

for category 1,2 and 3

$[-(b''_{32})^{(6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6)}(G_{35}, t)]$ are third detrition coefficients

for category 1,2 and 3

$[-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)]$ are fourth detrition coefficients for

category 1,2, and 3

$[-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)]$ are fifth detrition coefficients

for category 1,2, and 3

$[-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)]$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)]$ are seventh detrition

coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)]$ are eighth detrition

coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)]$ are ninth detrition coefficients

for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \left[\begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients

for category 1,2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation

coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation

coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation

coefficients

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$ sixth augmentation

coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\begin{aligned} \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} & 88 \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} & 89 \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} & 90 \end{aligned}$$

$$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t) \text{ are first detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t) \text{ are second detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t) \text{ are third detrition coefficients}$$

for category 1, 2 and 3

$$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t) \text{ are fourth detrition coefficients}$$

for category 1, 2, and 3

$$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) \text{ are fifth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \text{ are sixth detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \text{ are seventh detrition}$$

coefficients for category 1, 2, and 3

$$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1, 2, and 3

$$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \text{ are ninth detrition}$$

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)}$, $\boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)}$

are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$, $\boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \begin{bmatrix} \boxed{+(a'_{40})^{(8)}(T_{41}, t)} & \boxed{+(a'_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \begin{bmatrix} \boxed{+(a'_{41})^{(8)}(T_{41}, t)} & \boxed{+(a'_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a'_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a'_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} & \boxed{+(a'_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41}, t)$, $+(a''_{41})^{(8)}(T_{41}, t)$, $+(a''_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} \\ &- \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} \\ &- \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)} \text{ are third}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)} \text{ are fourth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)} \text{ are fifth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)} \text{ are sixth}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)} \text{ are Seventh}$$

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)} \text{ are eighth}$$

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)} \text{ are ninth}$$

augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{ccc} \boxed{(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} \boxed{(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \left[\begin{array}{ccc} \boxed{(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{44})^{(9)}(G_{47}, t)}, \boxed{-(b'_{45})^{(9)}(G_{47}, t)}, \boxed{-(b'_{46})^{(9)}(G_{47}, t)}$ are first detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}, \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} \text{ are second}$$

detrition coefficients for category 1, 2 and 3

$$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}, \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \text{ are third detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} \text{ are fourth}$$

detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)} t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \quad 109$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30$$

122

The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

123

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

127

The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

128

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(6)}(T'_{33}, t)$ and $(a''_i)^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a''_i)^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a''_i)^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(WWWWWWW) \quad (a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(XXXXXXXX) The functions $(a''_i)^{(7)}, (b''_i)^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(YYYYYYYY) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(ZZZZZZZZ)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(7)}(T'_{37}, t)$ and $(a''_i)^{(7)}(T_{37}, t)$. (T'_{37}, t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a''_i)^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a''_i)^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(AAAAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(BBBBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42$$

136

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$:

137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$$

138

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)}$$

139

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)}$$

140

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)}$$

141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)} t}$$

142

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$$

143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a''_i)^{(8)}(T_{41}, t)$. T'_{41}, t and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$, Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44, 45, 46 \quad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$

and $(a_i'')^{(9)}(T_{45}, t) \cdot (T_{45}', t)$ and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

166

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)}t} - 1 \right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ii) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} = \\ (1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \\ (1 + (a_{40})^{(8)} t) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right) \quad 180$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} = \\ (1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right) \quad 181$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\bar{P}_{13})^{(1)} + ((\bar{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\bar{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[((\bar{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{13})^{(1)} \right] \leq (\bar{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\bar{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)} + (\bar{P}_{13})^{(1)} (\bar{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\bar{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on

G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, $((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\widehat{P}_{16})^{(2)}$ and $(\widehat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\bar{P}_{16})^{(2)} + ((\bar{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\bar{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\bar{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{16})^{(2)} \right] \leq (\bar{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left((G_{19})^{(1)}, (T_{19})^{(1)}, (G_{19})^{(2)}, (T_{19})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16, 17, 18$ depend only on T_{17} and respectively on

(G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 202
analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad \text{205}$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\bar{P}_{20})^{(3)} + ((\bar{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\bar{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{20})^{(3)} \right] \leq (\bar{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23}$: $(\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)}) d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$ it follows $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is 218
analogous with the preceding one. An analogous property is true if G_{21} is bounded from below.

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), ((G_{27})^{(2)}, (T_{27})^{(2)}) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}, \widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \quad 226$$

$$\frac{1}{(\bar{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} + (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)}) d\left((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}\right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on

(G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$ it follows $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is 230
analogous with the preceding one. An analogous property is true if G_{25} is bounded from below.

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left((G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}, \widetilde{T_{31}})$: $(\widetilde{G_{31}}, \widetilde{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{-(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\hat{M}_{28})^{(5)} s_{(28)}} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\hat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\hat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) &d\left((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as 238
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ and $(\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is 241
analogous with the preceding one. An analogous property is true if G_{29} is bounded from below.

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)}((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , G_{34} , T_{32} , T_{33} , T_{34}

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}, \widetilde{T_{35}})$: $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{-(\hat{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\hat{M}_{32})^{(6)}s_{(32)}} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)}t} &\leq \\ \frac{1}{(\hat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$ and $(\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)})_1 \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 252
analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , G_{38} , T_{36} , T_{37} , T_{38} 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{39}}, \widehat{T_{39}}) : (\widehat{G_{39}}, \widehat{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{-(\hat{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a'_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\hat{M}_{36})^{(7)}s_{(36)}} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\hat{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\hat{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\tilde{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as 260
not conformal with the reality, however we have put this hypothesis, in order that we can postulate
condition necessary to prove the uniqueness of the solution bounded by
 $(\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{36})^{(7)})_1$, $((\widehat{M}_{36})^{(7)})_2$ and $((\widehat{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\widehat{M}_{36})^{(7)})_1 \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is 263
analogous with the preceding one. An analogous property is true if G_{37} is bounded from below.

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} ((b_{38}'')^{(7)}((G_{39})(t), t)) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} < 1$ and to choose $(\bar{P}_{40})^{(8)}$ and $(\bar{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[(\bar{P}_{40})^{(8)} + ((\bar{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\bar{P}_{40})^{(8)} + G_j^0}{G_j^0}\right)} \right] \leq (\bar{P}_{40})^{(8)} \quad 266$$

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[((\bar{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\bar{Q}_{40})^{(8)} \right] \leq (\bar{Q}_{40})^{(8)} \quad 267$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left((G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \quad 268$$

Indeed if we denote

$$\textbf{Definition of } (\bar{G}_{43}), (\bar{T}_{43}) : (\bar{G}_{43}), (\bar{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43})) \quad 269$$

It results

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t ((a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}) ds_{(40)} \end{aligned} \quad 270$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \quad 271$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by

$(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ and $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1$, $((\widehat{M}_{40})^{(8)})_2$ and $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)} ((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way, one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)} ((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

279
A

$$\frac{(a_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[(\hat{P}_{44})^{(9)} + ((\hat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\bar{M}_{44})^{(9)}} \left[((\hat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{44})^{(9)} \right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : ((\widetilde{G_{47}}), (\widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}) d \left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(9)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{44})^{(9)})_1$, $((\widehat{M}_{44})^{(9)})_2$ and $((\widehat{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < ((\widehat{M}_{44})^{(9)}) \text{ it follows } \frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)} ((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)} ((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b_{46}'')^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$$\begin{aligned} &(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)} \text{ four constants satisfying} \\ &-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)} \\ &-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)} \end{aligned}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

$$\begin{aligned} &\text{By } (v_1)^{(1)} > 0, (v_2)^{(1)} < 0 \text{ and respectively } (u_1)^{(1)} > 0, (u_2)^{(1)} < 0 \text{ the roots of the equations} \\ &(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0 \end{aligned}$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

$$\begin{aligned} &\text{By } (\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0 \text{ and respectively } (\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0 \text{ the roots of the equations} \\ &(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0 \end{aligned}$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

$$\begin{aligned} &\text{If we define } (m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)} \text{ by} \\ &(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)} \\ &(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}, \end{aligned}$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$\begin{aligned} &(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)} \\ &(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}, \end{aligned}$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying 293

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a'_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b'_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 302$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)}) - (S_2)^{(2)}} \right) \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 317$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)} (v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^0e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

337

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation

349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)} (T_{33}, t) + (a''_{33})^{(6)} (T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)} ((G_{35}), t) - (b''_{33})^{(6)} ((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

$$\text{roots of the equations } (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

354

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq$$

355

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

356

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation

371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$:

372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations
 $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
 and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} [e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad (C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case,

385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)} (\nu_2)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((\nu_1)^{(1)} - (\nu_2)^{(1)}) t]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)} (\bar{\nu}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}) t]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof: From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:-

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

388

It follows

389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)} G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t) \right) - (a_{25}'')^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$-\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .**

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:-
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_1)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)}e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)}e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case**.

Analogously if $(b_{28}''^{(5)} = (b_{29}''^{(5)})$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)} (T_{33}, t) \right) - (a_{33}'')^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_0)^{(6)}) t]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} ((v_1)^{(6)} - (v_2)^{(6)}) t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} ((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}) t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-

$$\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$- \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)} (v_2)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)} ((v_1)^{(8)} - (v_2)^{(8)}) t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)} (\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)} ((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}) t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case.**

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then

$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

424

Proof : From 99,20,44,22,23,44 we obtain

A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)} (T_{45}, t) \right) - (a_{45}'')^{(9)} (T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)} (v^{(9)})^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((\nu_1)^{(9)} - (\nu_2)^{(9)})t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)})t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, and definition of $(u_0)^{(9)}$.

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations

431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following

432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a_{44}')^{(9)}(a_{45}')^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a_{45}')^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b_{44}')^{(9)}(b_{45}')^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b_{44}')^{(9)}(r_{45})^{(9)} - (b_{45}')^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a_{13}')^{(1)} + (a_{13}'')^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a_{14}')^{(1)} + (a_{14}'')^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a_{15}')^{(1)} + (a_{15}'')^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b_{13}')^{(1)} - (b_{13}'')^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b_{14}')^{(1)} - (b_{14}'')^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b_{15}')^{(1)} - (b_{15}'')^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b_{16}')^{(2)} - (b_{16}'')^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b_{17}')^{(2)} - (b_{17}'')^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b_{18}')^{(2)} - (b_{18}'')^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a_{20}')^{(3)} + (a_{20}'')^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a_{21}')^{(3)} + (a_{21}'')^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a_{22}')^{(3)} + (a_{22}'')^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b_{20}')^{(3)} - (b_{20}'')^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof:

486

(ii) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof:

487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof:

488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

495

Definition and uniqueness of T_{21}^* :-

496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :-

497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :-

498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a'_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :-

499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :-

500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :-

501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501

A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26})$, G_{24}, G_{26} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30})$, G_{28}, G_{30} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34})$, G_{32}, G_{34} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi((G_{35})^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38})$, G_{36}, G_{38} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b'_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b'_{40})^{(8)}(G_{43})] + (b'_{40})^{(8)}(G_{43})(b'_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b'_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b'_{44})^{(9)}(G_{47})] + (b'_{44})^{(9)}(G_{47})(b'_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_{14}'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS

531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*G_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18}(s_{(17)(j)}T_{17}^*G_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18}(s_{(18)(j)}T_{18}^*G_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22}(s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{25})^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* T_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)} T_{24}^* \mathbb{G}_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)} T_{25}^* \mathbb{G}_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)} T_{26}^* \mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* T_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}(s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}(s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*\mathbb{G}_j) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*\mathbb{G}_j) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*\mathbb{G}_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a_{36}')^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a_{37}')^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a_{38}')^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b_{36}')^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b_{37}')^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b_{38}')^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a_{40}')^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a_{41}')^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a_{42}')^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b_{40}')^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42}(s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)} T_{41}^* \mathbb{G}_j) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)} T_{42}^* \mathbb{G}_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$ belong to $\mathcal{C}^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{45})^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b'_{47})^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{d\mathbb{G}_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^* \mathbb{T}_{45} \quad 586$$

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(44)(j)} T_{44}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46} (s_{(45)(j)} T_{45}^* \mathbb{G}_j) \quad 586$$

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46} (s_{(46)(j)} T_{46}^* \mathbb{G}_j) \quad 586$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right]$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*)$$

$$+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^*)$$

$$((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*)$$

$$((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)}$$

$$((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^*) \\
 & \left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}) \{ ((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 & \left(((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & \left(((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
 & \left(((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 &+ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 &\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 &\left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 &+ ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 &\left(((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &\left(((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 &+ ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 &+ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 &\left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)}) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 &\left[((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 &+ ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 &\left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 &\left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 &\left(((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 &+ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^*) \\
 &((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY EIGHT

Easy Quantum Physics

INTRODUCTION—VARIABLES USED

Quantum physics just got less complicated Fri, 12/19/2014 - 2:21pm Get today's R&D headlines and news - Sign up now!

- (1) Quantum physics says that particles can behave like waves, and vice versa. Researchers have now shown that this 'wave-particle duality' is (=) simply the quantum uncertainty principle in disguise. Source: Timothy Yeo / CQT, National University of Singapore
- (2) Here's a nice surprise: quantum physics is less complicated than we thought. An international team of researchers has proved that two peculiar features of the quantum world previously considered distinct are different manifestations of the same thing. The result is published in Nature Communications. Patrick Coles, Jędrzej Kaniewski, and Stephanie Wehner made the breakthrough while at the Centre for Quantum Technologies at the National University of Singapore. They found that 'wave-particle duality' is simply the quantum 'uncertainty principle' in disguise, reducing two mysteries to one.
- (3) "The connection between uncertainty and (e&eb) wave-particle duality comes out very naturally when you consider them as (=) questions about what information you can gain about a system. Our result highlights the power of thinking about physics from the perspective of information," says Wehner, who is now an Associate Professor at QuTech at the Delft University of Technology in the Netherlands. The discovery deepens our understanding of quantum physics and could prompt ideas for new applications of wave-particle duality.
- (4) Wave-particle duality is the idea that a quantum object can behave like a wave, but that the wave behaviour disappears (e) if you try to locate the object.
- (5) It's most simply seen in a double slit experiment, where single particles, electrons, say, are fired one by one at a screen containing two narrow slits. The particles pile up behind the slits not in two heaps as classical objects would, but in (eb) a stripy pattern like you'd expect for waves interfering. At least this is what happens until you sneak a look at which slit a particle goes through - do that and the (eb) interference pattern vanishes.
- (6) The quantum uncertainty principle is the idea that it's (=) impossible to know certain pairs of things about a quantum particle at once.
- (7) For example, the more precisely you know the position of an atom, (eb) the less precisely you can know the speed with which it's moving.
- (8) It's (=) a limit on the fundamental knowability of nature, not a statement on measurement skill.
- (9) The new work shows that how much you can learn about the wave versus the particle behaviour of a system is (=) constrained in exactly the same way. Wave-particle duality and uncertainty have

been fundamental concepts in quantum physics since the early 1900s. "We were guided by a gut feeling, and only a gut feeling, that there should be a connection," says Coles, who is now a Postdoctoral Fellow at the Institute for Quantum Computing in Waterloo, Canada.

- (10) It's possible to write equations that capture how much can be learned about pairs of properties that are affected by the uncertainty principle. Coles, Kaniewski and Wehner are experts in a form of such equations known as 'entropic uncertainty relations', and they discovered that all the maths previously used to describe wave-particle duality could be reformulated in terms of these relations.
- (11) "It was like we had discovered the 'Rosetta Stone' that connected two different languages," says Coles. "The literature on wave-particle duality was like hieroglyphics that we could now translate into our native tongue. We had several eureka moments when we finally understood what people had done," he says.
- (12) Because the entropic uncertainty relations used in their translation have also been used (e) in proving the security of quantum cryptography - schemes for secure communication using quantum particles - the researchers suggest the work could help inspire new cryptography protocols.
- (13) In earlier papers, Wehner and collaborators found connections between the uncertainty principle and other physics, namely (e&eb) quantum 'non-locality' and the second law of thermodynamics. The tantalizing next goal for the researchers is to think about how these pieces fit together and what bigger picture that paints of how nature is constructed. Source: Centre for Quantum Technologies

NOTATION

Module One

Quantum physics says that particles can behave like waves, and vice versa. Researchers have now shown that this

'wave-particle duality' is (=) simply the quantum uncertainty principle in disguise.

Source: Timothy Yeo / CQT, National University of Singapore

G_{13} : Category one of 'wave-particle duality'

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of quantum uncertainty principle in disguise

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

"The connection between uncertainty and (e&eb) wave-particle duality comes out very naturally when you consider them as (=) questions about what information you can gain about a system.

Our result highlights the power of thinking about physics from the perspective of information," says Wehner, who is now an Associate Professor at QuTech at the Delft University of Technology in the Netherlands. The discovery deepens our understanding of quantum physics and could prompt ideas for new applications of wave-particle duality.

G_{16} : Category one of uncertainty; wave-particle duality

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of wave-particle duality ;uncertainty

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Wave-particle duality is the idea that a quantum object can behave like a wave, but that the wave behaviour disappears (e) if you try to locate the object.

G_{20} : Category one of try to locate the object

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of disappearance of the wave behaviour

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

It's most simply seen in a double slit experiment, where single particles, electrons, say, are fired one by one at a screen containing two narrow slits. The particles pile up behind the slits not in two heaps as classical objects would, but in (eb) a stripy pattern like you'd expect for waves interfering.

At least this is what happens until you sneak a look at which slit a particle goes through - do that and the (eb) interference pattern vanishes

G_{24} : Category one of sneak a look at which slit a particle goes through

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of interference pattern vanishes

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The quantum uncertainty principle is the idea that it's (=) impossible to know certain pairs of things about a quantum particle at once.

G_{28} : Category one of possibility to know certain pairs of things

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of quantum uncertainty principle

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

For example, the

more precisely you know the position of an atom, (eb) the less precisely you can know the speed with which it's moving

G_{32} : Category one of more precisely you know the position of an atom

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of less precisely you can know the speed with which it's moving

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

The new work shows that

how much you can learn about the wave versus the particle behaviour of a system is (=) constrained in exactly the same way.

Wave-particle duality and uncertainty have been fundamental concepts in quantum physics since the early 1900s. "We were guided by a gut feeling, and only a gut feeling, that there should be a connection," says Coles, who is now a Postdoctoral Fellow at the Institute for Quantum Computing in Waterloo, Canada

G_{36} : Category one of how much you can learn about the wave versus the particle behaviour of a system

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of constrained in exactly the same way.

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

In earlier papers, Wehner and collaborators found connections between the uncertainty principle and other physics, namely (e&eb) quantum 'non-locality' and the second law of thermodynamics.

The tantalizing next goal for the researchers is to think about how these pieces fit together and what bigger picture that paints of how nature is constructed. Source: Centre for Quantum Technologies

G_{40} : Category one of uncertainty principle; quantum 'non-locality'

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of quantum 'non-locality'; uncertainty principle

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

G_{44} : Category one of uncertainty principle; second law of thermodynamics

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of second law of thermodynamics; uncertainty principle

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
 are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\begin{aligned}\frac{dG_{13}}{dt} &= (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} & 6 \\ &+ (a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\begin{aligned}\frac{dG_{16}}{dt} &= (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} & 12 \\ &+ (a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\begin{aligned}\frac{dG_{20}}{dt} &= (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} & 18 \\ &+ (a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor}\end{aligned}$$

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

42

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40}$$

43

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41}$$

44

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42}$$

45

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40}$$

46

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41}$$

47

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42}$$

48

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44}$$

49

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45}$$

50

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46}$$

51

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44}$$

52

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45}$$

53

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

55

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

56

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

57

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient

for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient

for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7)}(T_{37}, t)$, $+(a''_{36})^{(7,7)}(T_{37}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8)}(T_{41}, t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) \quad - (b''_{16})^{(2,2)}(G_{19}, t) \quad - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{28})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) \quad - (b''_{40})^{(8,8)}(G_{43}, t) \quad - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) \quad - (b''_{17})^{(2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{29})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) \quad - (b''_{41})^{(8,8)}(G_{43}, t) \quad - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} (b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t) \quad - (b''_{18})^{(2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4)}(G_{27}, t) \quad - (b''_{30})^{(5,5,5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7)}(G_{39}, t) \quad - (b''_{42})^{(8,8)}(G_{43}, t) \quad - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G, t)$, $-(b''_{14})^{(1)}(G, t)$, $-(b''_{15})^{(1)}(G, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} (G_{19}, t) & - (b''_{18})^{(2)} (G_{19}, t) & - (b'_{15})^{(1,1)} (G, t) & - (b''_{22})^{(3,3,3)} (G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)} (G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)} (G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)} (G_{35}, t) & \\ - (b''_{38})^{(7,7,7)} (G_{39}, t) & - (b''_{42})^{(8,8,8)} (G_{43}, t) & - (b''_{46})^{(9,9)} (G_{47}, t) & \end{array} \right] T_{18}$$

where $-(b'_{16})^{(2)} (G_{19}, t)$, $-(b'_{17})^{(2)} (G_{19}, t)$, $-(b'_{18})^{(2)} (G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)} (G, t)$, $-(b'_{14})^{(1,1)} (G, t)$, $-(b'_{15})^{(1,1)} (G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)} (G_{23}, t)$, $-(b''_{21})^{(3,3,3)} (G_{23}, t)$, $-(b''_{22})^{(3,3,3)} (G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)} (G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)} (G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)} (G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)} (G_{39}, t)$, $-(b''_{37})^{(7,7,7)} (G_{39}, t)$, $-(b''_{38})^{(7,7,7)} (G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)} (G_{43}, t)$, $-(b''_{41})^{(8,8,8)} (G_{43}, t)$, $-(b''_{42})^{(8,8,8)} (G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)} (G_{47}, t)$, $-(b''_{46})^{(9,9)} (G_{47}, t)$, $-(b''_{45})^{(9,9)} (G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} (T_{21}, t) & + (a''_{20})^{(3)} (T_{21}, t) & + (a'_{16})^{(2,2,2)} (T_{17}, t) & + (a'_{13})^{(1,1,1)} (T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)} (T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)} (T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)} (T_{33}, t) & \\ + (a''_{36})^{(7,7,7,7)} (T_{37}, t) & + (a''_{40})^{(8,8,8,8)} (T_{41}, t) & + (a''_{44})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} (T_{21}, t) & + (a''_{21})^{(3)} (T_{21}, t) & + (a'_{17})^{(2,2,2)} (T_{17}, t) & + (a'_{14})^{(1,1,1)} (T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)} (T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)} (T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)} (T_{33}, t) & \\ + (a''_{37})^{(7,7,7,7)} (T_{37}, t) & + (a''_{41})^{(8,8,8,8)} (T_{41}, t) & + (a''_{45})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{21}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} (T_{21}, t) & + (a''_{22})^{(3)} (T_{21}, t) & + (a'_{18})^{(2,2,2)} (T_{17}, t) & + (a'_{15})^{(1,1,1)} (T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)} (T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)} (T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)} (T_{33}, t) & \\ + (a''_{38})^{(7,7,7,7)} (T_{37}, t) & + (a''_{42})^{(8,8,8,8)} (T_{41}, t) & + (a''_{46})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{22}$$

$+(a'_{20})^{(3)} (T_{21}, t)$, $+(a'_{21})^{(3)} (T_{21}, t)$, $+(a'_{22})^{(3)} (T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)} (T_{17}, t)$, $+(a'_{17})^{(2,2,2)} (T_{17}, t)$, $+(a'_{18})^{(2,2,2)} (T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)} (T_{14}, t)$, $+(a'_{14})^{(1,1,1)} (T_{14}, t)$, $+(a'_{15})^{(1,1,1)} (T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b''_{16})^{(2,2,2)}(G_{19}, t) & -(b''_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b''_{17})^{(2,2,2)}(G_{19}, t) & -(b''_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b''_{18})^{(2,2,2)}(G_{19}, t) & -(b''_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b''_{30})^{(5,5,5)}(G_{31}, t) & -(b''_{26})^{(4,4,4)}(G_{27}, t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35}, t)$, $-(b''_{33})^{(6)}(G_{35}, t)$, $-(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & -(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{13})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & -(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{14})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & -(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & -(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & -(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1,1,1,1)}(G, t) & -(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{bmatrix} T_{15}$$

Where $-(b'_{36})^{(7)}(G_{39}, t)$, $-(b'_{37})^{(7)}(G_{39}, t)$, $-(b'_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$

95

$$= (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$\frac{dG_{41}}{dt}$

$$= (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$\frac{dG_{42}}{dt}$

$$= (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'|| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T'_{29}, t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(CCCCCCCC) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(DDDDDDDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(EEEEEEEEEE) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(FFFFFFFFF)$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(GGGGGGGGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(HHHHHHHHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \quad 169$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(jj) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$| (G_{19})^{(1)} - (G_{19})^{(2)} | e^{-(\widehat{M}_{16})^{(2)} t} \leq \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

$$\text{Then } \frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17} \text{ which leads to} \quad 204$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

$$\text{It is now sufficient to take } \frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1 \text{ and to choose} \quad 207$$

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \}$$

Indeed if we denote 212

Definition of $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $(\widetilde{G_{27}}, \widetilde{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} &\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), (G_{27})^{(2)}, (T_{27})^{(2)} \right) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3 :$ 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)} t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}, \widetilde{T}_{39}) : (\widetilde{G}_{39}, \widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)} s_{(36)}} e^{(\widetilde{M}_{36})^{(7)} s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) &d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ and $(\widehat{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$ and $((\widetilde{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}]$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t) :-$

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t) :-$

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,}$$

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain}$$

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(jj) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15})$, G_{13}, G_{15} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18})$, G_{16}, G_{18} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22})$, G_{20}, G_{22} must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)} (G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)} (G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)} (T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)} ((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)} ((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)} (T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)} ((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)} ((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)}T_{13}^*\mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)}T_{14}^*\mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)}T_{15}^*\mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)}T_{16}^*\mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^* T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^* T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^* T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^* T_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS 586

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
 & \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 & + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
 & \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
 & \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} (\lambda)^{(6)} \right) \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right) (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\ & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\ & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\ & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} (\lambda)^{(7)} \right) \\ & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} (\lambda)^{(7)} \right) \\ & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\ & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\ & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\ & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\ & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\ & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\ & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} (\lambda)^{(8)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \} \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY NINE

Entanglement Witnesses

INTRODUCTION—VARIABLES USED

Dariusz Chruściński and Gniewomir Sarbicki 2014 J. Phys A: Math. Theor 47 483001
doi:10.1088/1751-8113/47/48/483001 Entanglement witnesses: construction, analysis and classification
REVIEW ARTICLE

- (1) From the physical point of view entanglement witnesses (EWs) define (eb) a universal tool for analysis and classification (e&eb) of quantum entangled states.
- (2) From the mathematical perspective bipartite EWs provide (eb) highly non-trivial generalization of

positive operators and establish (eb) elegant correspondence with the theory of positive maps in (eb) matrix algebras.

- (3) Authors concentrate on theoretical analysis of various important notions like decomposability, atomicity, optimality, extremality and exposedness. Several methods of construction are provided as well. Discussion is illustrated by many examples enabling the reader to see the intricate structure of these objects. It is shown that the theory of EWs finds elegant geometric formulation in terms of (e&eb) convex cones and related geometric structures.

NOTATION

Module One

From the physical point of view

entanglement witnesses (EWs) define (eb) a universal tool for analysis and classification (e&eb) of quantum entangled states

G_{13} : Category one of entanglement witnesses (EWs)

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of universal tool for analysis and classification (e&eb) of quantum entangled states

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

entanglement witnesses (EWs) define a universal tool for analysis and classification (e&eb) of quantum entangled states

G_{16} : Category one of entanglement witnesses (EWs) define a universal tool; quantum entangled states

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of quantum entangled states ;entanglement witnesses (EWs) define a universal tool

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

From the mathematical perspective

bipartite EWs provide (eb) highly non-trivial generalization of positive operators and establish (eb) elegant correspondence with the theory of positive maps in (eb) matrix algebras

G_{20} : Category one of bipartite EWs

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of highly non-trivial generalization of positive operators and establish (eb) elegant correspondence with the theory of positive maps in (eb) matrix algebras

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

bipartite EWs provide highly non-trivial generalization of positive operators and establish (eb) elegant correspondence with the theory of positive maps in (eb) matrix algebras

G_{24} : Category one of bipartite EWs provide highly non-trivial generalization of positive operators

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of correspondence with the theory of positive maps in (eb) matrix algebras

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

bipartite EWs provide highly non-trivial generalization of positive operators and establish (eb) elegant correspondence with the theory of positive maps in matrix algebras

G_{28} : Category one of bipartite EWs; theory of positive maps in matrix algebras

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of theory of positive maps in matrix algebras ;bipartite EWs

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Authors concentrate on theoretical analysis of various important notions like

Decomposability, atomicity, optimality, extremality and exposedness.

Several methods of construction are provided as well.

Discussion is illustrated by many examples enabling the reader to see the intricate structure of these objects

It is shown that the

theory of EWs finds elegant geometric formulation in terms of (e&eb) convex cones and related geometric structures

G_{32} : Category one of Decomposability; atomicity,

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of atomicity,; Decomposability

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

G_{36} : Category one of atomicity; extremality and exposedness

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of extremality and exposedness ;atomicity

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

optimality, extremality and exposedness

G_{40} : Category one of optimality; extremality and exposedness

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of extremality and exposedness ;optimality

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

theory of EWs finds elegant geometric formulation in terms of (e&eb) convex cones and related geometric structures

G_{44} : Category one of theory of EWs; convex cones and related geometric structures

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of convex cones and related geometric structures; theory of EWs

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION FORTY

Chern–Simons–Landau–Ginzburg Theory

INTRODUCTION—VARIABLES USED

SHOU CHENG ZHANG, Int. J. Mod. Phys. B, 06, 25 (1992) DOI: 10.1142/S0217979292000037 THE CHERN–SIMONS–LANDAU–GINZBURG THEORY OF THE FRACTIONAL QUANTUM HALL EFFECT International Journal of Modern Physics B Condensed Matter Physics; Statistical Physics; Atomic, Molecular and Optical Physics

- (1) This paper gives a systematic review of a field theoretical approach to the fractional quantum Hall effect (FQHE) that has been developed in the past few years. Authors first illustrate some simple physical ideas to motivate such an approach and then present a systematic derivation of the Chern–Simons–Landau–Ginzburg (CSLG) action for the FQHE, starting from (e) the microscopic Hamiltonian.
- (2) It is demonstrated that all the phenomenological aspects of the FQHE can be derived from (e) the mean field solution and the small fluctuations of the CSLG action.
- (3) Although this formalism is logically independent of (e) Laughlin's wave function approach, their physical consequences are equivalent.
- (4) The CSLG theory demonstrates a deep connection between the phenomena of superfluidity and (e&eb) the FQHE, and can provide a simple and direct formalism to address many new macroscopic phenomena of the FQHE.

Martin Ammon et al 2013 J. Phys A: Math. Theor 46 214001 doi:10.1088/1751-8113/46/21/214001 Black holes in three dimensional higher spin gravity a review Martin Ammon, Michael Gutperle, Per Kraus and Eric Perlmutter

- (5) Authors review recent progress in the construction of black holes in (eb) three dimensional higher spin gravity theories.
- (6) Starting from spin-3 gravity and working our way toward the theory of an infinite tower of higher spins coupled to (e&eb) matter, they show (eb) how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes.
- (7) They review the construction of black holes with conserved higher spin charges and (e&eb) the computation of their partition functions to (e&eb) leading asymptotic order.
- (8) In view of the anti-de Sitter/conformal field theory (CFT) correspondence as applied to (e&eb) certain vector-like conformal field theories with extended conformal symmetry, they successfully compare to CFT calculations in a generalized Cardy regime. A brief recollection of pertinent aspects of ordinary gravity is also given. **This article is part of a special issue of Journal of**

Physics A: Mathematical and Theoretical devoted to 'Higher spin theories and holography'.

NOTATION

Module One

This paper gives a systematic review of a field theoretical approach to the fractional quantum Hall effect (FQHE) that has been developed in the past few years. Authors first illustrate some simple physical ideas to motivate such an approach and then present a systematic derivation of the

Chern–Simons–Landau–Ginzburg (CSLG) action for the FQHE, starting from (e) the microscopic Hamiltonian.

G_{13} : Category one of microscopic Hamiltonian.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Chern–Simons–Landau–Ginzburg (CSLG) action for the FQHE

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

It is demonstrated that

all the phenomenological aspects of the FQHE can be derived from (e) the mean field solution and the small fluctuations of the CSLG action

G_{16} : Category one of mean field solution and the small fluctuations of the CSLG action

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of all the phenomenological aspects of the FQHE

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Although this

formalism is logically independent of (e) Laughlin's wave function approach,

their physical consequences are equivalent.

G_{20} : Category one of Laughlin's wave function approach

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of formalism is logically independent

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

The

CSLG theory demonstrates a deep connection between the phenomena of superfluidity and (e&eb) the FQHE, and can provide a simple and direct formalism to address many new macroscopic phenomena of the FQHE

G_{24} : Category one of phenomena of superfluidity; FQHE

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of FQHE; phenomena of superfluidity

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Authors review recent progress in the construction of black holes in (eb) three dimensional higher spin gravity theories

G_{28} : Category one of construction of black holes

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of three dimensional higher spin gravity theories

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Starting from spin-3 gravity and working way toward the theory of an infinite tower of higher spins coupled to (e&eb) matter, they show (eb) how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes

G_{32} : Category one of theory of an infinite tower of higher spins; matter, they show (eb) how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of matter, they show (eb) how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes ;theory of an infinite tower of higher spins

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

theory of an infinite tower of higher spins coupled to matter, they show (eb) how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes

G_{36} : Category one of theory of an infinite tower of higher spins coupled to matter

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

theory of an infinite tower of higher spins coupled to matter, they show how to harness higher spin gauge invariance to consistently generalize (e&eb) familiar notions of black holes

G_{40} : Category one of theory of an infinite tower of higher spins coupled to matter, they show how to harness higher spin gauge invariance; notions of black holes

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of notions of black holes ;theory of an infinite tower of higher spins coupled to matter, they show how to harness higher spin gauge invariance

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

They review the

construction of black holes with conserved higher spin charges and (e&eb) the computation of their

partition functions to (e&eb) leading asymptotic order

G_{44} : Category one of construction of black holes with conserved higher spin charges; computation of their partition functions to (e&eb) leading asymptotic order

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of computation of their partition functions to (e&eb) leading asymptotic order; construction of black holes with conserved higher spin charges

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION FORTY ONE

Chiral Viscoelastic Response

INTRODUCTION—VARIABLES USED

Liang Sun and Shaolong Wan 2014 EPL 108 37007 doi:10.1209/0295-5075/108/37007 Chiral viscoelastic response in Weyl semimetals

- (1) Authors investigate the viscoelastic response of the electronic degrees of freedom in (eb) three-dimensional Weyl semimetals and obtain the effective action for this response.
- (2) This action is derived by adapting Fujikawa's method on chiral anomaly in curved space for (e) the most general Hamiltonian of Weyl semimetals.
- (3) **They clarify** that the Weyl semimetal phase will give rise to (eb) the nontrivial chiral viscoelastic effect due to (e) the perturbations of the tetrad field in the elastic media, which indicates (eb) that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response. And this nontrivial viscoelastic response can also be used to detect (eb) the properties of Weyl semimetals in the future..

Mitsutoshi Fujita et al JHEP06 (2009)066 doi:10.1088/1126-6708/2009/06/066 Fractional quantum Hall effect via holography: Chern-Simons, edge states and hierarchy

- (4) Authors present three holographic constructions of fractional quantum Hall effect (FQHE) via (e&eb) string theory.
- (5) The first model studies edge states in FQHE using (e) supersymmetric domain walls in $N = 6$ Chern-Simons theory.
- (6) They show that D4-branes wrapped on \mathbb{CP}^1 or D8-branes wrapped on \mathbb{CP}^3 create (eb) edge states that shift (e&eb) the rank or the level of the gauge group, respectively. These holographic edge states correctly reproduce the Hall conductivity.
- (7) The second model presents a holographic dual to (e) the pure $U(N)_k$ (Yang-Mills-)Chern-Simons theory based on (e) a D3-D7 system.
- (8) Its holography is equivalent to the (=) level-rank duality, which enables us (eb) to compute the Hall

conductivity and the topological entanglement entropy.

- (9) The third model introduces the first string theory embedding of (e) hierarchical FQHEs, using (e) IIA string on $\mathbb{C}^2/\mathbb{Z}_n$.

NOTATION

Module One

Authors investigate the viscoelastic response of the electronic degrees of freedom in (eb) three-dimensional Weyl semimetals and obtain the effective action for this response

G_{13} : Category one of viscoelastic response of the electronic degrees of freedom

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of three-dimensional Weyl semimetals and obtain the effective action for this response

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

This action is derived by adapting

Fujikawa's method on chiral anomaly in curved space for (e) the most general Hamiltonian of Weyl semimetals

G_{16} : Category one of adapting Fujikawa's method on chiral anomaly in curved space for (e) the most general Hamiltonian of Weyl semimetals

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of viscoelastic response of the electronic degrees of freedom in three-dimensional Weyl semimetals and obtain the effective action for this response

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Fujikawa's method on chiral anomaly in curved space for (e) the most general Hamiltonian of Weyl semimetals

G_{20} : Category one of most general Hamiltonian of Weyl semimetals

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Fujikawa's method on chiral anomaly in curved space

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

They clarify that the

Weyl semimetal phase will give rise to (eb) the nontrivial chiral viscoelastic effect due to (e) the perturbations of the tetrad field in the elastic media, which indicates (eb) that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response.

And this nontrivial viscoelastic response can also be used to detect (eb) the properties of Weyl semimetals in the future

G_{24} : Category one of Weyl semimetal phase

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of nontrivial chiral viscoelastic effect due to (e) the perturbations of the tetrad field in the elastic media, which indicates (eb) that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to (e) the perturbations of the tetrad field in the elastic media, which indicates (eb) that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

G_{28} : Category one of perturbations of the tetrad field in the elastic media, which indicates (eb) that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates (eb) that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

G_{32} : Category one of Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates that the crystal dislocations in Weyl semimetals will generate (eb) the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

G_{36} : Category one of Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates that the crystal dislocations in Weyl semimetals

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates that the crystal dislocations in Weyl semimetals will generate the energy-momentum current along the direction of dislocations which is (=) analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

G_{40} : Category one of Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates that the crystal dislocations in Weyl semimetals will generate the energy-momentum current along the direction of dislocations

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates that the crystal dislocations in Weyl semimetals will generate the energy-momentum current along the direction of dislocations which is analogous to the chiral magnetic effect predicted in (eb) electromagnetic response

G_{44} : Category one of Weyl semimetal phase will give rise to the nontrivial chiral viscoelastic effect due to the perturbations of the tetrad field in the elastic media, which indicates that the crystal dislocations in Weyl semimetals will generate the energy-momentum current along the direction of dislocations which is analogous to the chiral magnetic effect

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of electromagnetic response

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION FORTY TWO

Gravity Dual Of A Quantum Hall Plateau Transition

INTRODUCTION—VARIABLES USED

Joshua L. Davis et al JHEP11(2008)020 doi:10.1088/1126-6708/2008/11/020 Gravity dual of a quantum Hall plateau transition

- (1) Authors show how to model the transition between distinct quantum Hall plateaus in terms of (e&eb) D-branes in string theory.
- (2) A low energy theory of 2+1 dimensional fermions is obtained by (e) considering the D3-D7 system, and the plateau transition corresponds to (e&eb) moving the branes through one another.
- (3) They study the transition at strong coupling using (e) gauge/gravity duality and the probe approximation.

- (4) Strong coupling leads to (eb) a novel kind of plateau transition: at low temperatures the transition remains discontinuous due to (e) the effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out.
- (5) Strong coupling leads to (eb) a novel kind of plateau transition: at low temperatures the transition remains discontinuous due to (e) the effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out.

Clifford V. Johnson and Arnab Kundu JHEP12 (2008)053 doi:10.1088/1126-6708/2008/12/053
External fields and chiral symmetry breaking in the Sakai-Sugimoto model

- (6) Using the Sakai–Sugimoto model authors study the effect of an external magnetic field on (e&eb) the dynamics of fundamental flavours in (eb) both the confined and deconfined phases of a large N_c gauge theory.
- (7) They find that an external magnetic field promotes (eb) chiral symmetry breaking, consistent with the “magnetic catalysis” observed in the field theory literature, and seen in other studies using holographic duals.
- (8) The external field increases (eb) the separation between the deconfinement temperature and (e&eb) the chiral symmetry restoring temperature.
- (9) In the deconfined phase they investigate the temperature-magnetic field phase diagram and observe, for example, existence of (e) a maximum critical temperature (at which symmetry is restored) for (e) very large magnetic field.
- (10) They also find that this and certain other phenomena persist for (e) the Sakai–Sugimoto type models with (e&eb) probe branes of diverse dimensions.
- (11) They comment briefly on the dynamics in (eb) the presence of an external electric field

NOTATION

Module One

Authors show how to model the

transition between distinct quantum Hall plateaus in terms of (e&eb) D-branes in string theory

G_{13} : Category one of transition between distinct quantum Hall plateaus; D-branes in string theory

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of D-branes in string theory ;transition between distinct quantum Hall plateaus

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

A low energy theory of 2+1 dimensional fermions is obtained by (e) considering the D3-D7 system, and the plateau transition corresponds to (e&eb) moving the branes through one another

G_{16} : Category one of D3-D7 system, and the plateau transition corresponds to (e&eb) moving the branes through one another

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of low energy theory of 2+1 dimensional fermions

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

low energy theory of 2+1 dimensional fermions is obtained by considering the D3-D7 system, and the plateau transition corresponds to (e&eb) moving the branes through one another

G_{20} : Category one of low energy theory of 2+1 dimensional fermions is obtained by considering the D3-D7 system, and the plateau transition; moving the branes through one another

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of moving the branes through one another ;low energy theory of 2+1 dimensional fermions is obtained by considering the D3-D7 system, and the plateau transition

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

They study the

transition at strong coupling using (e) gauge/gravity duality and the probe approximation

G_{24} : Category one of transition at strong coupling; gauge/gravity duality and the probe approximation

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of gauge/gravity duality and the probe approximation; transition at strong coupling

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Strong coupling leads to (eb) a novel kind of **plateau transition**: at low temperatures the transition remains discontinuous due to (e) the effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out

G_{28} : Category one of Strong coupling

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of novel kind of plateau transition: at low temperatures the transition remains discontinuous due to (e) the effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Strong coupling leads to a novel kind of plateau transition:

transition remains discontinuous at low temperatures due to (e) the effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out

G_{32} : Category one of effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Strong coupling leads to a novel kind of plateau transition: at low temperatures the transition remains discontinuous

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

transition remains discontinuous at low temperatures due to (e) the effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out

G_{36} : Category one of effects of dynamical symmetry breaking and mass generation and at high temperatures is only partially smoothed out

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of transition remains discontinuous at low temperatures

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

authors study the

effect of an external magnetic field Using the Sakai–Sugimoto model on (e&eb) the dynamics of fundamental flavours in (eb) both the confined and deconfined phases of a large N_c gauge theory

G_{40} : Category one of effect of an external magnetic field Using the Sakai–Sugimoto model

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of dynamics of fundamental flavours in (eb) both the confined and deconfined phases of a large N_c gauge theory

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

effect of an external magnetic field Using the Sakai–Sugimoto model on the dynamics of fundamental flavours in (eb) both the confined and deconfined phases of a large N_c gauge theory

G_{44} : Category one of effect of an external magnetic field Using the Sakai–Sugimoto model on the dynamics of fundamental flavours

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of confined and deconfined phases of a large N_c gauge theory

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)};$
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)] G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)] G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)] G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)} T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)] T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)} T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)] T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)} T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)] T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)] G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)] G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)] G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)] T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)] T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)] T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)] G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)] G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)] G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)] T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)] T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)} T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)] T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)} G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)] G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)} G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)] G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)} G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)] G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)} T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)] T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)} T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)] T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)] T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)] G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)] G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)] G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)] T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)] T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)] T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)] T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)] T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

42

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40}$$

43

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41}$$

44

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42}$$

45

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40}$$

46

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41}$$

47

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42}$$

48

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44}$$

49

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45}$$

50

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46}$$

51

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44}$$

52

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45}$$

53

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46}$$

54

$$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$$

$$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

55

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

56

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

57

Where $(a''_{13})^{(1)}(T_{14}, t)$, $(a''_{14})^{(1)}(T_{14}, t)$, $(a''_{15})^{(1)}(T_{14}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$, $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b''_{37})^{(7,7)}(G_{39}, t)}$, $\boxed{-(b''_{38})^{(7,7)}(G_{39}, t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43}, t)}$, $\boxed{-(b''_{41})^{(8,8)}(G_{43}, t)}$, $\boxed{-(b''_{42})^{(8,8)}(G_{43}, t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) & \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) & \end{array} \right] G_{18} \quad 63$$

Where $+(a'_{16})^{(2)}(T_{17}, t)$, $+(a'_{17})^{(2)}(T_{17}, t)$, $+(a'_{18})^{(2)}(T_{17}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1)}(T_{14}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8)}(T_{41}, t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9)}(T_{45}, t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) & \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) & \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} (G_{19}, t) & -(b''_{18})^{(2)} (G_{19}, t) & -(b'_{15})^{(1,1)} (G, t) & -(b''_{22})^{(3,3,3)} (G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4)} (G_{27}, t) & -(b''_{30})^{(5,5,5,5,5)} (G_{31}, t) & -(b''_{34})^{(6,6,6,6,6)} (G_{35}, t) & \\ -(b''_{38})^{(7,7,7)} (G_{39}, t) & -(b''_{42})^{(8,8,8)} (G_{43}, t) & -(b''_{46})^{(9,9)} (G_{47}, t) & \end{array} \right] T_{18}$$

where $-(b'_{16})^{(2)} (G_{19}, t)$, $-(b'_{17})^{(2)} (G_{19}, t)$, $-(b'_{18})^{(2)} (G_{19}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1)} (G, t)$, $-(b'_{14})^{(1,1)} (G, t)$, $-(b'_{15})^{(1,1)} (G, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)} (G_{23}, t)$, $-(b''_{21})^{(3,3,3)} (G_{23}, t)$, $-(b''_{22})^{(3,3,3)} (G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)} (G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)} (G_{27}, t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)} (G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)} (G_{31}, t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)} (G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)} (G_{35}, t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)} (G_{39}, t)$, $-(b''_{37})^{(7,7,7)} (G_{39}, t)$, $-(b''_{38})^{(7,7,7)} (G_{39}, t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)} (G_{43}, t)$, $-(b''_{41})^{(8,8,8)} (G_{43}, t)$, $-(b''_{42})^{(8,8,8)} (G_{43}, t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)} (G_{47}, t)$, $-(b''_{46})^{(9,9)} (G_{47}, t)$, $-(b''_{45})^{(9,9)} (G_{47}, t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} (T_{21}, t) & +(a''_{20})^{(3)} (T_{21}, t) & +(a'_{16})^{(2,2,2)} (T_{17}, t) & +(a'_{13})^{(1,1,1)} (T_{14}, t) \\ +(a''_{24})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{28})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{32})^{(6,6,6,6,6)} (T_{33}, t) & \\ +(a''_{36})^{(7,7,7,7)} (T_{37}, t) & +(a''_{40})^{(8,8,8,8)} (T_{41}, t) & +(a''_{44})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} (T_{21}, t) & +(a''_{21})^{(3)} (T_{21}, t) & +(a'_{17})^{(2,2,2)} (T_{17}, t) & +(a'_{14})^{(1,1,1)} (T_{14}, t) \\ +(a''_{25})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{29})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{33})^{(6,6,6,6,6)} (T_{33}, t) & \\ +(a''_{37})^{(7,7,7,7)} (T_{37}, t) & +(a''_{41})^{(8,8,8,8)} (T_{41}, t) & +(a''_{45})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{21}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} (T_{21}, t) & +(a''_{22})^{(3)} (T_{21}, t) & +(a'_{18})^{(2,2,2)} (T_{17}, t) & +(a'_{15})^{(1,1,1)} (T_{14}, t) \\ +(a''_{26})^{(4,4,4,4,4)} (T_{25}, t) & +(a''_{30})^{(5,5,5,5,5)} (T_{29}, t) & +(a''_{34})^{(6,6,6,6,6)} (T_{33}, t) & \\ +(a''_{38})^{(7,7,7,7)} (T_{37}, t) & +(a''_{42})^{(8,8,8,8)} (T_{41}, t) & +(a''_{46})^{(9,9,9)} (T_{45}, t) & \end{array} \right] G_{22}$$

$+(a'_{20})^{(3)} (T_{21}, t)$, $+(a'_{21})^{(3)} (T_{21}, t)$, $+(a'_{22})^{(3)} (T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2)} (T_{17}, t)$, $+(a'_{17})^{(2,2,2)} (T_{17}, t)$, $+(a'_{18})^{(2,2,2)} (T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a'_{13})^{(1,1,1)} (T_{14}, t)$, $+(a'_{14})^{(1,1,1)} (T_{14}, t)$, $+(a'_{15})^{(1,1,1)} (T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41}, t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9)}(T_{45}, t)$, $+(a''_{46})^{(9,9,9)}(T_{45}, t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}[-(b''_{20})^{(3)}(G_{23}, t)] & -(b''_{16})^{(2,2,2)}(G_{19}, t) & -(b''_{13})^{(1,1,1)}(G, t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39}, t) & -(b''_{40})^{(8,8,8,8)}(G_{43}, t) & -(b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)}[-(b''_{21})^{(3)}(G_{23}, t)] & -(b''_{17})^{(2,2,2)}(G_{19}, t) & -(b''_{14})^{(1,1,1)}(G, t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39}, t) & -(b''_{41})^{(8,8,8,8)}(G_{43}, t) & -(b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)}[-(b''_{22})^{(3)}(G_{23}, t)] & -(b''_{18})^{(2,2,2)}(G_{19}, t) & -(b''_{15})^{(1,1,1)}(G, t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39}, t) & -(b''_{42})^{(8,8,8,8)}(G_{43}, t) & -(b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$, $-(b''_{21})^{(3)}(G_{23}, t)$, $-(b''_{22})^{(3)}(G_{23}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$, $(a''_{25})^{(4)}(T_{25}, t)$, $(a''_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$ are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)} T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27}, t) & -(b''_{30})^{(5,5)}(G_{31}, t) & -(b''_{34})^{(6,6)}(G_{35}, t) \\ -(b'_{15})^{(1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{26}$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b'_{36})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b'_{40})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b'_{46})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29}, t) & +(a'_{24})^{(4,4)}(T_{25}, t) & +(a'_{32})^{(6,6,6)}(T_{33}, t) \\ +(a'_{13})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{16})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{36})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{40})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{28}$$

79

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29}, t) & +(a'_{25})^{(4,4)}(T_{25}, t) & +(a'_{33})^{(6,6,6)}(T_{33}, t) \\ +(a'_{14})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{17})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{37})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{41})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{29}$$

80

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29}, t) & +(a'_{26})^{(4,4)}(T_{25}, t) & +(a'_{34})^{(6,6,6)}(T_{33}, t) \\ +(a'_{15})^{(1,1,1,1,1)}(T_{14}, t) & +(a'_{18})^{(2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3)}(T_{21}, t) & \\ +(a'_{38})^{(7,7,7,7,7)}(T_{37}, t) & +(a'_{42})^{(8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{30}$$

81

Where $+(a'_{28})^{(5)}(T_{29}, t)$, $+(a'_{29})^{(5)}(T_{29}, t)$, $+(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a'_{24})^{(4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \text{ are fifth augmentation}$$

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \text{ are sixth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) \text{ are seventh augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) \text{ are eighth augmentation}$$

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \text{ are ninth augmentation}$$

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31}, t)$, $-(b''_{29})^{(5)}(G_{31}, t)$, $-(b''_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33}$$

$$- \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$

seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b'_{30})^{(5,5,5)}(G_{31}, t) & -(b'_{26})^{(4,4,4)}(G_{27}, t) \\ -(b'_{15})^{(1,1,1,1,1,1)}(G, t) & -(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ -(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & -(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & -(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t) & \end{array} \right] T_{34} \quad 90$$

$-(b'_{32})^{(6)}(G_{35}, t)$, $-(b'_{33})^{(6)}(G_{35}, t)$, $-(b'_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients
for category 1, 2 and 3

$-(b'_{28})^{(5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients
for category 1, 2 and 3

$-(b'_{24})^{(4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients
for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients
for category 1, 2, and 3

$-(b'_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition
coefficients for category 1, 2, and 3

$-(b'_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition
coefficients for category 1, 2, and 3

$-(b'_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ are seventh detrition
coefficients for category 1, 2, and 3

$-(b'_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$
are eighth detrition coefficients for category 1, 2, and 3

$-(b'_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$, $-(b'_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37}, t) & +(a'_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{44})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[\begin{array}{ccc} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37}, t) & +(a'_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{45})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \left[\begin{array}{ccc} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37}, t) & +(a'_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & +(a'_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ +(a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & +(a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & +(a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ +(a'_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & +(a'_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & +(a'_{46})^{(9,9,9,9,9,9)}(T_{45}, t) & \end{array} \right] G_{15}$$

Where $(a'_{36})^{(7)}(T_{37}, t)$, $(a'_{37})^{(7)}(T_{37}, t)$, $(a'_{38})^{(7)}(T_{37}, t)$ are first augmentation coefficients for
category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \left[\begin{array}{ccc} \boxed{(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \left[\begin{array}{ccc} \boxed{(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$, $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$

are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} \quad 95$$

$$- \left[\begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where $+(a'_{40})^{(8)}(T_{41}, t)$, $+(a'_{41})^{(8)}(T_{41}, t)$, $+(a'_{42})^{(8)}(T_{41}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39}, t)$, $-(b''_{37})^{(7)}(G_{39}, t)$, $-(b''_{38})^{(7)}(G_{39}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b''_{38})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) & + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) & + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) & + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) & + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where $+(a'_{44})^{(9)}(T_{45}, t)$, $+(a'_{45})^{(9)}(T_{45}, t)$, $+(a'_{46})^{(9)}(T_{37}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$, $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$, $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$, $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$, $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$, $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$, $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$, $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$, $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} -$$

$$\begin{aligned}
 & \left[\begin{array}{ccc} (b'_{44})^{(9)} \left[- (b''_{44})^{(9)}(G_{47}, t) \right] & - (b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\
 \frac{dT_{45}}{dt} = & \\
 (b_{45})^{(9)} T_{44} - & \left[\begin{array}{ccc} (b'_{45})^{(9)} \left[- (b''_{45})^{(9)}(G_{47}, t) \right] & - (b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\
 \frac{dT_{46}}{dt} = & \\
 (b_{46})^{(9)} T_{45} - & \left[\begin{array}{ccc} (b'_{46})^{(9)} \left[- (b''_{46})^{(9)}(G_{47}, t) \right] & - (b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15}
 \end{aligned}$$

Where $-(b'_{44})^{(9)}(G_{47}, t)$, $-(b'_{45})^{(9)}(G_{47}, t)$, $-(b'_{46})^{(9)}(G_{47}, t)$ are first detrition coefficients for category 1, 2 and 3
 $-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ are second detrition coefficients for category 1, 2 and 3
 $-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1, 2 and 3
 $-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$, $-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ are fourth detrition coefficients for category 1, 2 and 3
 $-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ are fifth detrition coefficients for category 1, 2 and 3
 $-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)$ are sixth detrition coefficients for category 1, 2 and 3
 $-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$ are seventh detrition coefficients for category 1, 2 and 3
 $-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$, $-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ are eighth detrition coefficients for category 1, 2 and 3
 $-(b'_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$, $-(b'_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)$ are ninth detrition coefficients for category 1, 2 and 3
Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

97

The functions $(a'_i)^{(1)}, (b'_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a'_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T'_{21}, t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T'_{25}, t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \quad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T_{29}'| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||G_{31} - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||G_{35} - (G_{35})'| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(IIIIIIII) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38 \quad 131$$

(JJJJJJJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(KKKKKKKKKK) \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$(LLLLLLLLLL)$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$:

134

(MMMMMMMMMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$:

135

(NNNNNNNNNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$ and $(\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions $(a''_i)^{(8)}, (b''_i)^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(8)}(T'_{41}, t)$ and $(a'_i)^{(8)}(T_{41}, t)$. (T'_{41}, t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a''_i)^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a''_i)^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146
A

The functions $(a''_i)^{(9)}, (b''_i)^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, (r_i)^{(9)}$:

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$ are positive constants and $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(9)}(T'_{45}, t)$ and $(a''_i)^{(9)}(T_{45}, t)$. (T'_{45}, t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a''_i)^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a''_i)^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ and $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

A

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions

153

B

Definition of $G_i(0), T_i(0)$:

$$\begin{aligned} G_i(t) &\leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{G_i(0) = G_i^0 > 0} \\ T_i(t) &\leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}, \quad \boxed{T_i(0) = T_i^0 > 0} \end{aligned}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + (a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + (a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{P}_{13})^{(1)} + G_{14}^0) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[((\hat{P}_{24})^{(4)} + G_{25}^0) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(kk) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)} t) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}} + (\hat{P}_{44})^{(9)} \right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \}$$

Indeed if we denote

Definition of \tilde{G}, \tilde{T} : $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + \\ G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \quad 186$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$ and $((\widehat{M}_{13})^{(1)})_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})$ it follows $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$.

189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

191

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

192

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

194

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)} t} \}$$

Indeed if we denote

195

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{(\bar{M}_{16})^{(2)} s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)} s_{(16)}} e^{-(\bar{M}_{16})^{(2)} s_{(16)}} +$$

$$(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) | G_{16}^{(1)} - G_{16}^{(2)} | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} \} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)} t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$ respectively of \mathbb{R}_+ . 199

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \rightarrow \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} &\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned} \quad 214$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ and $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of \mathbb{R}_+ . 215

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1$, $((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b_{22}'')^{(3)}((G_{23})(t), t)) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose 221

$(\bar{P}_{24})^{(4)}$ and $(\bar{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\bar{P}_{24})^{(4)} + ((\bar{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\bar{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\bar{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\bar{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\bar{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\bar{Q}_{24})^{(4)} \right] \leq (\bar{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

Definition of $(\bar{G}_{27}), (\bar{T}_{27})$: $(\bar{G}_{27}), (\bar{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)} t} &\leq \\ \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) &d((G_{27})^{(1)}, (T_{27})^{(1)}; (G_{27})^{(2)}, (T_{27})^{(2)}) \end{aligned} \quad 226$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of \mathbb{R}_+ . 227

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}\} (T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)}) \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \rightarrow \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)} t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) &d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned} \quad 237$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of \mathbb{R}_+ . 238

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[(\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \}$$

Indeed if we denote

Definition of $(\widehat{G_{35}}, \widehat{T_{35}})$: $(\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \end{aligned} \quad 248$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a'_{32})^{(6)}$ and $(b'_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of \mathbb{R}_+ . 249

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose $(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[(\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d \left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : (\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{-(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} + \\ &G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)} s_{(36)}} e^{(\bar{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)} t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} + (\bar{P}_{36})^{(7)} (\bar{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned} \quad 259$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ and $(\bar{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$ respectively of \mathbb{R}_+ . 260

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)} t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < (\bar{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \rightarrow \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for T_{38} if $\lim_{t \rightarrow \infty} (b''_{38})^{(7)} ((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[(\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_{i \in \mathbb{R}_+} \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \} \quad 269$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}, \widetilde{T_{43}})$: $(\widetilde{G_{43}}, \widetilde{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)} \end{aligned}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)}) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) \end{aligned} \quad 273$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ and $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$ respectively of \mathbb{R}_+ . 274

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}$, $i = 40, 41, 42$ depend only on T_{41} and respectively on (G_{43}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{40})^{(8)})_1, ((\bar{M}_{40})^{(8)})_2$ and $((\bar{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} and G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If G_{41} or G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} is bounded, from below, the same property holds for G_{41} and G_{42} . The proof is 277
analogous with the preceding one. An analogous property is true if G_{41} is bounded from below.

Remark 40: If T_{40} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \rightarrow \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for T_{42} if $\lim_{t \rightarrow \infty} (b''_{42})^{(8)}((G_{43})(t), t(t), t) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279
A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[(\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} |(a'_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a'_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\bar{A}_{44})^{(9)} + (\bar{P}_{44})^{(9)} (\bar{k}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\bar{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ and $(\bar{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(9)}$ and $(b'_i)^{(9)}$, $i = 44, 45, 46$ depend only on T_{45} and respectively on (G_{47}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

Definition of $((\bar{M}_{44})^{(9)})_1, ((\bar{M}_{44})^{(9)})_2$ and $((\bar{M}_{44})^{(9)})_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)})_1 - (a'_{45})^{(9)} G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$ then $T_{45} \rightarrow \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for T_{46} if $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation

280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$:

281

By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$:

282

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:-

283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right)$$

$$T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)} (m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation

291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$:

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$:

295

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots

296

of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and

298

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$:

299

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the

300

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:-

303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

$$\text{and } (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

320

By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the

roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$:

361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$:

362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \quad 366$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$ and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^0e^{-(a'_{42})^{(8)}t}$$

$$T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^0e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}]+T_{42}^0e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}]+T_{42}^0e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$ the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

Definition of $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$ the

roots of the equations $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof: From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:-
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

$$\text{For } 0 < \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right] < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \left[(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}} \right]$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \left[(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}} \right]$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \left[(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} \right]$, we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

Proof : From global equations we obtain

398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

$$\text{it follows } (v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

$$\text{If } 0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)} \text{ we find like in the previous case,} \quad 402$$

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

$$\text{If } 0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} , \text{ we obtain} \quad 403$$

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

Proof: From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines** $(v_0)^{(4)}$ **for the special case**.

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 409

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)} (\nu_2)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((\nu_1)^{(5)} - (\nu_2)^{(5)}) t]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)} (\bar{\nu}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}) t]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}''^{(5)} = (a_{29}''^{(5)})$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines $(\nu_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, and definition of $(u_0)^{(5)}$.

Proof: From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

417

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:-
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)}e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)}e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}$, then $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines $(v_0)^{(8)}$ for the special case .**

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}$, then $(\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of $(u_0)^{(8)}$.**

Proof : From 99,20,44,22,23,44 we obtain

424
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left((a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)} (\nu_2)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)} ((\nu_1)^{(9)} - (\nu_2)^{(9)}) t]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)} (\bar{\nu}_2)^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)} ((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}) t]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}$, then $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines $(\nu_0)^{(9)}$ for the special case.**

Analogously if $(b_{44}'')^{(9)} = (b_{45}'')^{(9)}$, then $(\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then

$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$ if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of $(u_0)^{(9)}$.**

We can prove the following

425

Theorem : If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a_{13}')^{(1)}(a_{14}')^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a_{14}')^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b_{13}')^{(1)}(r_{14})^{(1)} - (b_{14}')^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions with the notations

426

$$(a_{16}')^{(2)}(a_{17}')^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied, then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied, then the system

Theorem : If $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied, then the system

Theorem If $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with $(p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436
45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0$	479
$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0$	480
$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0$	481
$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0$	482
$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0$	483
$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0$	484
$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0$	484
$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$	A
$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$	
$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$	
$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$	
$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$	

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(kk) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof:

489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof:

490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof:

491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof:

492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof:

492

A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :-

493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :-

494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

Definition and uniqueness of T_{21}^* :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

Definition and uniqueness of T_{25}^* :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a_{40})^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a_{42})^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :-

501
A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44})^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46})^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b'_{14})^{(1)}(G) + (b'_{14})^{(1)}(b'_{13})^{(1)}(G)] + (b'_{13})^{(1)}(G)(b'_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b'_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b'_{16})^{(2)}(G_{19})] + (b'_{16})^{(2)}(G_{19})(b'_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions G_{20}, G_{21} if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b'_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b'_{20})^{(3)}(G_{23})] + (b'_{20})^{(3)}(G_{23})(b'_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
 $[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
 $[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)} ((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)} ((G_{31})^*)]}$$

519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)} (T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)} ((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)} ((G_{39})^*)]}$$

Finally we obtain the unique solution

523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)} ((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)} ((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99

523

A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)} ((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)} ((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS

524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)} T_{13}^* \mathbb{G}_j) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)} T_{14}^* \mathbb{G}_j) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)} T_{15}^* \mathbb{G}_j) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)} T_{16}^* \mathbb{G}_j) \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)} T_{17}^* \mathbb{G}_j) \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)} T_{18}^* \mathbb{G}_j) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)} T_{20}^* \mathbb{G}_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)} T_{21}^* \mathbb{G}_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)} T_{22}^* \mathbb{G}_j) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of \mathbb{G}_i, T_i :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*\mathbb{G}_j) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*\mathbb{G}_j) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*\mathbb{G}_j) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)}T_{28}^*\mathbb{G}_j) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)}T_{29}^*\mathbb{G}_j) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)}T_{30}^*\mathbb{G}_j) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)}T_{32}^*G_j) \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)}T_{33}^*G_j) \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)}T_{34}^*G_j) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)}T_{36}^*G_j) \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)}T_{37}^*G_j) \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)}T_{38}^*G_j) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)}T_{40}^*G_j) \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)}T_{41}^*G_j) \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)}T_{42}^*G_j) \quad 586$$

ASYMPTOTIC STABILITY ANALYSIS 586

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ and $(b_i'')^{(9)}$ belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{d\mathbb{G}_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45}$$

586
C

$$\frac{d\mathbb{G}_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45}$$

586
D

$$\frac{d\mathbb{T}_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(44)(j)}T_{44}^*\mathbb{G}_j)$$

586
E

$$\frac{d\mathbb{T}_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}(s_{(45)(j)}T_{45}^*\mathbb{G}_j)$$

586
F

$$\frac{d\mathbb{T}_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}(s_{(46)(j)}T_{46}^*\mathbb{G}_j)$$

586
G

The characteristic equation of this system is

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \\ &+ ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \\ &((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \\ &((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \\ &+ ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) ((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*) \\ &((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\ &+ ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\ &((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\ &((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \} \end{aligned}$$

$$\begin{aligned}
& \left(((\lambda)^{(2)})^2 + (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
& + \left(((\lambda)^{(2)})^2 + (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
& + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^*) \\
& \left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0 \\
& + \\
& ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)}) \{ ((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
& \left[((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
& \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
& + \left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)}) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \right) \\
& \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right) \\
& \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
& \left(((\lambda)^{(3)})^2 + (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
& + \left(((\lambda)^{(3)})^2 + (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
& + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^*) \\
& \left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)}) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0 \\
& + \\
& ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)}) \{ ((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
& \left[((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
& \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right) \\
& + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)}) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right) \\
& \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right) \\
& \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)}
\end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)})^2 + (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^*) \\
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)}) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\
 & + \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} (\lambda)^{(6)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\
 & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}) \{ ((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \\
 & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\
 & \left(((\lambda)^{(7)})^2 + (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\
 & + \left(((\lambda)^{(7)})^2 + (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^*) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \} = 0 \\
 & + \\
 & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)}) \{ ((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 & \left[((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \\
 & + \left(((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)}) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \\
 & \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^*) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(8)})^2 + (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} \right) (\lambda)^{(8)} \\
 & + \left(((\lambda)^{(8)})^2 + (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} \right) (\lambda)^{(8)} (q_{42})^{(8)} G_{42} \\
 & + \left((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left((a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left(((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right) \} = 0 \\
 & + \\
 & \left((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \{ ((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 & \left[\left(((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \right] \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left(((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)}) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & \left(((\lambda)^{(9)})^2 + (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} \right) (\lambda)^{(9)} \\
 & + \left(((\lambda)^{(9)})^2 + (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} \right) (\lambda)^{(9)} (q_{46})^{(9)} G_{46} \\
 & + \left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left((a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left(((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

References

- (1) Bousso, Raphael (2002). "The holographic principle". *Reviews of Modern Physics* 74 (3): 825–874. arXiv:hep-th/0203101. Bibcode:2002RvMP...74..825B. doi:10.1103/RevModPhys.74.825.
- (2) 't Hooft, Gerard (1993). "Dimensional Reduction in Quantum Gravity". p. 10026. arXiv:gr-qc/9310026. Bibcode:1993gr.qc....10026T.. 't Hooft's original paper.
- (3) ^ Jump up to: a b Susskind, Leonard (1995). "The World as a Hologram". *Journal of Mathematical Physics* 36 (11): 6377–6396. arXiv:hep-th/9409089. Bibcode:1995JMP....36.6377S. doi:10.1063/1.531249.
- (4) Jump up ^ Thorn, Charles B. (27–31 May 1991). "Reformulating string theory with the 1/N expansion". *International A.D. Sakharov Conference on Physics. Moscow*. pp. 447–54. arXiv:hep-th/9405069. ISBN 978-1-56072-073-7.
- (5) Jump up ^ Bousso, Raphael (2002). "The Holographic Principle". *Reviews of Modern Physics* 74 (3): 825–874. arXiv:hep-th/0203101. Bibcode:2002RvMP...74..825B. doi:10.1103/RevModPhys.74.825
- (6) Jump up ^ Lloyd, Seth (2002-05-24). "Computational Capacity of the Universe". *Physical Review Letters* 88 (23): 237901. arXiv:quant-ph/0110141. Bibcode:2002PhRvL..88w7901L. doi:10.1103/PhysRevLett.88.237901. PMID 12059399.
- (7) Jump up ^ Susskind, L. (2008). *The Black Hole War – My Battle with Stephen Hawking to Make the World Safe for Quantum Mechanics*. Little, Brown and Company
- (8) Jump up ^ Marolf, Donald (April 2009). "Black Holes, AdS, and CFTs". *General Relativity and Gravitation* 41 (4): 903–17. Bibcode:2009GReGr..41..903M. doi:10.1007/s10714-008-0749-7.
- (9) Qingming Cheng (2001), "De Sitter space", in Hazewinkel, Michiel, *Encyclopedia of Mathematics*, Springer, ISBN 978-1-55608-010-4
- (10) de Sitter, W. (1917), "On the relativity of inertia: Remarks concerning Einstein's latest hypothesis", *Proc. Kon. Ned. Acad. Wet.* 19: 1217–1225
- (11) de Sitter, W. (1917), "On the curvature of space", *Proc. Kon. Ned. Acad. Wet.* 20: 229–243
- (12) Nomizu, Katsumi (1982), "The Lorentz–Poincaré metric on the upper half-space and its extension", *Hokkaido Mathematical Journal* 11 (3): 253–261
- (13) Coxeter, H. S. M. (1943), "A geometrical background for de Sitter's world", *American Mathematical Monthly (Mathematical Association of America)* 50 (4): 217–228, doi:10.2307/2303924, JSTOR 2303924
- (14) Susskind, L.; Lindesay, J. (2005), *An Introduction to Black Holes, Information and the String Theory Revolution: The Holographic Universe*, p. 119(11.5.25)
- (15) Coussaert, Oliver; Henneaux, Marc; van Driel, Peter (1995). "The asymptotic dynamics of three-dimensional Einstein gravity with a negative cosmological constant". *Classical and Quantum Gravity* 12 (12): 2961. arXiv:gr-qc/9506019. Bibcode:1995CQGr..12.2961C. doi:10.1088/0264-9381/12/12/012.
- (16) Cowen, Ron (2013). "Simulations back up theory that Universe is a hologram". *Nature News & Comment*. doi:10.1038/nature.2013.14328
- (17) de Haro, Sebastian; Dieks, Dennis; 't Hooft, Gerard; Verlinde, Erik (2013). "Forty Years of String Theory Reflecting on the Foundations". *Foundations of Physics* 43 (1): 1–7. Bibcode:2013FoPh...43....1D. doi:10.1007/s10701-012-9691-3.
- (18) Deligne, Pierre; Etingof, Pavel; Freed, Daniel; Jeffery, Lisa; Kazhdan, David; Morgan, John; Morrison, David; Witten, Edward, eds. (1999). *Quantum Fields and Strings: A Course for Mathematicians*. American Mathematical Society. ISBN 978-0-8218-2014-8.
- (19) Fefferman, Charles; Graham, Robin (1985). "Conformal invariants". *Asterisque*: 95–116.
- (20) Fefferman, Charles; Graham, Robin (2011). *The Ambient Metric*. Princeton University Press. ISBN 978-1-4008-4058-8.
- (21) Giombi, Simone; Yin, Xi (2010). "Higher spin gauge theory and holography: the three-point functions". *Journal of High Energy Physics* 2010 (9): 1–80. arXiv:0912.3462. Bibcode:2010JHEP...09..115G. doi:10.1007/JHEP09(2010)115.
- (22) Greene, Brian (2000). *The Elegant Universe: Superstrings, Hidden Dimensions, and the Quest for the Ultimate Theory*. Random House. ISBN 978-0-9650888-0-0.

- (23) Griffiths, David (2004). Introduction to Quantum Mechanics. Pearson Prentice Hall. ISBN 978-0-13-111892-8.
- (24) Gubser, Steven; Klebanov, Igor; Polyakov, Alexander (1998). "Gauge theory correlators from non-critical string theory". Physics Letters B 428: 105–114. arXiv:hep-th/9802109. Bibcode:1998PhLB..428..105G. doi:10.1016/S0370-2693(98)00377-3.
- (25) Guica, Monica; Hartman, Thomas; Song, Wei; Strominger, Andrew (2009). "The Kerr/CFT Correspondence". Physical Review D 80 (12). arXiv:0809.4266. Bibcode:2009PhRvD..80l4008G. doi:10.1103/PhysRevD.80.124008.
- (26) Hawking, Stephen (1975). "Particle creation by black holes". Communications in mathematical physics 43 (3): 199–220. Bibcode:1975CMaPh..43..199H. doi:10.1007/BF02345020.
- (27) Hawking, Stephen (2005). "Information loss in black holes". Physical Review D 72 (8). arXiv:hep-th/0507171. Bibcode:2005PhRvD..72h4013H. doi:10.1103/PhysRevD.72.084013.
- (28) Klebanov, Igor; Maldacena, Juan (2009). "Solving Quantum Field Theories via Curved Spacetimes" (PDF). Physics Today 62: 28. Bibcode:2009PhT....62a..28K. doi:10.1063/1.3074260. Retrieved May 2013.
- (29) Klebanov, Igor; Polyakov, Alexander (2002). "The AdS dual of the critical O(N) vector model". Physics Letters B 550 (3–4): 213–219. arXiv:hep-th/0210114. Bibcode:2002PhLB..550..213K. doi:10.1016/S0370-2693(02)02980-5.
- (30) Kovtun, P. K.; Son, Dam T.; Starinets, A. O. (2001). "Viscosity in strongly interacting quantum field theories from black hole physics". Physical review letters 94 (11): 111601. arXiv:hep-th/0405231. Bibcode:2005PhRvL..94k1601K. doi:10.1103/PhysRevLett.94.111601. PMID 15903845.
- (31) Luzum, Matthew; Romatschke, Paul (2008). "Conformal relativistic viscous hydrodynamics: Applications to RHIC results at $\sqrt{s_{NN}}=200$ GeV". Physical Review C 78 (3). arXiv:0804.4015. doi:10.1103/PhysRevC.78.034915.
- (32) Maldacena, Juan (1998). "The Large N limit of superconformal field theories and supergravity". Advances in Theoretical and Mathematical Physics 2: 231–252. arXiv:hep-th/9711200. Bibcode:1998AdTMP...2..231M. doi:10.1063/1.59653.
- (33) Maldacena, Juan (2005). "The Illusion of Gravity". Scientific American 293 (5): 56–63. Bibcode:2005SciAm.293e..56M. doi:10.1038/scientificamerican1105-56. PMID 16318027. Retrieved July 2013.
- (34) McLerran, Larry (2007). "Theory Summary : Quark Matter 2006". Journal of Physics G: Nuclear and Particle Physics 34 (8): S583. arXiv:hep-ph/0702004. Bibcode:2007JPhG...34..583M. doi:10.1088/0954-3899/34/8/S50.
- (35) Merali, Zeeya (2011). "Collaborative physics: string theory finds a bench mate". Nature 478 (7369): 302–304. Bibcode:2011Natur.478..302M. doi:10.1038/478302a. PMID 22012369.
- (36) Moore, Gregory (2012). "Applications of the six-dimensional (2,0) theories to Physical Mathematics". Retrieved 14 August 2013.
- (37) Perlmutter, Saul (2003). "Supernovae, dark energy, and the accelerating universe". Physics Today 56 (4): 53–62. Bibcode:2003PhT....56d..53P. doi:10.1063/1.1580050.
- (38) Peskin, Michael; Schroeder, Daniel (1995). An Introduction to Quantum Field Theory. Westview Press. ISBN 978-0-201-50397-5.
- (39) Polyakov, Alexander (2008). "From Quarks to Strings". arXiv:0812.0183 [hep-th].
- (40) Rangamani, Mukund (2009). "Gravity and Hydrodynamics: Lectures on the fluid-gravity correspondence". Classical and quantum gravity 26 (22): 4003. arXiv:0905.4352. Bibcode:2009CQGra..26v4003R. doi:10.1088/0264-9381/26/22/224003.
- (41) Sachdev, Subir (2013). "Strange and stringy". Scientific American 308 (44): 44. Bibcode:2012SciAm.308a..44S. doi:10.1038/scientificamerican0113-44.
- (42) Jump up ^ Bekenstein, Jacob D. (January 1981). "Universal upper bound on the entropy-to-energy ratio for bounded systems". Physical Review D 23 (215): 287–298. Bibcode:1981PhRvD..23..287B. doi:10.1103/PhysRevD.23.287.

@@@@
 @@@@
 @@@@